

Journal of Visualized Experiments

In situ transverse rectus abdominis myocutaneous flap. A rat model of myocutaneous ischaemia reperfusion injury.

--Manuscript Draft--

Manuscript Number:	JoVE50473R2
Full Title:	In situ transverse rectus abdominis myocutaneous flap. A rat model of myocutaneous ischaemia reperfusion injury.
Article Type:	Methods Article - Author Produced Video
Corresponding Author:	Marie-Claire Edmunds, BSc (hons) MBChB (hons) MRCSEd University of Edinburgh Edinburgh, Midlothian UNITED KINGDOM
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	University of Edinburgh
Corresponding Author's Secondary Institution:	
First Author:	Marie-Claire Edmunds, BSc (hons) MBChB (hons) MRCSEd
First Author Secondary Information:	
Order of Authors:	Marie-Claire Edmunds, BSc (hons) MBChB (hons) MRCSEd Stephen J Wigmore, MBChB BSc PhD FRCS David Charles Kluth, MBChB BSc PhD FRCP
Order of Authors Secondary Information:	
Abstract:	<p>Free tissue transfer is the gold standard of reconstructive surgery to repair complex defects not amenable to local options or those requiring composite tissue. Ischemia reperfusion injury (IRI) is a known cause of free flap failure and has no effective treatment. Establishing a laboratory model of this injury can prove costly both financially as larger mammals are typically used and in the expertise required by the technical difficulty of these procedures, typically requiring an experienced microsurgeon. This publication and video demonstrate the effective use of a model of IRI in rats which does not require microsurgical expertise. This procedure is an in situ model of a transverse abdominis myocutaneous (TRAM) flap where atraumatic clamps are utilized to reproduce the ischemia-reperfusion injury. Laser Doppler Imaging (LDI) scanner is employed to assess perfusion in the flap and the image processing software Image J to assess percentage area skin survival as a primary outcome measure of injury.</p>

W2.29 Centre for Inflammation Research
Queen's Medical Research Institute
47 Little France Cres
Edinburgh
EH16 4TJ
Marie-claire.haddock@ed.ac.uk
03.01.13

Dear Dr L Jarzylo,

The revised manuscript and figures have now been uploaded onto the Jove site following your instructions.

Please do not hesitate to contact me should you need any further information.

Kind regards

Miss Marie-Claire Haddock BSc (hons) MBChB (hons) MRCSEd

***In situ* transverse rectus abdominis myocutaneous flap. A rat model of myocutaneous ischemia reperfusion injury**

Authors: Edmunds Marie-Claire,¹ Wigmore J Stephen² and Kluth David.³

Authors: institution(s)/affiliation(s) for each author:

All: MRC Centre for Inflammation Research, Queen's Medical Research Institute, 47 Little France Crescent, Edinburgh EH16 4TJ, UK

¹ MRC Clinical Research Fellow, Department of Surgery, Royal Infirmary of Edinburgh 51 Little France Crescent, Edinburgh, Midlothian EH16 4SA, UK

² Professor of Transplant Surgery and Honorary Consultant Surgeon, Department of Surgery, Royal Infirmary of Edinburgh 51 Little France Crescent, Edinburgh, Midlothian EH16 4SA

³ Senior Lecturer and Honorary Consultant Physician, Department of Nephrology, Royal Infirmary of Edinburgh 51 Little France Crescent, Edinburgh, Midlothian EH16 4SA, UK

Corresponding author: Marie-Claire Edmunds marie-claire.edmunds@ed.ac.uk, W2.29, Queen's Medical Research Institute. Centre for Inflammation Research, 47 Little France Crescent, Edinburgh, EH16 4TJ, UK. 00 44 1312426672

Keywords: Myocutaneous flap; preconditioning; ischemia reperfusion injury.

Short Abstract: (49/50)

Free tissue transfer is widely employed in reconstructive surgery to restore form and function following oncological resection and trauma. Preconditioning this tissue prior to surgery may improve outcome. This article describes an *in situ* transverse rectus abdominis myocutaneous flap (TRAM) in rats as a means for testing preconditioning strategies.

Long Abstract: (159/400)

Free tissue transfer is the gold standard of reconstructive surgery to repair complex defects not amenable to local options or those requiring composite tissue. Ischemia reperfusion injury (IRI) is a known cause of partial free flap failure and has no effective treatment. Establishing a laboratory model of this injury can prove costly both financially as larger mammals are conventionally used and in the expertise required by the technical difficulty of these procedures typically requires employing an experienced microsurgeon. This publication and video demonstrate the effective use of a model of IRI in rats which does not require microsurgical expertise. This procedure is an *in situ* model of a transverse abdominis myocutaneous (TRAM) flap where atraumatic clamps are utilized to reproduce the ischemia-reperfusion injury associated with this surgery. A laser Doppler Imaging (LDI) scanner is employed to assess flap perfusion and the image processing software, Image J to assess percentage area skin survival as a primary outcome measure of injury.

Introduction: 538

The goal of this protocol is to demonstrate a reliable and reproducible model of the ischemia-reperfusion injury observed in free tissue transfer to enable interventional strategies to be investigated.

Free tissue transfer is defined as the vascular detachment of an isolated block of tissue followed by autologous transplant of that tissue with anastomosis of the flap's transected

vessels to native vessels at the recipient site. The procedure is known as FTT and the tissue being transferred referred to as the free flap.

Free tissue transfer is the gold standard approach for the correction of complex, composite defects where local options are unsuitable or unavailable.¹⁻⁴ Ischemia reperfusion injury (IRI) is inevitable in free tissue transfer, contributes to flap failure^{5,6} and has no effective treatment. The elective nature of free flap surgeries permits administration of pharmacological agents to precondition against IRI.

IRI results in impaired flow through the microcirculation by endothelial activation and metabolic dysfunction,⁷ increased capillary permeability and subsequent interstitial edema⁷, influx of inflammatory cells,⁸ release of inflammatory mediators, reactive oxygen species⁹ and complement deposition.¹⁰ This complex process of hypoxia and subsequent reperfusion injury ultimately leads to cell death. A model of myocutaneous IRI enables the effectiveness of preconditioning strategies on clinical outcomes to be assessed. Recent work has validated the use of animal models of IRI studies as a surrogate for human IRI by comparing the molecular changes observed in human subjects and existing animal data.^{10,11}

The rat transverse rectus abdominis myocutaneous (TRAM) flap was first described in 1987 in German¹² and in 1993¹³ in English. This model gained wide popularity¹³⁻²⁵ as a cheap, robust model to investigate different strategies to reduce IRI associated with free tissue transfer.^{14,17-22} The majority of these studies were designed as unipedicled TRAM flaps based on the deep, inferior, epigastric vascular pedicle.^{15-18,20-22} Comparison of the data from these studies is complicated by the use of different sized cutaneous islands (10.5 - 30 cm²) and different lengths of postoperative follow-up (2 - 10 days). The average total percentage area flap necrosis in the control arm of these studies is 69 ± 6.2 % (*mean \pm SEM*). It should be noted that these six papers all employ the rectus abdominis muscle as a carrier for the vascular pedicle but do not expose, divide and microanastomose or clamp the vessels. Zhang *et al*²³ have described a true, free rat TRAM flap based on the superior epigastric vessels in which the flaps were raised, vessels divided and the myocutaneous flap transferred and microanastomosed to the groin vessels. This difficult technique required the microanastomosis of 0.45 - 0.5 mm caliber vessels. Only fifteen were performed and of these 67% survived.²³ The model described by Zhang *et al*²³ is an excellent model for the human free TRAM flap as it truly mirrors the injury incurred during FTT. The other published models of a rat TRAM flap more accurately reflect the injuries incurred during a human pedicled TRAM but do not accurately reflect the IRI as these flap in do not undergo an ischemic period followed by reperfusion as the vascular pedicle is never clamped or divided and microanastomosis performed. This protocol and video describe a new model of free tissue transfer using the rat TRAM in which the IRI is replicated using microclamps. This more faithfully replicates IRI than the pedicle TRAM predecessors but is technically easier than performing the microanastomosis. Microclamps have been widely employed by transplant researchers to recreate IRI associated with solid organ transplant;²⁶⁻³³ however, this is the first time it has been described in the rat TRAM flap.

PROTOCOL:

All surgery is performed in accordance with guidelines set out by the United Kingdom's Home Office and the University of Edinburgh's Veterinary Services Department

1. Surgical procedure set-up notes

1.1 Change into clean surgical scrubs, gown, scrub cap and mask. Clean all surfaces of the operating room including equipment with 2% chlorhexidine in 70% isopropyl alcohol.

1.2 Prior to surgery, autoclave all surgical supplies and instruments that will be used in the procedure. Sterile packs per operation should include: 3 drapes, gauze, cotton tip applicators, silicone sheeting and the surgical instruments, see the Table of Specific Surgical Materials and Tools and Figure 1. Weigh the rat and measure out the appropriate volume of buprenorphine (0.04mg/kg) to be administered subcutaneously 1h before completing the procedure. Lay out; 3 x 1 ml syringes for subcutaneous fluid administration during surgery, 2 x 6-0 Vicryl sutures, 1 x 5-0 Ethilon sutures, a sterile marker with ruler and a 10-blade disposable scalpel, 4 - 5 pairs of sterile gloves and a hand-held cautery unit.

1.3 Place 4 x 10 ml sterile, 0.9 % saline vials in a water bath warmed to 37°C. This will be used for subcutaneous fluid replacement (1ml/kg/h) and to rinse the surgical site. Set up the homeothermic blanket, rectal probe, heat lamp, operating microscope and anesthetic rig. Turn on the laser and its software, set-up another anesthetic rig and place a heat pad underneath the black mat on which the animal will lay during the scan.

1.4 Use male, Lewis rats weighing 250-300 g. House rats for 7 days with food and water *ad libitum* with 12 h light-dark cycles before any surgery is performed.

2. Anesthesia and skin preparation.

2.1 Place the rat in the anesthetic rig's induction chamber and administer 4% isoflurane with 1L/min O₂ for 2-3 min to induce anesthesia. Remove the anesthetized rat from the chamber and place it supine on the clean, heated mat. Maintain isoflurane at 1.5% using a nose cone. Apply lacrilube or similar agent to prevent corneal abrasion during the procedure. Perform a foot-pad pinch test to ensure the animal is adequately anesthetized before proceeding. Repeat this last test before each major step in the procedure and adjust the inhalational anesthetic concentration accordingly.

2.2 Closely shave the anterior abdominal using an electric shaver so that the entire abdominal surface is exposed. Apply depilating cream for the duration recommended by the supplier. Remove the cream and thoroughly rinse the skin with warmed sterile saline to remove all traces of the cream. Apply 2% chlorhexidine in 70% isopropyl alcohol to the skin and allow to dry before proceeding. This is the standard skin preparation in our unit based on current evidence for surgical site infection.³⁴ Please discuss with your Veterinary Department what is standard procedure in your unit before electing a skin preparation protocol.

2.3 Place 2 drapes either side of the rat and take care to keep them sterile. Put on sterile gloves and with the aid of an assistant open up the sterile packs. Place all the instruments on one drape and the sutures, gauze, cotton applicators, silicone sheeting and sterile marker pen with ruler on the other (Figure 2).

2.4 Identify the midline using xiphisternum and the tail as reference points. Mark the midline. Measure 0.8 cm below the xiphisternum and mark this point. Draw a line perpendicular to the midline from this point. Taking the midline as the center of the flap mark out 1 cm and 2 cm to the left and the right of the midline. Draw vertical lines parallel to the midline from the points. Measure 4 cm below the original horizontal line and draw another parallel to it. Following these instructions a 4 x 4 cm flap divided into 4 equal strips is delineated (see Figure 3).

3. Laser Doppler imaging.

3.1 Carefully move the rat to the nose cone of the second anesthetic rig at the LDI scanner. Continue anesthesia at 1.5% isoflurane, 1L/min O₂. Turn on the laser and follow the manufacturer's instructions to start scanning. After saving the scanned file return the rat back to the first rig and re-insert the rectal probe of the homeothermic blanket using soft white paraffin as a lubricant.

4. *In situ* TRAM flap - myocutaneous model of IRI

4.1 Re-scrub hands and put on fresh sterile gloves. Cut out a 5 cm diameter circle in the center of the remaining sterile drape and place this over the exposed abdomen to create a draped sterile-field.

4.2 Make an incision down the left lateral marked edge (Figure 4. A & B). Achieve hemostasis. Make similar incisions down the horizontal lines to the left of the midline. Achieve hemostasis.

4.3 The fat overlying the left inferior rectus sheath should be visible. Using forceps and fine iris scissors carefully get underneath this fat. Take care not to damage the perforators coming through the left, anterior rectus sheath. The plane opened up by such dissection is that immediately above the anterior abdominal wall fascia. Continue dissecting in this plane around the demarcated margins. In the left iliac fossa lies the large, superficial circumflex iliac vessels- these can be tied or cauterized. Then extend the dissection medially with caution and only as far as the lateral margin of the left rectus muscle. There is an obvious color change at this point from pink to near white (Figure 4. C). Carefully irrigate the area with sterile saline and check that hemostasis has been achieved before placing moist gauze over the area.

4.4 Repeat the procedure on the contralateral side but this time extend to the *linea alba* (mid line). Take care to identify and cauterize all the musculocutaneous perforators arising in the center of the right rectus abdominis muscle (Figure 4. D). If this is not done properly it can result in a postoperative hematoma and spurious results. Similarly achieve hemostasis, irrigate and place moist gauze over the raised flap.

4.5 Return to the inferior margin of the left anterior rectus (Figure 4. E & F). Cauterize the most inferior perforator seen. Proceed to cut a small (approx. 0.6 cm x 0.6 cm) window in the anterior rectus sheath using microscissors and pointed curved

Graeffe forceps. Blunt dissect slowly down the lateral margin of the muscle until the muscle thins out but before the posterior sheath is breached. Then rotate the forceps and blunt dissect medially until the belly of the muscle is atop the curved edge of the forceps and the tips are free at the medial margin. Feed approximately 6 cm of 5-0 Ethilon into the jaws of the forceps and tie off the inferior rectus sheath. Upon completion of this step the myocutaneous flap is isolated on one dominant vessel- the deep superior epigastric vessels. Cover with a moist gauze.

4.6 Cut the silicone sheeting into ovals with smooth corners, (Figure 4 G). These should be large enough to cover most of the area exposed under the fasciocutaneous portions of the flap. However, caution must be taken to ensure that the skin edge can be closed without any tension and the medial curve of the oval may have to be paired back to prevent it impairing flow through the remaining perforators. These are then sutured in place with 6-0 Vicryl (Figure 4 H). Cover with moist gauze.

4.7 Using simple interrupted 5-0 Ethilon sutures ‘peg out’ the flap to reduce heat and water loss (Figure 4 I). Cover with a moist gauze.

4.8 Extend the wound superiorly to the left of the xiphisternum (Figure 4 I). Suture this to the upper left quadrant to improve the field-of-view.

4.9 Cut away any overlying fat to reveal the superior, left, anterior rectus sheath. Cut a small (0.6 cm x 0.6 cm) window in this sheath (Figure 4 J). Extend the wound medially until a change in muscle fiber trajectory from vertical to oblique and consistency from tightly packed to loose fibrils is seen.

4.10 Insert the curved forceps carefully between these two muscles and open up a plane by blunt dissection. Carefully cut down only as far as the superior surface of these curved forceps cutting through the belly of the left rectus abdominis muscle to reveal the underlying deep, superior epigastric artery and vein (Figure 4 K).

4.11 Using micro-instruments and high power on the operating microscope, carefully separate the artery and vein and strip off any surrounding fat.

4.12 Apply atraumatic Acland clamps to the artery and vein (B-1, “V” type) and start the timer to count down the 30 min ischemic period. Irrigate the clamped pedicle and cover with gauze. We did not employ vessel dilators such as verapamil or pabavarine but should vessel spasm be a problem, these drugs should be considered.

4.13 Administer the buprenorphine (0.04 mg/kg) and warmed, sterile saline (1ml/kg/h).

4.14 Starting at the top left corner, suture the flap in place with 6-0 Vicryl subcuticular sutures stopping and tying off at the xiphisternum.

4.15 When the 30 min ischemic time is over, carefully remove the clamps and irrigate the pedicle with warmed saline. Check that flow has been re-established. Please note that this ischaemic time was stipulated by the UK Home Office authority. Researchers working in other authorities may be able to extend this time. Extending the ischemic time will likely lead to worse clinical outcome.

4.16 Suture the cut edges of the rectus back in place with 6-0 Vicryl. Take care not to apply too much tension as this can lead to kinking of the vessels.

4.17 Complete the subcutaneous suturing taking care to bury all knots below the skin (Figure 4 K).

4.18 Clean the wound area and allow it to dry. Re-draw the zones on the flap.

4.19 Re-scan the animal to obtain a postoperative image.

4.20 Re-apply lacrilube to the animal's eyes and place in a warmed incubator (37 °C) for 1 h to recover before returning to the husbandry unit.

Critical steps within the protocol

The crux of the procedure is in identifying the deep, superior epigastric vessels. This is shown clearly in the accompanying film. In brief, a window is cut in the anterior rectus sheath to expose the underlying muscle fibers which will be running longitudinally. By extending the superficial dissection of the anterior rectus sheath medially a change in muscle fiber trajectory is observed from longitudinal to oblique. Insert blunt ended, curved, Graeffe forceps (or similar) at the intersection of these two bundles of muscle fibers. Blunt dissect laterally. Cut down, using micro scissors, onto the upper surface of the curved forceps held in this plane between the muscle fiber bundles. On removal of the Graeffe forceps the deep, superior epigastric artery and vein will be observed at the midpoint of the rectus abdominis muscle body. Strip off the fat overlying the vessels using micro instruments and apply the clamps.

The fasciocutaneous portions of the rat TRAM flap are thin enough to permit the flap to take as a full thickness skin graft. To prevent this and to ensure that this is a true model of IRI a thin, flexible silicon sheet is placed underneath the fasciocutaneous portions of the flap.³⁵ This step has been adopted by other researchers undertaking rat TRAM models.^{17,21,25}

Rats chew through knots so make sure all sutures are subcuticular and all knots are buried. In carrying out meticulous suturing autocannibalism of flaps as reported by other researchers can be avoided.²⁴

Following administration of Buprenorphine reduce maintenance anesthesia to 1 % Isoflurane (1L/min O₂).

RESULTS

Rat models are more economical than larger animals models,³⁶ are disease-resistant in nature and may be genetically manipulated. Loose skinned animals, such as rodents, were thought to have a different arrangement of cutaneous blood supply compared to fixed skinned animals such as humans and pigs. In loose skinned animals, skin is supplied primarily by direct cutaneous blood vessels passing through the subcutaneous fat to the overlying skin (Figure 5) By contrast, fixed skinned animals derive cutaneous blood supply through vessels that course through the underlying muscles to supply the overlying integument via musculocutaneous perforators (Figure 5). Consequently there were concerns whether loose-skinned animals

could be used in flap research. However, Taylor's work on angiosomes showed that there were discreet areas of rat skin that are supplied in an analogous manner as humans via musculocutaneous perforators. The anterior abdominal wall on which the transverse rectus myocutaneous (TRAM) flap is based is one such area.^{37,36,15}

Relevant anatomy

The superior, deep, epigastric vessels are the dominant vascular pedicle in rats and six to ten perforators pass through the anterior rectus sheath to supply the overlying integument.^{13,15} The superior deep epigastric vessels in the rat enter the rectus abdominis muscle at the level of the xiphoid and continue, diminishing in caliber, toward the pubis. The lateral margins of the anterior abdomen are supplied by the superficial inferior and superior epigastric and circumflex iliac vessels.³⁷ There is physiological overlap between the territories supplied by these direct cutaneous branches and those areas of integument supplied by musculocutaneous perforators via choke vessels.³⁸ This is consistent with the anatomical and physiological vascular territories described in the human, though in the human the dominant vascular pedicle is the inferior rather than the superior epigastric artery.³⁹

Transverse rectus abdominis myocutaneous (TRAM) flap

The transverse rectus abdominis flap was first described for reconstruction following radical resection of breast cancer in 1974.⁴⁰ This myocutaneous flap is based on deep epigastric blood vessels and incorporates a portion of the rectus abdominis muscles and overlying integument. Throughout this article the TRAM flap will be divided into four equal areas called zones. They are numbered I-IV as per Schlegel *et al.* such that: Zone I (ZI) is the integument overlying the rectus abdominis muscle directly supplied by the vascular pedicle; Zone II (ZII) describes the integument overlying the contralateral rectus abdominis; Zone III (ZIII) the area lateral to Zone I; and Zone IV (ZIV) the area lateral to Zone II (see Figure 6).⁴¹

Laser Doppler imaging- assessing blood perfusion

Laser Doppler imaging provides a non-invasive means of assessing blood flow in the flap.⁴²⁻
⁴⁵ A monochromatic light source is emitted from the laser head. This incident light (blue in Figure 7) is shifted by erythrocytes within the tissue. The degree of shift is related to the velocity of the erythrocytes. The shifted light (green in Figure 7) is detected by the photo detector within the scanner head and converted into a measurement of perfusion. These are given in arbitrary units, perfusion units (PU), and the data converted into an image much like a weather map wherein perfusion is graded from high to low and each value assigned a color (Figure 2). The color map generated illustrates relative perfusion between the different areas of the flap. Each instrument is carefully calibrated such that comparisons can be made between subjects when the same scanner is employed

Rats underwent laser Doppler perfusion imaging using a Moor LD12 (Moor Instruments, Essex, UK) scanner pre-operatively, immediately postoperatively and at 24 and 48 h after surgery.

Using the software provided with the LDI scanner a region of interest (R.O.I) can be superimposed onto the LDI image and the average perfusion of that area calculated (Figure 2).

Image J analysis of percentage area necrosis- primary outcome measure

Image J is an open-access image processing program courtesy of the National Institutes of Health.⁴⁶ This can be used to measure areas and subsequently calculate the percentage skin area of each zone that is normal or fully necrosed at each time point (Figure 8).

Assessment of injury

The highest rates of skin necrosis were found in zone IV (see representative data in Figures 9 & 10) consistent with other studies.^{16,22,24,25,47} These findings correspond with the pattern of necrosis reported clinically in human TRAM flaps confirming that this is a faithful representation of the clinical problem.¹⁴ The total percentage area flap necrosis was 37.86 ± 5.4 % (*mean \pm SEM*).

Changes in skin blood

LDI perfusion scanning was employed to assess blood flow in the TRAM flap model. This is a simple, non-invasive and reproducible means of assessing perfusion (Figures 9 and 11). Perfusion decreased to 58.4 ± 0.49 % ($n = 10$, *mean \pm SEM*) immediately postoperatively, 56.98 ± 0.41 % at 24 h and 92.4 ± 0.6 % compared to the pre-operative values for the whole flap. The areas of the flap with lowest perfusion in the immediate postoperative and 24 h scans indicate areas where necrosis will subsequently develop at 48 h (see Figure 9).

Figure Captions:

Figure 1: [Equipment set-up]

The anesthetic rig with red induction chamber are seen behind desk. The rat is lying supine with anesthesia maintained via a nose cone. A heat lamp is employed to increase ambient temperature. Above the rat is the operating microscope. To the left of the rat is a sterile drape with gauze, sutures etc. To the right of the rat is a sterile drape with the surgical instruments. Core temperature is maintained using a homeothermic blanket (underneath the rat) and rectal probe attached to the Harvard apparatus device (black box in front of the sharps bin).

Figure 2: [Assessing average perfusion using LDI software]

Select the polygon icon from the tool bar (red arrow) then the region of interest (ROI) selection tool (rectangle with the blue cross, 2 icons to the right of the polygon tool). Using the mouse draw around the ROI, in this figure all 4 zones are marked. Before moving to the next ROI click on the rectangle with the blue square again. Once all the desired ROIs are selected press the stats icon in the center of the tool bar (the icon of a notepad with numbers on it) and the average perfusion statistics for each ROI will pop-up in a new window as shown.

Figure 3: [Marking out the flap boundaries and zones]

The depilated rat is placed supine. The midline is identified and marked (blue dashed line). A line is marked perpendicular to the midline 0.8 cm below the xiphisternum. 4 lines are drawn parallel to the midline, 1 cm apart. A last line is drawn parallel and 4 cm below the second line to complete the square.

Figure 4: [Step-by-step surgical approach]

The flap is delineated as previously described (4 A). The left lateral margin is incised (4 B) and dissection continued medially in the plane immediately superficial to the anterior abdominal wall fascia to the lateral margin of the left rectus (4 C). The same steps are conducted on the contralateral side but the dissection continued medially to the *linea alba* (midline), (4 D). Cauterize the musculocutaneous perforators arising from the center of the right rectus abdominis muscle. A small window is cut in the inferior aspect of left rectus sheath (4 E) and the inferior rectus tied off (4 F). Silicone sheeting is then cut and sutured in place underneath the fasciocutaneous portions of the flap (4 G & H). The flap is then

‘pegged’ out (4-I). Steps (4 G & I) can be performed before or after steps (4 E & F). A small window is cut in the superior aspect of the left, anterior rectus sheath (4-J). The exposed muscle is then carefully examined. A change in muscle fiber trajectory from parallel to oblique and tightly packed to loosely packed fibrils will be seen medially. Pass the curved Graeffe forceps between these muscle planes and blunt dissect laterally. Cut down onto the closed, upper surface of these forceps to expose the vascular pedicle. Remove surrounding fat and expose the vessels for clamping. Place Acland clamps on the artery and vein (4 K) and count down the ischemic time period. Start subcuticular suturing leaving the area immediately above the clamps until last. Remove the clamps after the allotted period and appose the free ends of the left rectus abdominis muscle. Complete the subcuticular sutures (4 L).

Figure 5: [Cutaneous blood supply in fixed and loose skinned animals]. Cutaneous blood supply in loose skinned mammals such as rats is predominantly via direct cutaneous branches rather than musculocutaneous perforators as in fixed skin mammals such as humans and pigs. For this reason rats have not historically been favored for plastic surgical research. This has been shown to be an out-dated notion and specific areas of the rat such as the anterior abdominal wall are supplied by musculocutaneous perforators and are therefore suitable areas to use for flap models.

Figure 6: [Zones of the transverse abdominis myocutaneous flap as described by *Schlefen et al.* in 1983]

The red arrow indicates the vascular pedicle (in this case the left, superior, deep epigastric vessels). The blue roman numerals show the 4 zones numbered I-IV based on their position relative to the vascular pedicle such that: Zone I (ZI) is the integument overlying the rectus abdominis muscle directly supplied by the vascular pedicle; Zone II (ZII) describes the integument overlying the contralateral rectus abdominis; Zone III (ZIII) the area lateral to Zone I; and Zone IV (ZIV) lateral to zone II.

Figure 7: [Laser Doppler imaging scanner]

Moor LD12 scanner assesses perfusion by sending out a monochromatic light sources which is shifted by the moving erythrocytes within the skin. The degree of shift is related to the velocity of the erythrocytes. This shifted light is detected by the photo-scanner and perfusion in that area calculated. A mirror then moves the beam in a sequential manner so that the whole anterior abdominal wall can be scanned in approximately 7 min.

Figure 8: [Image J analysis]

Figure 8-1) [Image J- Select straight-line tool] Select the straight-line tool, draw a line down the center of the flap as shown.

Figure 8-2. [Image J- Set scale 1]. Select Analyze from the tool bar and from the drop down menu select set scale.

Figure 8 -3. [Image J- Set scale 2]. On the pop up window set the scale to 4 cm.

Figure 8-4 [Image J- Select the polygon tool & outline zone of interest]. Select polygon tool (highlighted icon) and outline the zone of interest. The total perimeter of zone IV is outlined in this example.

Figure 8-5. [Image J- Measure area 1]. Select Analyze from the tool bar and on the drop down menu select Measure.

Figure 8-6. [Image J- Measure area 2]. The area will be displayed in a separate results window.

Figure 8-7. [Image J- Repeat for area of full necrosis] Repeat the previous 2 steps but this time only outline the necrosed area. This example shows the full necrosis in zone IV outlined. To calculate the percentage area full necrosis divide the latter value by the former and multiply by 100.

Figure 9 [Montage of representative images of this procedure.]

Caption: Each row represents a different subject. Photographs (left) and corresponding LDI image (right) are shown at the 4 different time points (from left to right: pre-operatively, postoperatively, at 24 h and at 48 h after surgery). It is clear that necrosis occurs consistently in zones ZIV and III. The color scale on the bottom right shows the colors and their corresponding perfusion units. Red- high perfusion, blue- low perfusion).

Figure 10: [Representative results- skin necrosis expressed as a percentage of total flap area at 48h.]

Caption: Percentage area full necrosis of the flap assessed clinically and measured using image J software at 48 h. The mean and SEM shown, $n = 10$

Figure 11: [Representative results- Laser Doppler imaging]

Caption: Laser Doppler imaging to show the average perfusion measured in perfusion units of the flap in control subjects pre-operatively, postoperatively, at 24 and 48h. The mean and SEM shown, $n = 10$

DISCUSSION

Modifications and trouble shooting

The protocol presented here reproduces the IRI seen in free tissue transfer in an experimental system enabling further understanding of that process and provides a means to investigate means of ameliorating IRI and improving outcome. This could easily be modified to produce a more severe injury if it were based on the non-dominant, deep, inferior epigastric pedicle or if the ischemic time were increased.

Limitations of technique

The anterior abdominal wall of the rat has significantly less subcutaneous fat than the anterior abdominal wall of most women undergoing TRAM flap surgery for breast reconstruction. The model outlined in this text is designed specifically as a model of ischemia reperfusion injury in myocutaneous flaps so that the effects of preconditioning therapies can be evaluated using skin necrosis and perfusion as outcome measures. The procedure detailed in this article does not specifically model problems such as fat necrosis which are incurred in human TRAM flaps when flaps with significant fat components are deliberately harvested to create projection for large breast reconstructions.

Direct *in-vivo* observation of the microcirculation is not demonstrated in this protocol but has been described in the cremaster muscle model⁴⁸ and in an osteomyocutaneous flap.^{7,49-51} The TRAM model is a myocutaneous flap model, if

researchers are particularly interested in osteomyocutaneous flaps this model is not appropriate but an alternative model has been described in the literature.⁵⁰

Significance with respect to other methods

Most published rat TRAM models use the rectus abdominis muscle surrounding the chosen vascular pedicle as a carrier for the vascular pedicle.^{13-22,24,25} They do not accurately reflect the IRI as the flap is never undergoes a true period of ischemia followed by reperfusion. Therefore, compared to these papers the model detailed in this protocol gives reproducible, controlled myocutaneous IRI. Researchers have also performed this as a free flap to the groin vessels²³ however this is technically extremely demanding as the deep superior, epigastric artery and vein measure 0.45 and 0.5 mm respectively. This protocol represents a simpler model.

Future applications

Research in improving outcome in free tissue transfer has focused primarily on preconditioning strategies. These strategies are employed or initiated before the surgery with the aim of 'training' the transferred tissue to withstand better the free tissue transfer surgery and this improve outcome. There are two main ways in which this can be achieved: pharmacological or ischemic preconditioning.⁵² A lot of this work has been performed on pigs which are more expensive to house and more difficult to work with than rats. The protocol described in this paper can be used to test these strategies in a laboratory animal which is easy to house and work with and in which there is possibility to work with genetically manipulated animals.

Acknowledgments:

This work was funded by the Medical Research Council grant G1000299.

The corresponding author would like to thank Gary Borthwick, University of Edinburgh, for assisting during the surgery.

The authors would like to acknowledge advice from Helen Douglas and Iain Mackay and allowing us to observe their Deep Inferior Epigastric (DIEP) flap procedure (Canniesburn Plastic Surgery Unit, Glasgow Royal Infirmary, 84 Castle Street, Glasgow G4 0SF, UK).

The authors would also like to thank Gary Blackie at the University of Edinburgh for his help in producing the video for this article.

Disclosures:


We have no disclosures.


References:

- 1 Wang, X. *et al.* Free anterolateral thigh adipofascial flap for hemifacial atrophy. *Ann. Plast. Surg.* **55** (6), 617-622 (2005).
- 2 Eckardt, A. & Fokas, K. Microsurgical reconstruction in the head and neck region: An 18-year experience with 500 consecutive cases. *J. Cranio. Maxill. Surg.* **31** (4), 197-201, doi:http://dx.doi.org/10.1016/S1010-5182(03)00039-8 (2003).
- 3 Yazar, S. *et al.* Safety and reliability of microsurgical free tissue transfers in paediatric head and neck reconstruction – a report of 72 cases. *J. Plast. Reconstr. Aes.* **61** (7), 767-771, doi:http://dx.doi.org/10.1016/j.bjps.2007.10.022 (2008).
- 4 Blondeel, P.N., Landuyt, K.H.V. & Monstrey, S.J. Surgical-technical aspects of the free diep flap for breast reconstruction. *Operat. Tech. Plast. Reconstr. Surg.* **6** (1), 27-37, doi:http://dx.doi.org/10.1016/S1071-0949(99)80017-1 (1999).
- 5 Siemonow, M. & Arslan, E. Ischaemia/reperfusion injury: A review in relation to free tissue transfers. *Microsurgery.* **24**, 468-475 (2004).
- 6 Wang, W.Z. Investigation of reperfusion injury and ischaemic preconditioning in microsurgery. *Microsurgery.* **29**, 72-79 (2009).
- 7 Rucker, M. *et al.* Reduction of inflammatory response in composite flap transfer by local stress conditioning-induced heat-shock protein 32. *Surgery.* **129** (3), 292-301, doi:10.1067/msy.2001.111079 (2001).
- 8 Cetinkale, O. *et al.* Involvement of neutrophils in ischemia-reperfusion injury of inguinal island skin flaps in rats. *Plast. Reconstr. Surg.* **102** (1), 153-160 (1998).
- 9 Korthuis, R.J., Granger, D.N., Townsley, M.I. & Taylor, A.E. The role of oxygen-derived free radicals in ischemia-induced increases in canine skeletal muscle vascular permeability. *Circ. Res.* **57** (4), 599-609 (1985).
- 10 Eisenhardt, S.U. *et al.* Monitoring molecular changes induced by ischemia/reperfusion in human free muscle flap tissue samples. *Ann. Plast. Surg.* **68** (2), 202-208, doi:10.1097/SAP.0b013e3181f77ba5 (2012).
- 11 Dragu, A. *et al.* Gene expression analysis of ischaemia and reperfusion in human microsurgical free muscle tissue transfer. *J. Cell. Mol. Med.* **15** (4), 983-993, doi:10.1111/j.1582-4934.2010.01061.x (2011).
- 12 Tilgner, A. & Herrberger, U. [myocutaneous flap models in the rat. Anatomy, histology and preparation technic of the myocutaneous rectus abdominis flap]. *Z. Versuchstierkd.* **29** (5-6), 231-236 (1987).
- 13 Dunn, R.M., Huff, W. & Mancoll, J. The rat rectus abdominis myocutaneous flap: A true myocutaneous flap model. *Ann. Plast. Surg.* **31** (4), 352-357 (1993).
- 14 Clugston, P.A., Perry, L.C., Fisher, J. & Maxwell, G.P. A rat transverse rectus abdominis musculocutaneous flap model: Effects of pharmacological manipulation. *Ann. Plast. Surg.* **34** (2), 154-161 (1995).
- 15 Ozgentas, H.E., Shenaq, S. & Spira, M. Development of a tram flap model in the rat and study of vascular dominance. *Plast. Reconstr. Surg.* **94** (7), 1012-1017; 1025-1016 discussion (1994).
- 16 Doncatto, L.F., da Silva, J.B., da Silva, V.D. & Martins, P.D. Cutaneous viability in a rat pedicled tram flap model. *Plast. Reconstr. Surg.* **119** (5), 1425-1430, doi:10.1097/01.prs.0000258534.46634.e7 (2007).
- 17 Lineaweaver, W.C. *et al.* Vascular endothelium growth factor, surgical delay, and skin flap survival. *Ann. Surg.* **239** (6), 866-873; discussion 873-865 (2004).
- 18 Rezende, F.C. *et al.* Electroporation of vascular endothelial growth factor gene in a unipedicle transverse rectus abdominis myocutaneous flap reduces necrosis. *Ann. Plast. Surg.* **64** (2), 242-246, doi:10.1097/SAP.0b013e318196cbe0 (2010).


- 19 Zacchigna, S. *et al.* Improved survival of ischemic cutaneous and musculocutaneous flaps after vascular endothelial growth factor gene transfer using adeno-associated virus vectors. *Am. J. Pathol.* **167** (4), 981-991, doi:10.1016/s0002-9440(10)61188-1 (2005).
- 20 Zhang, F. *et al.* Improvement of skin paddle survival by application of vascular endothelial growth factor in a rat tram flap model. *Ann. Plast. Surg.* **46**, 314-319 (2010).
- 21 Hijjawi, J. *et al.* Platelet-derived growth factor β , but not fibroblast growth factor 2, plasmid DNA improves survival of ischemic myocutaneous flaps. *Arch. Surg.* **139** (2), 142-147, doi:10.1001/archsurg.139.2.142 (2004).
- 22 Wong, M.S. *et al.* Basic fibroblast growth factor expression following surgical delay of rat transverse rectus abdominis myocutaneous flaps. *Plast. Reconstr. Surg.* **113** (7), 2030-2036 (2004).
- 23 Zhang, F. *et al.* Microvascular transfer of the rectus abdominis muscle and myocutaneous flap in rats. *Microsurgery.* **14** (6), 420-423 (1993).
- 24 Hallock, G.G. & Rice, D.C. Comparison of tram and diep flap physiology in a rat model. *Plast Reconstr Surg.* **114** (5), 1179-1184 (2004).
- 25 Qiao, Q. *et al.* Patterns of flap loss related to arterial and venous insufficiency in the rat pedicled tram flap. *Annals of Plastic Surgery August.* **43** (2), 171 (1999).
- 26 Persy, V.P., Verhulst, A., Ysebaert, D.K., De Greef, K.E. & De Broe, M.E. Reduced postischemic macrophage infiltration and interstitial fibrosis in osteopontin knockout mice. *Kidney Int.* **63** (2), 543-553, doi:10.1046/j.1523-1755.2003.00767.x (2003).
- 27 Li, Y. *et al.* Overexpression of cgmp-dependent protein kinase i (pkg-i) attenuates ischemia-reperfusion-induced kidney injury. *Am. J. Physiol. Ren. Physiol.* **302** (5), F561-570, doi:10.1152/ajprenal.00355.2011 (2012).
- 28 Hunter, J.P. *et al.* Effects of hydrogen sulphide in an experimental model of renal ischaemia-reperfusion injury. *Brit. J. Surg.* **99** (12), 1665-1671, doi:10.1002/bjs.8956 (2012).
- 29 Hamada, T., Fondevila, C., Busuttil, R.W. & Coito, A.J. Metalloproteinase-9 deficiency protects against hepatic ischemia/reperfusion injury. *Hepatology.* **47** (1), 186-198, doi:10.1002/hep.21922 (2008).
- 30 Duarte, S., Hamada, T., Kuriyama, N., Busuttil, R.W. & Coito, A.J. Timp-1 deficiency leads to lethal partial hepatic ischemia and reperfusion injury. *Hepatology.* **56** (3), 1074-1085, doi:10.1002/hep.25710 (2012).
- 31 Shen, X.D. *et al.* Cd154-cd40 t-cell costimulation pathway is required in the mechanism of hepatic ischemia/reperfusion injury, and its blockade facilitates and depends on heme oxygenase-1 mediated cytoprotection. *Transplantation.* **74** (3), 315-319 (2002).
- 32 Liu, J. *et al.* Endoplasmic reticulum stress modulates liver inflammatory immune response in the pathogenesis of liver ischemia and reperfusion injury. *Transplantation.* **94** (3), 211-217, doi:10.1097/TP.0b013e318259d38e (2012).
- 33 Pan, G.Z. *et al.* Bone marrow mesenchymal stem cells ameliorate hepatic ischemia/reperfusion injuries via inactivation of the mek/erk signaling pathway in rats. *J. Surg. Res.* **178** (2), 935-948, doi:10.1016/j.jss.2012.04.070 (2012).
- 34 Darouiche, R.O. *et al.* Chlorhexidine–alcohol versus povidone–iodine for surgical-site antisepsis. *New. Engl. J. Med.* **362** (1), 18-26, doi:10.1056/NEJMoa0810988 (2010).
- 35 Fukui, A., Inada, Y., Murata, K. & Tamai, S. "Plasmatic imbibition" in the rabbit flow-through venus flap, using horseradish peroxidase and fluorescein. *J Reconstr Miro surg.* **11**, 255-264 (1995).

- 36 Dunn, R.M. & Mancoll, J. Flap models in the rat: A review and reappraisal. *Plast. Reconstr. Surg.* **90** (2), 319-328 (1992).
- 37 Taylor, G. & Minabe, T. The angiosomes of the mammals and other vertebrates. *Plast. Reconstr. Surg.* **89** (2), 181-215 (1992).
- 38 Taylor, G., Corlett, R. & Boyd, J. The versatile deep inferior epigastric (inferior rectus abdominis) flap. *Brit. J. Plast. Surg.* **37** (3), 330-350 (1984).
- 39 Taylor, G., Corlett, R. & Boyd, J. The extended deep inferior epigastric flap: A clinical technique. *Plast. Reconstr. Surg.* **72** (6), 751-765 (1983).
- 40 Tai, Y. & Hasegawa, H. A transverse abdominal flap for reconstruction after radical operations for recurrent breast cancer. *Plast. Reconstr. Surg.* **53** (1), 52-54 (1974).
- 41 Schefflan, M. & Dinner, M.I. The transverse abdominal island flap: Part i. Indications, contraindications, results, and complications. *Ann. Plast. Surg.* **10**, 24-35 (1983).
- 42 Tindholdt, T.T., Saidian, S., Pripp, A.H. & Tonseth, K.A. Monitoring microcirculatory changes in the deep inferior epigastric artery perforator flap with laser doppler perfusion imaging. *Ann. Plast. Surg.* **67** (2), 139-142, doi:10.1097/SAP.0b013e3181f3e39b (2011).
- 43 Tindholdt, T.T., Saidian, S. & Tonseth, K.A. Microcirculatory evaluation of deep inferior epigastric artery perforator flaps with laser doppler perfusion imaging in breast reconstruction. *J. Plast. Surg. Hand. Surg.* **45** (3), 143-147, doi:10.3109/2000656x.2011.579721 (2011).
- 44 Booi, D.I., Debats, I.B.J.G., Boeckx, W.D. & van der Hulsi, R.R.W.J. A study of perfusion of the distal free-tram flap using laser doppler flowmetry. *J. Plast. Reconstr. Aes.* **61**, 282-288 (2008).
- 45 Hallock, G.G. Physiological studies using laser doppler flowmetry to compare blood flow to the zones of the free tram flap. *Ann. Plast. Surg.* **47** (3), 229-233 (2001).
- 46 Collin, T. Image j for microscopy. *Biotechniques. Suppl.* **43** (1), 25-30 (2007).
- 47 Hallock, G. & Rice, D. Physiologic superiority of the anatomic dominant pedicle of the tram flap in a rat model. *Plast. Reconstr. Surg.* **96**, 111-118 (1995).
- 48 Ozmen, S., Ayhan, S., Demir, Y., Siemionow, M. & Atabay, K. Impact of gradual blood flow increase on ischaemia-reperfusion injury in the rat cremaster microcirculation model. *J. Plast. Reconstr. Aes.* **61** (8), 939-948, doi:http://dx.doi.org/10.1016/j.bjps.2007.05.017 (2008).
- 49 Rucker, M., Vollmar, B., Roesken, F., Spitzer, W.J. & Menger, M.D. Microvascular transfer-related abrogation of capillary flow motion in critically reperfused composite flaps. *Brit J Plast Surg.* **55** (2), 129-135, doi:10.1054/bjps.2001.3748 (2002).
- 50 Rucker, M., Kadirogullari, B., Vollmar, B., Spitzer, W.J. & Menger, M.D. Improvement of nutritive perfusion after free tissue transfer by local heat shock-priming-induced preservation of capillary flowmotion. *J. Surg. Res.* **123**, 102-108 (2005).
- 51 Rucker, M. *et al.* New model for *in vivo* quantification of microvascular embolization, thrombus formation, and recanalization in composite flaps. *J. Surg. Res.* **108** (1), 129-137 (2002).
- 52 Wang, W.Z., Baynosa, R.C. & Zamboni, W.A. Update on ischemia-reperfusion injury for the plastic surgeon: 2011. *Plast. Reconstr. Surg.* **128** (6), 685e-692e, doi:10.1097/PRS.0b013e318230c57b (2011).






*Figure
[Click here to download high resolution image](#)



*Figure

[Click here to download high resolution image](#)


Image J-Select straight line tool



*Figure

[Click here to download high resolution image](#)


Image J- Set scale 1



*Figure

[Click here to download high resolution image](#)


Image J- Set scale 2



*Figure

[Click here to download high resolution image](#)


Image J- Select polygon tool & outline zone



*Figure

[Click here to download high resolution image](#)


Image J- Measure area-1



*Figure

[Click here to download high resolution image](#)


Image J- Measure area-2



*Figure

[Click here to download high resolution image](#)


Image J- Repeat for area of full necrosis





*Figure
[Click here to download high resolution image](#)




*Figure
[Click here to download high resolution image](#)









*Figure
[Click here to download high resolution image](#)



*Figure
[Click here to download high resolution image](#)



*Figure
[Click here to download high resolution image](#)



*Figure

[Click here to download high resolution image](#)



Animated/Video Figure

[Click here to download Animated/Video Figure: 50473jove-22-apr-13-MCE.m4v](#)

Name of Reagent/Material	Company
Moor LD12 laser doppler imaging scanner	http://gb.moor.co.uk/product/moorldi2-laser-doppler-imager/8
Complete homeothermic blanket system with flexible probe. Small. 230 VAC, 50 Hz.	www.harvardapparatus.com
Graeffe forceps 0.8 mm tips curved	http://www.finescience.de
Acland clamps	http://www.merciansurgical.com/acland-clamps.pdf
Clamp applicator	http://www.merciansurgical.com/acland-clamps.pdf
Gemini cautery unit	www.harvardapparatus.com
Micro-vessel dilators 11cm 0.3mm tips 00124	http://www.merciansurgical.com
Micro Jewellers Forceps 11cm angulated 00109	http://www.merciansurgical.com
Micro Jewellers Forceps 11cm straight 00108	http://www.merciansurgical.com
Acland Single Clamps B-1V (Pair)	http://www.merciansurgical.com
Micro Scissors Round Handles 15 cm Straight	http://www.merciansurgical.com
Iris Scissors 11.5cm Curves EASY-CUT	http://www.merciansurgical.com
Mayo Scissors 14cm Straight Chamfered Blades EASY-CUT	http://www.merciansurgical.com
Derf Needle Holders 12cm TC	http://www.merciansurgical.com
Ethilon 5-0	http://www.farlamedical.co.uk/
Vicryl rapide 6-0	http://www.millermedicalsupplies.com/
Instrapac - Adson Toothed Forceps (Extra Fine)	http://www.millermedicalsupplies.com/
Castroviejo needle holders	http://s-and-t.ne
Heat Lamp	http://www.chicken-house.co.uk
Silicone sheeting 0.3mm translucent	http://www.silex.co.uk/
Image J software	http://rsbweb.nih.gov/ij/

Zeiss OPMI pico Operating microscope	http://www.zeiss.co.uk/
Vet tech solution isofluorane rig	http://www.vet-tech.co.uk/
Vet tech solution isofluorane rig	http://www.vet-tech.co.uk/

Catalog Number	Comments
507221F	
11052-10	2
00398 V	B-1 'V' pattern clamps used on both
CAF-4	
726067	
D-5a.2	
JFA-5b	
JF-5	
396	
67	
EA7613-11	
EA7652-14	
703DE12	
W1618	
W9913	
7973	
12565-14	



17 Sellers Street
Cambridge, MA 02139
tel. +1.617.946.9061
www.jove.com

ARTICLE AND VIDEO LICENSE AGREEMENT

Title of Article:

INSITU TRANSVERSE RECTUS ABDOMINIS MYOCUTANEOUS FLAP
MYOCUTANEOUS ISCHAEMIA REPERFUSION INJURY.
HADDOK M-C, WIGMORE SJ, KLVTH D.

Author(s):

Item 1 (check one box): The Author elects to have the Materials be made available (as described at

<http://www.jove.com/publish>) via: ☒ Standard Access ☐ Open Access

Item 2 (check one box):

- ☒ The Author is NOT a United States government employee.
- ☐ The Author is a United States government employee and the Materials were prepared in the course of his or her duties as a United States government employee.
- ☐ The Author is a United States government employee but the Materials were NOT prepared in the course of his or her duties as a United States government employee.

ARTICLE AND VIDEO LICENSE AGREEMENT

1. **Defined Terms.** As used in this Article and Video License Agreement, the following terms shall have the following meanings: "Agreement" means this Article and Video License Agreement; "Article" means the article specified on the last page of this Agreement, including any associated materials such as texts, figures, tables, artwork, abstracts, or summaries contained therein; "Author" means the author who is a signatory to this Agreement; "Collective Work" means a work, such as a periodical issue, anthology or encyclopedia, in which the Materials in their entirety in unmodified form, along with a number of other contributions, constituting separate and independent works in themselves, are assembled into a collective whole; "CRC License" means the Creative Commons Attribution-Non Commercial-No Derivs 3.0 Unported Agreement, the terms and conditions of which can be found at: <http://creativecommons.org/licenses/by-nc-nd/3.0/legalcode>; "Derivative Work" means a work based upon the Materials or upon the Materials and other pre-existing works, such as a translation, musical arrangement, dramatization, fictionalization, motion picture version, sound recording, art reproduction, abridgment, condensation, or any other form in which the Materials may be recast, transformed, or adapted; "Institution" means the institution, listed on the last page of this Agreement, by which the Author was employed at the time of the creation of the Materials; "JoVE" means MyJoVE Corporation, a Massachusetts corporation and the publisher of *The Journal of Visualized Experiments*; "Materials" means the Article and / or the Video; "Parties" means the Author and JoVE; "Video" means any video(s) made by the Author, alone or in conjunction with any other parties, or by JoVE or its affiliates or agents, individually or in collaboration with the Author or any other parties, incorporating all or any portion of the Article, and in which the Author may or may not appear.

2. **Background.** The Author, who is the author of the Article, in order to ensure the dissemination and protection of the Article, desires to have the JoVE publish the Article and create and transmit videos based on the Article. In furtherance of such goals, the Parties desire to memorialize in this Agreement the respective rights of each Party in and to the Article and the Video.

3. **Grant of Rights in Article.** In consideration of JoVE agreeing to publish the Article, the Author hereby grants to JoVE, subject to Sections 4 and 7 below, the exclusive, royalty-free, perpetual (for the full term of copyright in the Article, including any extensions thereto) license (a) to publish, reproduce, distribute, display and store the Article in all forms, formats and media whether now known or hereafter developed (including without limitation in print, digital and electronic form) throughout the world, (b) to translate the Article into other languages, create adaptations, summaries or extracts of the Article or other Derivative Works (including, without limitation, the Video) or Collective Works based on all or any portion of the Article and exercise all of the rights set forth in (a) above in such translations, adaptations, summaries, extracts, Derivative Works or Collective Works and (c) to license others to do any or all of the above. The foregoing rights may be exercised in all media and formats, whether now known or hereafter devised, and include the right to make such modifications as are technically necessary to exercise the rights in other media and formats. If the "Open Access" box has been checked in Item 1 above, JoVE and the Author hereby grant to the public all such rights in the Article as provided in, but subject to all limitations and requirements set forth in, the CRC License.

4. **Retention of Rights in Article.** Notwithstanding the exclusive license granted to JoVE in Section 3 above, the

ARTICLE AND VIDEO LICENSE AGREEMENT

Author shall, with respect to the Article, retain the non-exclusive right to use all or part of the Article for the non-commercial purpose of giving lectures, presentations or teaching classes, and to post a copy of the Article on the Institution's website or the Author's personal website, in each case provided that a link to the Article on the JoVE website is provided and notice of JoVE's copyright in the Article is included. All non-copyright intellectual property rights in and to the Article, such as patent rights, shall remain with the Author.

5. Grant of Rights in Video – Standard Access. This Section 5 applies if the "Standard Access" box has been checked in Item 1 above or if no box has been checked in Item 1 above. In consideration of JoVE agreeing to produce, display or otherwise assist with the Video, the Author hereby acknowledges and agrees that, Subject to Section 7 below, JoVE is and shall be the sole and exclusive owner of all rights of any nature, including, without limitation, all copyrights, in and to the Video. To the extent that, by law, the Author is deemed, now or at any time in the future, to have any rights of any nature in or to the Video, the Author hereby disclaims all such rights and transfers all such rights to JoVE.

6. Grant of Rights in Video – Open Access. This Section 6 applies only if the "Open Access" box has been checked in Item 1 above. In consideration of JoVE agreeing to produce, display or otherwise assist with the Video, the Author hereby grants to JoVE, subject to Section 7 below, the exclusive, royalty-free, perpetual (for the full term of copyright in the Article, including any extensions thereto) license (a) to publish, reproduce, distribute, display and store the Video in all forms, formats and media whether now known or hereafter developed (including without limitation in print, digital and electronic form) throughout the world, (b) to translate the Video into other languages, create adaptations, summaries or extracts of the Video or other Derivative Works or Collective Works based on all or any portion of the Video and exercise all of the rights set forth in (a) above in such translations, adaptations, summaries, extracts, Derivative Works or Collective Works and (c) to license others to do any or all of the above. The foregoing rights may be exercised in all media and formats, whether now known or hereafter devised, and include the right to make such modifications as are technically necessary to exercise the rights in other media and formats. For any Video to which this Section 6 is applicable, JoVE and the Author hereby grant to the public all such rights in the Video as provided in, but subject to all limitations and requirements set forth in, the CRC License.

7. Government Employees. If the Author is a United States government employee and the Article was prepared in the course of his or her duties as a United States government employee, as indicated in Item 2 above, and any of the licenses or grants granted by the Author hereunder exceed the scope of the 17 U.S.C. 403, then the rights granted hereunder shall be limited to the maximum rights permitted under such statute. In such case, all provisions contained herein that are not in conflict with such statute shall remain in full force and effect, and all provisions contained herein that do so conflict

shall be deemed to be amended so as to provide to JoVE the maximum rights permissible within such statute.

8. Likeness, Privacy, Personality. The Author hereby grants JoVE the right to use the Author's name, voice, likeness, picture, photograph, image, biography and performance in any way, commercial or otherwise, in connection with the Materials and the sale, promotion and distribution thereof. The Author hereby waives any and all rights he or she may have, relating to his or her appearance in the Video or otherwise relating to the Materials, under all applicable privacy, likeness, personality or similar laws.

9. Author Warranties. The Author represents and warrants that the Article is original, that it has not been published, that the copyright interest is owned by the Author (or, if more than one author is listed at the beginning of this Agreement, by such authors collectively) and has not been assigned, licensed, or otherwise transferred to any other party. The Author represents and warrants that the author(s) listed at the top of this Agreement are the only authors of the Materials. If more than one author is listed at the top of this Agreement and if any such author has not entered into a separate Article and Video License Agreement with JoVE relating to the Materials, the Author represents and warrants that the Author has been authorized by each of the other such authors to execute this Agreement on his or her behalf and to bind him or her with respect to the terms of this Agreement as if each of them had been a party hereto as an Author. The Author warrants that the use, reproduction, distribution, public or private performance or display, and/or modification of all or any portion of the Materials does not and will not violate, infringe and/or misappropriate the patent, trademark, intellectual property or other rights of any third party. The Author represents and warrants that it has and will continue to comply with all government, institutional and other regulations, including, without limitation all institutional, laboratory, hospital, ethical, human and animal treatment, privacy, and all other rules, regulations, laws, procedures or guidelines, applicable to the Materials, and that all research involving human and animal subjects has been approved by the Author's relevant institutional review board.

10. JoVE Discretion. If the Author requests the assistance of JoVE in producing the Video in the Author's facility, the Author shall ensure that the presence of JoVE employees, agents or independent contractors is in accordance with the relevant regulations of the Author's institution. If more than one author is listed at the beginning of this Agreement, JoVE may, in its sole discretion, elect not take any action with respect to the Article until such time as it has received complete, executed Article and Video License Agreements from each such author. JoVE reserves the right, in its absolute and sole discretion and without giving any reason therefore, to accept or decline any work submitted to JoVE. JoVE and its employees, agents and independent contractors shall have full, unfettered access to the facilities of the Author or of the Author's institution as necessary to make the Video, whether actually published or not. JoVE has sole discretion as to the method of making and publishing the Materials, including,

ARTICLE AND VIDEO LICENSE AGREEMENT

without limitation, to all decisions regarding editing, lighting, filming, timing of publication, if any, length, quality, content and the like.

11. **Indemnification.** The Author agrees to indemnify JoVE and/or its successors and assigns from and against any and all claims, costs, and expenses, including attorney's fees, arising out of any breach of any warranty or other representations contained herein. The Author further agrees to indemnify and hold harmless JoVE from and against any and all claims, costs, and expenses, including attorney's fees, resulting from the breach by the Author of any representation or warranty contained herein or from allegations or instances of violation of intellectual property rights, damage to the Author's or the Author's institution's facilities, fraud, libel, defamation, research, equipment, experiments, property damage, personal injury, violations of institutional, laboratory, hospital, ethical, human and animal treatment, privacy or other rules, regulations, laws, procedures or guidelines, liabilities and other losses or damages related in any way to the submission of work to JoVE, making of videos by JoVE, or publication in JoVE or elsewhere by JoVE. The Author shall be responsible for, and shall hold JoVE harmless from, damages caused by lack of sterilization, lack of cleanliness or by contamination due to the making of a video by JoVE its employees, agents or independent contractors. All sterilization, cleanliness or decontamination procedures shall be solely the responsibility of the Author and shall be undertaken at the Author's expense. All indemnifications provided herein shall include JoVE's attorney's fees and costs related to said losses or

damages. Such indemnification and holding harmless shall include such losses or damages incurred by, or in connection with, acts or omissions of JoVE, its employees, agents or independent contractors.

12. **Fees.** To cover the cost incurred for publication, JoVE must receive payment before production and publication the Materials. Payment is due in 21 days of invoice. Should the Materials not be published due to an editorial or production decision, these funds will be returned to the Author. Withdrawal by the Author of any submitted Materials after final peer review approval will result in a US\$1,200 fee to cover pre-production expenses incurred by JoVE. If payment is not received by the completion of filming, production and publication of the Materials will be suspended until payment is received.

13. **Transfer, Governing Law.** This Agreement may be assigned by JoVE and shall inure to the benefits of any of JoVE's successors and assignees. This Agreement shall be governed and construed by the internal laws of the Commonwealth of Massachusetts without giving effect to any conflict of law provision thereunder. This Agreement may be executed in counterparts, each of which shall be deemed an original, but all of which together shall be deemed to be one and the same agreement. A signed copy of this Agreement delivered by facsimile, e-mail or other means of electronic transmission shall be deemed to have the same legal effect as delivery of an original signed copy of this Agreement.

A signed copy of this document must be sent with all new submissions. Only one Agreement required per submission.

AUTHOR:

Name:

MARIE-CLAIRE HADDOCK.

Department:

SURGERY

Institution:

UNIVERSITY OF EDINBURGH.

Article Title:

IN SITU TRANSVERSE RECTUS ABDOMINIS MYOCUTANEOUS FLAP. A RAT MODEL OF MYOCUTANEOUS ISCHAEMIA REPERFUSION INJURY.

Signature:

Mc

Date:

16. October 2012.

Please submit a signed and dated copy of this license by one of the following three methods:

- 1) Upload a scanned copy as a PDF to the JoVE submission site upon manuscript submission (preferred);
- 2) Fax the document to +1.866.381.2236; or
- 3) Mail the document to JoVE / Attn: JoVE Editorial / 17 Sellers St / Cambridge, MA 02139

For questions, please email editorial@jove.com or call +1.617.945.9051.

MS # (internal use):

W2.29

Queens' Medical Research Institute,
47 Little France Cres,
Edinburgh,
EH16 4TJ,
22nd April 2013.
marie-claire.edmunds@ed.ac.uk

Dear Miss Jarzylo,

Re: Peer Review comments for: JoVE50473R1.

I have addressed each of these comments in turn (black) under the original comment from the reviewer.

Veterinary Reviewer:

*0:59 - Indicates that animals are anesthetized but does not mention drug. Drugs are detailed in the PDF text.

Drug name now mentioned in new audio.

*1:08?- Absolutely not acceptable: "Chlorhexidine applied and allowed to dry?" It appears in both the text and video that the skin is only prepped once. It is standard practice to prep the skin 2-3 times. The current information provided insinuates that skin is only prepped once.

I have modified the video to include images of the skin preparation. This is how the skin was prepared. It is based on strong clinical evidence and is in-keeping current evidence-based guidelines in clinical practice and our local Department of Veterinary Services guidelines.¹⁻³ I have performed over 100 of these operations and have had no wound infections. I took this comment very seriously and performed an extensive literature search to find any evidence behind multiple chlorhexidine skin applications but was unable to find any evidence. I have subsequently amended the text to include the specific chlorhexidine-alcohol preparation that I used and have asked that future researchers clarify with their Veterinary Department what is standard practice in their unit and adopt that policy accordingly.

*5:23 - Improvement required: Use of the silicone sheeting. Please have the author add a statement the purpose of the silicone sheeting.

This is explained in the text:

“The fasciocutaneous portions of the rat TRAM flap are thin enough to permit the flap to take as a full thickness skin graft. To prevent this and to ensure that this is a true model of IRI a thin, flexible silicon sheet is placed underneath the fasciocutaneous portions of the flap.⁴ This step has been adopted by other researchers undertaking rat TRAM models.^{5-7”}

I have also added a statement to that effect in the audio accompanying the video.

*1.1 (TEXT) - Improvement required: "chlorhexidine in 70% alcohol". Please have author clarify this statement, is the chlorhexidine being mixed with alcohol or are both agents being used independently? Noticed there has been recent data to support the use of Chlorhexidine mixed in 70% alcohol.

I have changed this as requested.

*P 15-32 (TEXT) - Improvement required: The Figure #s are missing. Please add figure #s to each figure.

I believe this is a problem with the PDF created for peer review. I uploaded all Figures with their names in the correct order but this may not appear in the PDF. This is perhaps something that the technical team at JOVE could rectify.

Reviewers' comments:

Reviewer #1:

Major Concerns:

N/A

Minor Concerns:

These are very small calibre vessels - what Acland clamp pressure was used where these low pressure 3v and 3A clamps - was there a distinction between arterial and venous clamps used. Very often in microsurgical procedures before division and anastomosis of vessels and following anastomosis vessel dilators such as papavarin or indeed calcium channel blockers like Verapamil are used- was this the case after clamp release in your model? Further given that the vessel intima can easily be damaged by the use of these clamps were any vessels look at histologically or under the operating microscope to ascertain vessel intima damage? Why was the ischaemia reperfusion injury time 30 minutes given that in most microsurgical procedures the duration of ischaemia is 60 to 120 minutes - to re establish microsurgical flow in under 45 minutes in free tissue transfer is unusual. Finally given that the rat has a very low volume of anterior abdominal fat - will this model reflect the role of IRI in free tissue transfer for breast reconstruction given that Free TRAM flaps or pedicled or indeed DIEAP flaps contain large fat volumes required for breast reconstruction? -this rat model looks at the anterior abdominal skin and as an end point.

I have amended the reagents list to give more details of the clamps. B-1, "V"-type clamps were used on both the artery and vein. These are the smallest and gentlest Acland clamps available and are suitable for vessels 0.4-1 mm in diameter, they exert a pressure of between 5-15 gm/mm². We did not employ Verapamil or papavarine as we did not encounter prohibitive vasospasm but I thank the reviewer for the comment and have added it to the written protocol. After the clamps were removed re-establishment of arterial flow was confirmed by observing arterial flow over forceps. I was unable to get good video of this procedure otherwise I would have include it. The flaps were checked twice a day for any signs of venous congestion suggestive of venous thrombosis but none were found. I did assess a number of these vessels after

the animals were culled at 48 for signs of intimal damage. It was difficult to tell exactly where the clamps had been placed as they left no marks but no evidence of intimal damage was evident in those that I studied. The ischaemic time was limited to 30 minutes due to stipulations by the UK Home Office authority that we had to work under. I have added to the text that this can be extended to cause more significant injury. The rat does have a low fat volume compared to many women undergoing TRAM flaps but not all free flaps have thick subcutaneous fat depending on the site and the habitus of the patient. This is a model primarily of free tissue transfer rather than specifically free tissue transfer for breast reconstruction. My aim is that it will be used to evaluate methods for improving outcomes in myocutaneous free tissue transfer but I have gone on to do a study to assess a pharmacological preconditioning strategy in this model (soon to be submitted to *Am. J. Pathol.*). I have changed the text accordingly to indicate the limitations of this TRAM specifically as a model of breast reconstruction.

Additional Comments to Authors:

The video and the explanation of the model is very clear and will serve as excellent model

Reviewer #2:

Major Concerns:

Major concerns - written part:

* The authors are required to discuss "future applications" more in depth as this demonstrates the significance of the model.

This has been expanded.

* The authors used LDI perfusion scanning to assess blood flow in the TRAM flap however do not refer to former publications where LDI scan has been used.

These references have been added.

* In Figure 10 and 11, the authors provide representative results. The number of cases shown is low (n=3). A higher number and a statistical analysis would certainly strengthen the work.

I have changed these graphs (text and video) and text to reflect n =10.

Major concerns - video:

* As this is supposed to be a video publication, we expect at least some differentiated photographs instead of only one picture of the experimental setup in the first minute.

Please find the video amended accordingly.

* Anesthesia and skin preparation are well described in the written protocol however these steps are not shown in the video.

Please find the video amended accordingly.

* The authors provide a schematic drawing of flap dimensions. They do not demonstrate the steps in the animal though fundamental for a video publication.

Please find video amended accordingly.

* Magnification of the TRAM flap preparation should partially be reduced as the area of interest disappears out of sight.

Please accept my apologies but I am afraid this cannot be changed. I am no longer doing these surgeries and these are the best images that I have. If the JOVE video and manuscript review process, prior to peer review, had been shorter I could have taken more images. Equally, if the cost of their video assistance had been less I would have gladly used it.

* Ideally, the authors in person should briefly discuss indications and limitations of the model at the end of the video.

We are contractually bound by the University of Edinburgh to not appear in person linking us with live animal work nor to film any identifiable buildings. I cannot, therefore, comply with this request but have added some audio to the video file to briefly outline this area. This is due to genuine fear of safety of researchers doing work on animals in the UK from anti-vivisection campaigners. The University has had two attempted bombings on these grounds in the recent past.

* In summary non-microsurgical video sequences are essentially missing.

Please find updated images on the video.

Minor Concerns:

* Figure 1 looks rather confusing and unprofessional. Legends and arrows might help.

I have added arrows to this images and updated the sequence of images to take in other comments.

* The figures showing skin necrosis in the movie and the written part differ. Consistent graphs should be used.

Please accept my apology for this, I have updated these images.

* The first section of "critical steps" should rather be part of the protocol text than the discussion as surgical instructions are described.

This text has been transferred to the end of the Protocol section as requested.

* Explain "FTT". Free tissue transfer?

This has been added to the introduction.

* Write "Isoflurane" instead of "Isoflourane".

Corrected.

Additional Comments to Authors:
N/A

Yours sincerely,



Miss Marie-Claire Edmunds BSc (Hons) MBChB (Hons) MRCSEd

Bibliography

- 1 Darouiche, R.O. *et al.* Chlorhexidine–alcohol versus povidone–iodine for surgical-site antisepsis. *New. Engl. J. Med.* **362** (1), 18-26, doi:10.1056/NEJMoa0810988 (2010).
- 2 Macias, J.H. *et al.* Chlorhexidine is a better antiseptic than povidone iodine and sodium hypochlorite because of its substantive effect. *Am. J. Infect. Control.* doi:10.1016/j.ajic.2012.10.002 (2013).
- 3 Dumville, J.C., McFarlane, E., Edwards, P., Lipp, A. & Holmes, A. Preoperative skin antiseptics for preventing surgical wound infections after clean surgery. *Cochrane Database Syst Rev.* **3**, CD003949, doi:10.1002/14651858.CD003949.pub3 (2013).
- 4 Fukui, A., Inada, Y., Murata, K. & Tamai, S. "Plasmatic imbibition" in the rabbit flow-through venous flap, using horseradish peroxidase and fluorescein. *J Reconstr Microsurg.* **11**, 255-264 (1995).
- 5 Qiao, Q. *et al.* Patterns of flap loss related to arterial and venous insufficiency in the rat pedicled tram flap. *Annals of Plastic Surgery August.* **43** (2), 171 (1999).
- 6 Hijjawi, J. *et al.* Platelet-derived growth factor β , but not fibroblast growth factor 2, plasmid DNA improves survival of ischemic myocutaneous flaps. *Arch. Surg.* **139** (2), 142-147, doi:10.1001/archsurg.139.2.142 (2004).
- 7 Lineaweaver, W.C. *et al.* Vascular endothelium growth factor, surgical delay, and skin flap survival. *Ann. Surg.* **239** (6), 866-873; discussion 873-865 (2004).