

# Journal of Visualized Experiments

## Syntheses, crystallization and spectroscopic characterization of 3,5-lutidine N-oxide dehydrate.

--Manuscript Draft--

|  |  |
|--|--|
| <b>Article Type:</b>   | Invited Methods Article - JoVE Produced Video  |
| <b>Manuscript Number:</b>  | JoVE57233R3  |
| <b>Full Title:</b>   | Syntheses, crystallization and spectroscopic characterization of 3,5-lutidine N-oxide dehydrate. |
| <b>Keywords:</b>   | Crystallization, Lutidines, Spectroscopy, Basic Modulation, Supramolecular structures, Hydrates. |
| <b>Corresponding Author:</b>   | Yasmi Reyes Ortega<br>Benemerita Universidad Autonoma de Puebla<br>Puebla, Puebla MEXICO         |
| <b>Corresponding Author's Institution:</b>   | Benemerita Universidad Autonoma de Puebla  |
| <b>Corresponding Author E-Mail:</b>  | yasmi.reyes@correo.buap.mx   |
| <b>First Author:</b>   | Yasmi Reyes Ortega   |
| <b>Other Authors:</b>  | Samuel Hernández-Anzaldo<br>María del Rosario Merino-García                                      |
| <b>Author Comments:</b>  |  |
| <b>Additional Information:</b>   |  |
| <b>Question</b>  | <b>Response</b>  |
| If this article needs to be "in-press" by a certain date, please indicate the date below and explain in your cover letter. |  |

**TITLE:**

Syntheses, Crystallization and Spectroscopic Characterization of 3,5-Lutidine *N*-Oxide Dehydrate

**AUTHORS & AFFILIATIONS:**

María del Rosario Merino-García, Samuel Hernández-Anzaldo, Yasmi Reyes-Ortega  
Centro de Química, Instituto de Ciencias, Benemérita Universidad Autónoma de Puebla, Puebla, México.

**CORRESPONDENCE AUTHOR:**

Yasmi Reyes-Ortega

**EMAIL ADDRESSES:**

Yasmi Reyes-Ortega (yasmi.reyes@correo.buap.mx)  
Samuel Hernández-Anzaldo (samuel.hernandezan@correo.buap.mx)  
María del Rosario Merino-García (mar\_aura78@outlook.com)

**KEYWORDS:**

Crystallization, lutidines, spectroscopy, basic modulation, supramolecular structures, hydrates

**SUMMARY:**

Herein, we report the synthesis and crystallization of 3,5-lutidine *N*-oxide dehydrate by a simple protocol that differs from the classical synthesis of pyridine *N*-oxide. This protocol utilizes different starting material and involves less reaction time to yield a new solvated supramolecular structure, which crystallizes under slow evaporation.

**ABSTRACT:**

The synthesis of 3,5-lutidine *N*-oxide dehydrate, **1**, was achieved in the synthesis route of 2-amino-pyridine-3,5-dicarboxylic acid. Ochiai first used the methodology for non-substituted pyridines in 1957 in a 12 h process, but no X-ray suitable crystals were obtained. The substituted ring used in the methodology presented here clearly influenced the addition of water molecules into the asymmetric unit, which confers a different nucleophilic strength in **1**. The X-ray suitable crystal compound **1** was possible due to the stabilization of the negative charge in the oxygen by the presence of two water molecules where the hydrogen atoms donate positive charge into the ring; such water molecules serve well to construct a supramolecular interaction. The hydrated molecules may be possible for the alkaline system that is reached by adjusting the pH to 10. Importantly, the double methyl substituted ring and a reaction time of 5 h, makes it a more versatile method and with wider chemical applications for future ring insertions.

**INTRODUCTION:**

Nowadays, scientists around the globe have been investing resources into the development of new synthetic routes for the functionalization of aromatic groups, which are known for low reactivity front to addition reactions<sup>1-3</sup>. Pyridine, where a nitrogen atom substitutes a carbon atom, presents a similar chemical reactivity to analogue rings composed solely of carbon atoms<sup>3</sup>, and it usually undergoes a substitution mechanism rather than addition. *N*-oxides are distinctive

by the presence of a donor bond between nitrogen and oxygen formed by the overlap of the nonbonding electron pair on the nitrogen with an empty orbital on the oxygen atom<sup>3</sup>. Particularly, pyridine *N*-oxides are Lewis bases, because their N-O moiety may act as an electron donor, and they may combine with Lewis acids forming the corresponding Lewis acid-base pairs. This property has an essential chemical consequence, because it can increase the nucleophilicity of the Lewis acids towards potential electrophiles and thus allow them to react under conditions where normally the reaction would not occur. Probably the most frequent use of such compounds is in various oxidation reactions where they act as oxidants<sup>4</sup>. Pyridine *N*-oxides and many of their ring-functionalized derivatives are recurrent molecules of biologically active and pharmacological agents<sup>5</sup>, and a clear spatial distribution by different spectroscopic tools has been established for some of them<sup>6,7</sup>. In research on attaching different groups to the pyridine ring, scientists have tested various methodologies to produce an easy and conventional method, since isoxazolines requires a catalytic amount of base such as DBU in boiling xylene to form 6-substituted-2-aminopyridine *N*-oxides<sup>8,9</sup>. A variety of pyridine derivatives were converted into their corresponding *N*-oxides in the presence of a catalytic amount of manganese tetrakis(2,6-dichlorophenyl)porphyrin and ammonium acetate in CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>CN<sup>8,10</sup>. Other pyridines are oxidized to their oxides using H<sub>2</sub>O<sub>2</sub> in the presence of catalytic amounts of methyltrioxorhenium<sup>8,11</sup>, or by the addition of excess dimethyldioxirane in CH<sub>2</sub>Cl<sub>2</sub> at 0 °C, which leads to the corresponding *N*-oxides<sup>8,12-14</sup>. Bis(trimethylsilyl)peroxide in the presence of trioxorhenium in CH<sub>2</sub>Cl<sub>2</sub> has been used for the synthesis of pyridine *N*-oxides<sup>8,11</sup>. The synthesis of aminopyridine *N*-oxides involving acylation using Caro's acid (peroxomonosulfuric acid) has also been reported<sup>8</sup>. Nevertheless, the methodology reported here, and which uses part of the methodology reported by Ochiai<sup>1</sup>, provides very good results with the use of cheaper and accessible reagents, H<sub>2</sub>O<sub>2</sub> and glacial acetic acid. This practice is more suitable for use in large scale preparations that act on tertiary amines, it produces good yields in a reaction that only requires 30% hydrogen peroxide and glacial acetic acid in a temperature between 70–80 °C, and it uses a purification process that is available in most synthesis laboratories like distillation, without the use of catalyst or more expensive reagents<sup>1</sup>. The literature reports that other methodologies also frequently involve time frames from 10–24 h and temperatures above 100 °C<sup>4,8</sup>, and the yield of well-formed crystals for X-ray analyses is rarely reported.

Reactively, various *N*-oxide derivatives are used to adequately activate the lutidine ring, in either a nucleophilic or electrophilic way. The nucleophilic or electrophilic factor is affected by the substituents. With the pyridine ring being the electron-withdrawing groups, the main factor is the nucleophilic characteristic<sup>1</sup>. The free *N*-oxide compounds are rarely isolated as suitable crystals for X-ray analysis due to the delocalized charge in the aromatic ring. However, the solvation factor is critical to stabilize the negative density of the oxygen<sup>15</sup>.

## PROTOCOL:

### 1. Reaction

1.1. Place in a fume hood an opened round 100 mL flask with 0.5 mol (29.8 mL) glacial acetic acid and add 0.051 mol (5.82 mL) of 3,5-dimethylpyridine and 5 mL of H<sub>2</sub>O<sub>2</sub> (35%). Keep the

mixture reaction under constant magnetic stirring, at an inner temperature of 80 °C for 5 h.

1.2. After the reaction time, cool the flask to 24 °C with ice (do not expose the acetic acid gases to the ice), and plug it to a high vacuum distillation unit for 90–120 min to remove excess acetic acid.

Caution: Do not use hot material. Wait until the glassware reaches a manageable temperature. This will also avoid vapors entering the top of the distillation unit.

1.3. Add distilled water (10 mL) twice to ensure the removal of any trace of acetic acid and to concentrate the mixture much as possible.

## **2. Basicity Adjustment and Extraction**

2.1. Dissolve in bi-distilled water the isolated viscous and transparent product and use a potentiometer to adjust the pH to 10 with pure solid Na<sub>2</sub>CO<sub>3</sub>.

2.2. Place carefully the solution in a 250 mL separation funnel and extract it 5 times with 250 mL of CHCl<sub>3</sub> to improve the yield. Recover the organic layer and dry it over solid Na<sub>2</sub>SO<sub>4</sub> for 30 min maximum, which will contain the product. If necessary, re-extract the aqueous phase with the desired amount of CHCl<sub>3</sub>.

Caution: CHCl<sub>3</sub> may cause drowsiness and dizziness; handle with care and inside a fume hood.

2.3. Remove the solvent under reduced pressure with a high vacuum distillation unit, until the formation of a very hygroscopic clear beige crystalline powder (70%).

## **3. Crystallization Process**

3.1. Dissolve 4.3 g of the crystalline powder in 50 mL of cold high performed liquid chromatography (HPLC) grade diethyl ether. Vacuum filter the solution to remove any trace of solid starting material or even dust. Pour the filtrate into a glass Petri dish, leaving it to slow evaporate at 4 °C in a laboratory fridge.

3.2. Ensure that after two days, clear colorless crystals are obtained. Then measure the melting point, which should be in the range of 310–311 K.

## **4. Analysis of 3,5-Lutidine *N*-oxide Dehydrate**

4.1. Remove the crystals that are formed, of prismatic shape and colorless, by decantation from the flask's walls for further X-ray analysis. If not immediately used, keep the crystals in diethyl ether to avoid crystal hydration.

4.2. Dissolve 0.010 g of 3,5-lutidine *N*-oxide dehydrate in 0.4 mL of CDCl<sub>3</sub> to perform NMR H<sup>1</sup>

and  $C^{13}$  analysis to prove the effectiveness of the procedure.

#### REPRESENTATIVE RESULTS:

The protocol is essentially an extension of Ochiai's technique<sup>1</sup>. However, lower temperature and less time are applied. This simple method can be used to obtain a versatile ligand, which is a substituted pyridine *N*-oxide derivate. To confirm the formation of **1**, NMR  $^1H$  and  $^{13}C$  analysis are preferred to test the effectiveness of the procedure.

The chemical shift demonstrates the formation of **1**. The signal at 2.28 ppm (parts per million) corresponds to the six equivalent hydrogens of the two methyl groups in the 3 and 5 positions, which perceive the magnetic field in less proportion than the permanent magnetic. There are two sets of septuplets: one belongs to the proton in the c position at 7.9, which doubles the size of the other signal at 6.9 that belongs to the proton in the position a. **Figure 1** shows the chemical shifts provoked by the presence of the oxygen atom bonded to the nitrogen atom of the pyridine ring. The oxygen atom is electro-withdrawing and the closer hydrogen atoms to the oxygen atom (c and a) show displacement to a higher frequency than that for the methyl hydrogens (b).

The same process is plotted for the NMR  $^{13}C$  spectrum, **Figure 2**, where the signals for the closer carbons to the oxygen atom (c and a) show frequency separation between their signals of  $\Delta_c = 1,300$  Hz and  $\Delta_a = 200$  Hz. Once again, the methyl carbons do not show any change. The IR spectrum can be used to see the success of the method as well.

The ORTEP diagram, **Figure 4**, demonstrates the presence of two molecules of water surrounding the asymmetric molecule. These molecules are believed to stabilize the N—O bond. In similar cases, it has been described for pyridine *N*-oxide and related aromatic oxides. There is a significant stabilizing  $\pi$ -type  $O \rightarrow N$  back-donation, reflected in a calculated bond order higher than 1 and a number of electron lone pairs on the O atom lower than 3<sup>6</sup>.

#### FIGURE AND TABLE LEGENDS:

**Figure 1. TMS referenced  $CDCl_3$  500 MHz NMR  $^1H$  spectrum of **1**.** The integrations and chemical shifts of the three signals agree with three different types of hydrogen atoms present in lutine *N*-oxide.

**Figure 2. TMS referenced  $CDCl_3$  100 MHz NMR  $^{13}C$  spectrum of **1**.** Three signals are observed for the five aromatic carbons and one for the two methyl groups.

**Figure 3. IR spectrum of **1**.** The O-H bonds, above of  $3,300\text{ cm}^{-1}$ , are responsible for the supramolecular structure formation and the crystal formation.

**Figure 4. ORTEP diagram of **1** where two molecules of  $H_2O$  forms bridge hydrogen bonds with lutidine's oxygen, driving their hydrogen atoms toward the oxygen atom.** This Figure has been modified from Merino García *et al.*<sup>12</sup>

**Figure 5. Pictures of suitable X-ray diffraction crystals of **1** in diethyl ether (top) and at open air**

**(bottom).** One of these crystals was confirmed in the X-ray magnetometer and showed a diffraction path of X-ray, which was translated and refined by special computational programs in a molecular and crystalline structure<sup>24-28</sup>.

## DISCUSSION:

The protocol presented here is a conventional method to link an oxygen atom to the nitrogen atom of the 3,5-lutidine as a functionalization method of substrates. This technique is also well established to yield X-ray suitable dehydrated crystals (**Figure 5**, pictures taken with a DSC-HX300 Cyber-shot Sony camera). As far as we are concerned, not many reports have described the production of such crystals<sup>16</sup>. Many compounds grow ideal crystals for X-ray analysis when they are chelated by various metals<sup>17-20</sup>. Once the crystalline powder is formed, it is important to extract it from their mother liquors using a Kitasato flask and a Buchner funnel. Using rubber hoses, the Kitasato flask is connected to a vacuum line and on top of it the Buchner funnel is placed with a filter paper. Once the vacuum has been turned on, the filter paper is moistened with a small amount of solvent from which the product crystallized. This prevents the crystalline powder from trickling into the Buchner funnel by the vacuum effect. After securing the filter paper, the solution containing the crystalline powder is shaken to ensure that all the crystalline powder is filtered, and none remains in the bottom of the flask. The solution is quickly poured over the Buchner funnel. The crystalline powder obtained is left for about 10 min on the filter paper, and then the vacuum is turned off and the crystalline powder is detached from the paper and stored in an opaque glass vial, labeled with its code and kept at 4 °C until further analysis. The filtrated liquid is poured into a glass Petri dish, leaving it to slowly evaporation at 4 °C to improve the formation of adequate crystals for X-ray analysis.

It is important to notice that this protocol uses solvents and materials that are easily obtainable and generally are found at any research laboratory. The pH adjustment by the addition of Na<sub>2</sub>CO<sub>3</sub> and the consistent magnetic stirring are critical to the yield of the final product. However, it is important to pay extra careful attention in all process steps, especially in the extraction stage where no trace of starting material must be present to afford the formation of crystalline powder and subsequently crystals. Thus, this extraction/purification stage can be monitored by either NMR or IR spectroscopy to ensure the quality of the product.

To ensure the reproducibility of this protocol, NMR is an excellent tool. Even fine details are visible in the spectrum. All signals are shown as insets in **Figure 1**. These insets depict clearly a split, namely multiplicity, of all signals. For instance, the protons b ( $J_{\text{dip}}^b \sim 0.75 \text{ Hz}$ ) show four peaks at the crest of the signal, with a separation among them ( $\Delta_{\text{peak-peak}}$ ) queasy constant of  $\sim 0.0075 \text{ ppm}$ . The  $0.0075 \text{ ppm}$  may be transformed to energy using the following equation<sup>21</sup>:

$$E = \frac{(H_{\text{magnetometer}})(\Delta_{\text{peak-peak}})}{10^6} \quad \text{Equation 1}$$

$$E(\text{Hz}) = \frac{(500 \times 10^6 \text{ Hz})(0.0075 \text{ ppm})}{10^6 \text{ ppm}} = 0.75 \text{ Hz}$$

The transformation is recommended because the signals unfolding come from the dipolar spatial interaction among the three hydrogens nuclei of the methyl group, and even though they are further than the 4 single bonds with protons c and a, they are able to perceive their dipolar magnetic momentum interactions<sup>22</sup>. Additionally, the free sigma bonding rotation in the methyl group allows the super hyperfine proton-proton interaction to be visible in the multiplicity of the signal. The septuplets of protons a and c at 6.9 and 7.9 ppm, respectively are derived from the same dipolar nature phenomenon. In these cases, protons a and c can differentiate the protons in the methyl group for the same rotation dynamic. Last, as expected, the calculated  $J_{\text{dip}}$  for a, b, and c have barely the same value,  $\sim 0.75$  Hz. These quantities of the interactions confirm the hydrogen nuclei spatial arrangement throughout the magnetic anisotropy.

On the other hand, the  $C_{2v}$  symmetry of **1** makes equivalent carbons<sup>23</sup>. The  $^{13}\text{C}$  spectrum, **Figure 2**, shows the typical signal for methyl groups attached to aromatic rings, carbons d at 18 ppm. Moreover, a signal at 129 ppm is visible at this region due to the less electronegative element influenced carbon a. At high frequencies the signal for the more exposed carbons nuclei to the magnetic field are presented at 137 ppm<sup>22</sup>.

The presented methodology is very useful for the synthesis of pyridine N-oxides, providing good yields, in a reasonable time with soft reaction conditions and cheap and easy accessible reagents, that do not require additional catalysts. These conditions can be used for the scientific and educational community to obtain a broad range of pyridines N-oxides as precursors for other molecules of interest. The suitable methodology gives the opportunity to acquire basic experimental and conceptual tools in educational laboratories for students, proving a successful synthesis of compounds and the happiness to see the formation of crystals. However, it is important to emphasize that, like any chemical reaction, it is necessary to take all precautions since generally the reagents used are dangerous.

#### ACKNOWLEDGMENTS:

The present work has been supported by Vicerrectoría de Investigación y Estudios de Posgrado from BUAP, Divulgarion of Science, and Projects No. REOY-NAT14, 15, 16-G. HEAS-NAT17. RMG thanks CONACyT (Mexico) for scholarship 417887.

#### DISCLOSURES:

All authors declare no conflict of interest.


#### REFERENCES:


1. Ochiai, E. Recent Japanese work on the chemistry of pyridine 1-oxide and related compounds. *J. Org. Chem.* **18** (5), 534–551, doi: 10.1021/jo01133a010 (1953).
2. Solomons, T.W.G. Organic Chemistry. John Wiley & Sons. 2<sup>nd</sup> Edition, 1110 pp (1976).
3. Albin, A., Pietra, S. Heterocyclic N-Oxides. CRC Press. 328 pp, ISBN: 0849345529 (1991).
4. Koukal, P., Ulc, J., Necas, D., Kotora. Heterocyclic N-Oxides. *Topics in Heterocyclic Chemistry*. **53**, 29-58, doi: 10.1007/7081\_2017\_3 (2017).
5. Wen-Man, Z., Jian-Jun, D., Xu, J., Jun X., Huan-Jian, X. Visible-Light-Induced C2 alkylation of pyridine N- oxides. *J. Org. Chem.* **82** (4), 2059–2066, doi: 10.1021/acs.joc.6b02891, (2017).


6. Merino García, M. R., Ríos-Merino, F.J., Bernès, S., Reyes-Ortega, Y. Crystal structure of 3,5-di-methyl-pyridine *N*-oxide dihydrate. *Acta Cryst.* **E72** (12), 1687-1690, doi: 10.1107/S205698901601687X, (2016).
7. Sarma, R., Karmakar, A., Baruah, J. B. *N*-Oxides in Metal-Containing Multicomponent Molecular Complexes. *Inorg. Chem.* **47** (3), 763-765, doi: 10.1021/ic701639n (2008).
8. Youssif, S. Recent trends in the chemistry of pyridine *N*-oxides. *ARKIVOC.* **2001** (I), 242-268, ISSN 1424-6376 (2001).
9. Chucholowski, A. W., Uhlenndorf, S. Base catalyzed rearrangement of 5-cyanomethyl-2-isoxazolines; novel pathway for the formation of 2-aminopyridine *N*-oxides. *Tetrahedron Lett.* **31** (14), 1949-1952, doi: 10.1016/S0040-4039(00)88886-3, (1990).
10. Thellend, A, Battioni, P., Sanderson, W., Mansuy, D. Oxidation of *N*-Heterocycles by H<sub>2</sub>O<sub>2</sub> Catalyzed by a Mn-Porphyrin: An Easy Access to *N*-Oxides Under Mild Conditions. *Synthesis*, **1997** (12), 1387-1388, doi: 10.1055/s-1997-1383 (1997).
11. Copéret, C., Adolfson, H., Tinh-Alfredo, V. Kh., Yudin, A.K., Sharpless, K.B. A simple and Efficient Method for the Preparation of Pyridine *N*-Oxides. *J. Org. Chem.* **63** (5), 1740-1741, doi: 10.1021/jo9723467 (1998).
12. Ferrer, M., Sánchez-Baeza, F., Messeguer, A. On the preparation of amine *N*-oxides by using dioxiranes. *Tetrahedron*, **53** (46), 15877-15888, doi: 10.1016/S0040-4020(97)10048-5 (1997).
13. Adam, W., Briviba, K., Duschek, F., Golsch, D., Kiefer, W., Sies, H. Formation of singlet oxygen in the deoxygenation of heteroarene *N*-oxides by dimethyldioxirane. *J. Chem. Soc. Chem. Commun.* **1995** (18), 1831-1832, doi: 10.1039/C39950001831 (1995).
14. Murray, R. W.; Singh, M. A Facile One-Step Synthesis of C-Arylnitrones Using Dimethyldioxirane. *J. Org. Chem.*, **55** (9), 2954-2957 doi: 10.1021/jo00296a073 (1990).
15. Kim, S. W., Um, T., Shin, S. Brønsted acid-catalyzed  $\alpha$ -halogenation of ynamides from halogenated solvents and pyridine-*N*-oxides. *Chem. Commun.* **53** (18), 2733-2736, doi: 10.1039/C6CC10286G (2017).
16. Campeau, L., Rousseaux, R., Fagnou K. A solution to the 2-pyridyl organometallic cross-coupling problem: regioselective catalytic direct arylation of pyridine *N*-oxides. *J. Am. Chem. Soc.* **127** (51), 18020-18021, doi: 10.1021/ja056800x (2005).
17. Gang, L., et al. Metal-free methylation of a pyridine *N*-oxide C-H bond by using peroxides. *Org. Biomol. Chem.* **13** (46), 11184-11188, doi: 10.1039/C5OB01900A (2015).
18. May, D., Nyman, Hampden-Smith, M.J., Duesler E. N. Synthesis, characterization, and reactivity of group 12 metal thiocarboxylates, M(SOCR)<sub>2</sub>Lut<sub>2</sub> [M ) Cd, Zn; R ) CH<sub>3</sub>, C(CH<sub>3</sub>)<sub>3</sub>; Lut ) 3,5-Dimethylpyridine (Lutidine)]. *Inorg. Chem.* **36** (10), 2218-2224, doi: 10.1021/ic960784a (1997).
19. Cho, S.H., Hwang, S.J., Chang, S. Palladium-Catalyzed C-H Functionalization of Pyridine *N*-Oxides: Highly Selective Alkenylation and Direct Arylation with Unactivated Arenes. *J. Am. Chem. Soc.* **130** (29), 9254-9256, doi: 10.1021/ja8026295 (2008).
20. Ide, Y., et al. Spin-crossover between high-spin (*S* = 5/2) and low-spin (*S* = 1/2) states in six-coordinate iron(III) porphyrin complexes having two pyridine-*N* oxide derivatives. *Dalton Trans.* **46** (1), 242-249, doi: 10.1039/C6DT03859J (2017).
21. Drago, R.S. Physical Methods in Chemistry. Saunders College Publishing USA. 750 pp (1977).





22. Cervantes-Mejía, V., et al. Branched Polyamines Functionalized with Proposed Reaction Pathways Based on  $^1\text{H}$ -NMR, Atomic Absorption and IR Spectroscopies. *American Journal of Analytical Chemistry*. **5** (16), 1090-1101, doi: 10.4236/ajac.2014.516116 (2014).
23. Huheey, J. E., Keiter, E. A., Keiter, R. L. Inorganic Chemistry: Principles of Structure and Reactivity. Oxford University Press Mexico. 4<sup>th</sup> Edition, 1023 pp, ISBN: 9706131620 (1997).
24. Rigaku. *CrysAlisPRO*. (2013).
25. Sheldrick, G. M. *SHELXT* – Integrated space-group and crystal-structure determination *Acta Cryst.* **A71** (1), 3–8 doi: 10.1107/S2053273314026370 (2015).
26. Sheldrick, G. M. Crystal structure refinement with *SHELXL*. *Acta Cryst.* **C71** (1), 3–8, doi: 10.1107/S2053229614024218 (2015).
27. Sheldrick, G. M. A short history of *SHELX*. *Acta Cryst.* **A64** (1), 112–122, doi: 10.1107/S0108767307043930 (2008).
28. Macrae, C. F., et al. *Mercury CSD 2.0* - new features for the visualization and investigation of crystal structures. *J. Appl. Cryst.* **41** (2), 466-470, doi: 10.1107/S0021889807067908 (2008).
29. PerkinElmer. ChemBioDraw Ultra 13. (2013).











| Name of material/Equipment                | Company                          | Catalog number |
|---|----------------------------------|----------------|
| 3,5-lutidine                              | Sigma-Aldrich                    | L4206-500ML    |
| Glacial acetic acid                       | Fermont                          | 3015           |
| Hydrogen peroxide (35%)                   | Sigma-Aldrich                    | 349887-500ML   |
| Na <sub>2</sub> CO <sub>3</sub> anhydrous | Productos Químicos Monterrey     | 1792           |
| Na <sub>2</sub> SO <sub>4</sub> anhydrous | Alfa reactivos                   | 25051-C        |
| CHCl <sub>3</sub>                         | Fermont                          | 6205           |
| Ethyl eter                                | Mercury Chemist                  | QME0309        |
| Distilled water                           | Comercializadora Química Poblana | not-existent   |
|   |                                  |                |
|   |                                  |                |
|   |                                  |                |

[illegible]





1 Alewife Center #200  
 Cambridge, MA 02140  
 tel. 617.945.9051  
[www.jove.com](http://www.jove.com)

## ARTICLE AND VIDEO LICENSE AGREEMENT

Title of Article:

Syntheses, crystallization and spectroscopic characterization of 3,5-lutidine N-oxide dehydrate.

Author(s):

María del Rosario Merino-García, Samuel Hernández-Anzaldo and Yasmi Reyes-Ortega

Item 1 (check one box): The Author elects to have the Materials be made available (as described at

<http://www.jove.com/author>) via: ☒ Standard Access ☐ Open Access

Item 2 (check one box):

☒

The Author is NOT a United States government employee.

☐

The Author is a United States government employee and the Materials were prepared in the course of his or her duties as a United States government employee.

☐

The Author is a United States government employee but the Materials were NOT prepared in the course of his or her duties as a United States government employee.

### ARTICLE AND VIDEO LICENSE AGREEMENT

1. **Defined Terms.** As used in this Article and Video License Agreement, the following terms shall have the following meanings: “**Agreement**” means this Article and Video License Agreement; “**Article**” means the article specified on the last page of this Agreement, including any associated materials such as texts, figures, tables, artwork, abstracts, or summaries contained therein; “**Author**” means the author who is a signatory to this Agreement; “**Collective Work**” means a work, such as a periodical issue, anthology or encyclopedia, in which the Materials in their entirety in unmodified form, along with a number of other contributions, constituting separate and independent works in themselves, are assembled into a collective whole; “**CRC License**” means the Creative Commons Attribution-Non Commercial-No Derivs 3.0 Unported Agreement, the terms and conditions of which can be found at: <http://creativecommons.org/licenses/by-nc-nd/3.0/legalcode>; “**Derivative Work**” means a work based upon the Materials or upon the Materials and other pre-existing works, such as a translation, musical arrangement, dramatization, fictionalization, motion picture version, sound recording, art reproduction, abridgment, condensation, or any other form in which the Materials may be recast, transformed, or adapted; “**Institution**” means the institution, listed on the last page of this Agreement, by which the Author was employed at the time of the creation of the Materials; “**JoVE**” means MyJoVE Corporation, a Massachusetts corporation and the publisher of *The Journal of Visualized Experiments*; “**Materials**” means the Article and / or the Video; “**Parties**” means the Author and JoVE; “**Video**” means any video(s) made by the Author, alone or in conjunction with any other parties, or by JoVE or its affiliates or agents, individually or in collaboration with the Author or any other parties, incorporating all or any portion of the Article, and in which the Author may or may not appear.

2. **Background.** The Author, who is the author of the Article, in order to ensure the dissemination and protection of the Article, desires to have the JoVE publish the Article and create and transmit videos based on the Article. In furtherance of such goals, the Parties desire to memorialize in this Agreement the respective rights of each Party in and to the Article and the Video.

3. **Grant of Rights in Article.** In consideration of JoVE agreeing to publish the Article, the Author hereby grants to JoVE, subject to **Sections 4** and **7** below, the exclusive, royalty-free, perpetual (for the full term of copyright in the Article, including any extensions thereto) license (a) to publish, reproduce, distribute, display and store the Article in all forms, formats and media whether now known or hereafter developed (including without limitation in print, digital and electronic form) throughout the world, (b) to translate the Article into other languages, create adaptations, summaries or extracts of the Article or other Derivative Works (including, without limitation, the Video) or Collective Works based on all or any portion of the Article and exercise all of the rights set forth in (a) above in such translations, adaptations, summaries, extracts, Derivative Works or Collective Works and (c) to license others to do any or all of the above. The foregoing rights may be exercised in all media and formats, whether now known or hereafter devised, and include the right to make such modifications as are technically necessary to exercise the rights in other media and formats. If the “Open Access” box has been checked in **Item 1** above, JoVE and the Author hereby grant to the public all such rights in the Article as provided in, but subject to all limitations and requirements set forth in, the CRC License.

## ARTICLE AND VIDEO LICENSE AGREEMENT

4. Retention of Rights in Article. Notwithstanding the exclusive license granted to JoVE in **Section 3** above, the Author shall, with respect to the Article, retain the non-exclusive right to use all or part of the Article for the non-commercial purpose of giving lectures, presentations or teaching classes, and to post a copy of the Article on the Institution's website or the Author's personal website, in each case provided that a link to the Article on the JoVE website is provided and notice of JoVE's copyright in the Article is included. All non-copyright intellectual property rights in and to the Article, such as patent rights, shall remain with the Author.

5. Grant of Rights in Video – Standard Access. This **Section 5** applies if the "Standard Access" box has been checked in **Item 1** above or if no box has been checked in **Item 1** above. In consideration of JoVE agreeing to produce, display or otherwise assist with the Video, the Author hereby acknowledges and agrees that, Subject to **Section 7** below, JoVE is and shall be the sole and exclusive owner of all rights of any nature, including, without limitation, all copyrights, in and to the Video. To the extent that, by law, the Author is deemed, now or at any time in the future, to have any rights of any nature in or to the Video, the Author hereby disclaims all such rights and transfers all such rights to JoVE.

6. Grant of Rights in Video – Open Access. This **Section 6** applies only if the "Open Access" box has been checked in **Item 1** above. In consideration of JoVE agreeing to produce, display or otherwise assist with the Video, the Author hereby grants to JoVE, subject to **Section 7** below, the exclusive, royalty-free, perpetual (for the full term of copyright in the Article, including any extensions thereto) license (a) to publish, reproduce, distribute, display and store the Video in all forms, formats and media whether now known or hereafter developed (including without limitation in print, digital and electronic form) throughout the world, (b) to translate the Video into other languages, create adaptations, summaries or extracts of the Video or other Derivative Works or Collective Works based on all or any portion of the Video and exercise all of the rights set forth in (a) above in such translations, adaptations, summaries, extracts, Derivative Works or Collective Works and (c) to license others to do any or all of the above. The foregoing rights may be exercised in all media and formats, whether now known or hereafter devised, and include the right to make such modifications as are technically necessary to exercise the rights in other media and formats. For any Video to which this Section 6 is applicable, JoVE and the Author hereby grant to the public all such rights in the Video as provided in, but subject to all limitations and requirements set forth in, the CRC License.

7. Government Employees. If the Author is a United States government employee and the Article was prepared in the course of his or her duties as a United States government employee, as indicated in **Item 2** above, and any of the licenses or grants granted by the Author hereunder exceed the scope of the 17 U.S.C. 403, then the rights granted hereunder shall be limited to the maximum rights permitted under such

statute. In such case, all provisions contained herein that are not in conflict with such statute shall remain in full force and effect, and all provisions contained herein that do so conflict shall be deemed to be amended so as to provide to JoVE the maximum rights permissible within such statute.

8. Likeness, Privacy, Personality. The Author hereby grants JoVE the right to use the Author's name, voice, likeness, picture, photograph, image, biography and performance in any way, commercial or otherwise, in connection with the Materials and the sale, promotion and distribution thereof. The Author hereby waives any and all rights he or she may have, relating to his or her appearance in the Video or otherwise relating to the Materials, under all applicable privacy, likeness, personality or similar laws.

9. Author Warranties. The Author represents and warrants that the Article is original, that it has not been published, that the copyright interest is owned by the Author (or, if more than one author is listed at the beginning of this Agreement, by such authors collectively) and has not been assigned, licensed, or otherwise transferred to any other party. The Author represents and warrants that the author(s) listed at the top of this Agreement are the only authors of the Materials. If more than one author is listed at the top of this Agreement and if any such author has not entered into a separate Article and Video License Agreement with JoVE relating to the Materials, the Author represents and warrants that the Author has been authorized by each of the other such authors to execute this Agreement on his or her behalf and to bind him or her with respect to the terms of this Agreement as if each of them had been a party hereto as an Author. The Author warrants that the use, reproduction, distribution, public or private performance or display, and/or modification of all or any portion of the Materials does not and will not violate, infringe and/or misappropriate the patent, trademark, intellectual property or other rights of any third party. The Author represents and warrants that it has and will continue to comply with all government, institutional and other regulations, including, without limitation all institutional, laboratory, hospital, ethical, human and animal treatment, privacy, and all other rules, regulations, laws, procedures or guidelines, applicable to the Materials, and that all research involving human and animal subjects has been approved by the Author's relevant institutional review board.

10. JoVE Discretion. If the Author requests the assistance of JoVE in producing the Video in the Author's facility, the Author shall ensure that the presence of JoVE employees, agents or independent contractors is in accordance with the relevant regulations of the Author's institution. If more than one author is listed at the beginning of this Agreement, JoVE may, in its sole discretion, elect not take any action with respect to the Article until such time as it has received complete, executed Article and Video License Agreements from each such author. JoVE reserves the right, in its absolute and sole discretion and without giving any reason therefore, to accept or decline any work submitted to JoVE. JoVE and its employees, agents and independent contractors shall have

## ARTICLE AND VIDEO LICENSE AGREEMENT

full, unfettered access to the facilities of the Author or of the Author's institution as necessary to make the Video, whether actually published or not. JoVE has sole discretion as to the method of making and publishing the Materials, including, without limitation, to all decisions regarding editing, lighting, filming, timing of publication, if any, length, quality, content and the like.

11. **Indemnification.** The Author agrees to indemnify JoVE and/or its successors and assigns from and against any and all claims, costs, and expenses, including attorney's fees, arising out of any breach of any warranty or other representations contained herein. The Author further agrees to indemnify and hold harmless JoVE from and against any and all claims, costs, and expenses, including attorney's fees, resulting from the breach by the Author of any representation or warranty contained herein or from allegations or instances of violation of intellectual property rights, damage to the Author's or the Author's institution's facilities, fraud, libel, defamation, research, equipment, experiments, property damage, personal injury, violations of institutional, laboratory, hospital, ethical, human and animal treatment, privacy or other rules, regulations, laws, procedures or guidelines, liabilities and other losses or damages related in any way to the submission of work to JoVE, making of videos by JoVE, or publication in JoVE or elsewhere by JoVE. The Author shall be responsible for, and shall hold JoVE harmless from, damages caused by lack of sterilization, lack of cleanliness or by contamination due to the making of a video by JoVE its employees, agents or independent contractors. All sterilization, cleanliness or decontamination procedures shall be solely the responsibility of the Author and shall be undertaken at the Author's


expense. All indemnifications provided herein shall include JoVE's attorney's fees and costs related to said losses or damages. Such indemnification and holding harmless shall include such losses or damages incurred by, or in connection with, acts or omissions of JoVE, its employees, agents or independent contractors.

12. **Fees.** To cover the cost incurred for publication, JoVE must receive payment before production and publication the Materials. Payment is due in 21 days of invoice. Should the Materials not be published due to an editorial or production decision, these funds will be returned to the Author. Withdrawal by the Author of any submitted Materials after final peer review approval will result in a US\$1,200 fee to cover pre-production expenses incurred by JoVE. If payment is not received by the completion of filming, production and publication of the Materials will be suspended until payment is received.

13. **Transfer, Governing Law.** This Agreement may be assigned by JoVE and shall inure to the benefits of any of JoVE's successors and assignees. This Agreement shall be governed and construed by the internal laws of the Commonwealth of Massachusetts without giving effect to any conflict of law provision thereunder. This Agreement may be executed in counterparts, each of which shall be deemed an original, but all of which together shall be deemed to be one and the same agreement. A signed copy of this Agreement delivered by facsimile, e-mail or other means of electronic transmission shall be deemed to have the same legal effect as delivery of an original signed copy of this Agreement.

A signed copy of this document must be sent with all new submissions. Only one Agreement required per submission.

### CORRESPONDING AUTHOR:

|                |  |                  |
|----------------|--|------------------|
| Name:          | Yasmi Reyes-Ortega   |                  |
| Department:    | Centro de Química  |                  |
| Institution:   | Benemérita Universidad Autónoma de Puebla  |                  |
| Article Title: | Syntheses, crystallization and spectroscopic characterization of 3,5-lutidine N-oxide dehydrate. |                  |
| Signature:     |               | Date: 30/08/2017 |

Please submit a signed and dated copy of this license by one of the following three methods:

- 1) Upload a scanned copy of the document as a pdf on the JoVE submission site;
- 2) Fax the document to +1.866.381.2236;
- 3) Mail the document to JoVE / Attn: JoVE Editorial / 1 Alewife Center #200 / Cambridge, MA 02139

For questions, please email [submissions@jove.com](mailto:submissions@jove.com) or call +1.617.945.9051

Journal of Visualized Experiments

Syntheses, crystallization and spectroscopic characterization of 3,5-lutidine N-oxide dehydrate

Ref. JoVE57233R1

Dear Editor Bing Wu,

Thank you for your comments on our manuscript. We have considered fully your concerns and we hope to have appropriately answered all them; this is a fully revised manuscript which we are resubmitting to Journal of Visualized Experiments.

We have modified the manuscript accordingly, and detailed corrections are listed below point by point.

1. There are still some protocol steps which are not in imperative tense. Please rewrite those steps

[In page 3](#)

[line 92](#)

1.2 Cool the flask to 24 °C, after the reaction time, with ice to make it manageable without exposure to the acetic acid gases, and plugged it to a high vacuum distillation unit for 90 to 120 min to remove excess acetic acid. Caution, do not use hot material, wait until the glassware reaches a manageable temperature, this will also avoid that the vapors up stream in top of distillation unit.

[Line 98](#)

- 1.3 Add distilled water (10 mL) twice to ensure the removal of any trace of acetic acid and to concentrate the mixture as far as possible.

[Line 106](#)

2.2 Place carefully, the solution in a separation funnel of 250 mL and wash it 5 times with 250 mL of  $\text{CHCl}_3$  to improve the yield. Recover and dry over solid  $\text{Na}_2\text{SO}_4$  for 30 min maximum the organic phase, which containing the product. If necessary re-extract the aqueous phase with the desired amount of  $\text{CHCl}_3$ .

[Line 113](#)

2.3 Remove the solvent under reduced pressure with a high vacuum distillation unit, until the formation of a very hygroscopic clear beige crystalline powder (70%).

Line 121

... pour the filtrated out liquid into a glass Petri dish, leaving it to slow evaporation at 4 °C in a laboratory fridge.

Line 124

3.2 Ensure that, after two days, are obtained clear colorless crystals. Then measure the melting point which has to be in the range of 310–311 K.

Line 129

4.1 Remove by decantation from the flask's walls the crystals that are formed, of prismatic shape and colorless, for further X-ray analysis. Keep them into diethyl ether to avoid crystals hydration in case of not immediately use.