

Journal of Visualized Experiments

Whole Body Vibration Methods with Survivors of Polio

--Manuscript Draft--

Article Type:	Invited Methods Article - JoVE Produced Video
Manuscript Number:	JoVE58449R3
Full Title:	Whole Body Vibration Methods with Survivors of Polio
Keywords:	poliomyelitis; post-polio syndrome; whole body vibration; weight bearing exercise; therapeutic exercise; feasibility
Corresponding Author:	Carolyn P Da Silva Texas Woman's University - Houston Center Houston, Texas UNITED STATES
Corresponding Author's Institution:	Texas Woman's University - Houston Center
Corresponding Author E-Mail:	cdasilva@twu.edu
Order of Authors:	Carolyn P Da Silva
Additional Information:	
Question	Response
Please indicate whether this article will be Standard Access or Open Access.	Open Access (US\$4,200)
Please indicate the city, state/province, and country where this article will be filmed . Please do not use abbreviations.	6700 Fannin St, Houston, TX 77030

TITLE:

Whole Body Vibration Methods with Survivors of Polio

AUTHORS & AFFILIATIONS:

Carolyn P Da Silva^{1,2}

¹School of Physical Therapy, Texas Woman's University, Houston, TX, USA

²Outpatient Medical Clinic, TIRR Memorial Hermann Rehabilitation and Research, Houston, TX, USA

Corresponding Author:

Carolyn P Da Silva

Email: cdasilva@twu.edu

Tel: 713-794-2087

KEYWORDS:

Poliomyelitis, post-polio syndrome, whole body vibration, weight bearing exercise, therapeutic exercise, feasibility

SHORT ABSTRACT:

The goal of this article is to highlight the strengths, limitations, and applications of the method used with whole body vibration on polio survivors with and without post-polio syndrome as a feasible and safe form of weight bearing exercise.

LONG ABSTRACT:

The purpose of the original study was to examine the use of whole body vibration (WBV) on polio survivors with and without post-polio syndrome as a form of weight bearing exercise. The goal of this article is to highlight the strengths, limitations, and applications of the method used. Fifteen participants completed two intervention blocks with a wash-out period in between the blocks. Each block consisted of twice a week (four weeks) WBV interventions, progressing from 10 to 20 min per session. Low intensity (peak to peak displacement 4.53 mm, frequency 24 Hz, g force 2.21) and higher intensity (peak to peak displacement 8.82 mm, frequency 35 Hz, g force 2.76) WBV blocks were used. Pain significantly improved in both groups following higher intensity vibration. Walking speed significantly improved in the group who participated in higher intensity intervention first. No study-related adverse events occurred. Even though this population can be at risk of developing overuse-related muscle weakness, fatigue, or pain from excessive physical activity or exercise, the vibration intensity levels utilized did not cause significant muscle weakness, pain, fatigue, or sleep disturbances. Therefore, WBV appears to provide a safe method of weight bearing exercise for this population. Limitations included the lack of measurement of reflexes, muscular activity, or circulation, the difficulty in participant recruitment and insufficient strength of some participants to stand in recommended position. Strengths included a standard, safe protocol with intentional monitoring of symptoms and the heterogeneity of the participants in their physical abilities. An application of the methods is the home use of WBV to reduce the barriers associated with going to a facility for weight bearing exercise for longer term interventions, and benefits for conditions such as osteoporosis, particularly for aging adults with

mobility difficulties due to paralysis or weakness. Presented method may serve as a starting point in future studies.

INTRODUCTION:

Whole body vibration (WBV) is reported to give benefits similar to customary exercise in adults, those who are normal or with medical conditions. A vibration platform provides motorized, oscillatory movement to the whole body while the person stands on it. Improvements in pain^{1,2}, strength^{3,4}, balance⁵⁻⁷, bone density^{4,6,8,9}, and flexibility⁵ have been reported. In del Pozo-Cruz *et al.*'s¹⁰ systematic review of WBV studies with people with neurologic conditions, vibration frequencies ranged from 2 to 50 Hz, amplitudes from 0 to 14 mm, 1 to 11 bouts with 1-3 min of rest between bouts, total vibration time per sequence of 0.5 to 15 min, and number of sessions from 1 to 240. WBV acute significant effects were seen in functional balance tests and quadriceps isometric strength in people with multiple sclerosis, postural control and movement items on the Unified Parkinson Disease Rating Scale (UPDRS) in people with Parkinson's disease, and quadriceps isometric and eccentric strength in people post stroke. The only significant change that lasted through the follow-up period was in the UPDRS in people with Parkinson's disease. The reviewers reported that there are few high quality methodological studies available for review and poor consistency of the data between studies¹⁰. A pilot study with adults with chronic incomplete spinal cord injury showed significantly faster walking speed and improvement in various gait parameters¹¹. However, a small study with survivors of polio was discontinued due to preliminary data analysis failing to show significant improvement in walking speed and strength¹².

Because of the large variations of frequencies and magnitudes used in the WBV intervention studies in the neurologic populations¹⁰, the studies reporting the effects of WBV with bone mineral density in post-menopausal women^{4,6,8,9}, older adults^{6,13,14}, and people with fibromyalgia¹⁵ were reviewed due to most of the current study's population of survivors of polio being of similar ages, experiencing problems in participating in traditional weight bearing exercises, and having problems with pain and fatigue. In these review studies, 10 to 40 Hz (most in 30 to 35 Hz range) vibration frequencies, with 1-8 mm amplitude, and variable g forces were used^{4,6,8-10,13-15}. In 2010, Rauch *et al.*¹⁶ published a manuscript describing the need for the standardization and recommendations for reporting WBV studies.

Survivors of polio, with or without post-polio syndrome (PPS), may be challenged in their determination of methods in which to exercise in weight bearing positions to maintain bone density or for other wellness purposes because of their residual weakness or other symptoms of PPS. Exercise has been reported to potentially overly fatigue already overused muscles and worsen symptoms of pain, fatigue, and increased weakness^{17,18}. Few studies with small sample sizes have shown the exercise to be safe and beneficial in this population¹⁹⁻²⁴.

The goal of the previously conducted study was to examine the use of WBV on polio survivors, with and without PPS, as a possible safe form of weight bearing exercise. Only one prior published study had addressed the use of WBV in people with PPS. The post-polio population has unique issues of chronic muscle weakness or paralysis and joint involvement that impacts their ability to

participate in weight bearing exercise; therefore, an exploratory limited-efficacy feasibility study was determined to be necessary²⁵. This article expands upon and demonstrates the method used in the previously published study²⁶. The purpose of this article is to highlight the strengths, limitations, and applications of the method used in that study.

PROTOCOL:

The protocol being described in this article follows the protocol used in the previously published study that abided by the ethical standards for human research of the Internal Review Boards of Texas Woman's University and Baylor College of Medicine in Houston, Texas, USA²⁶.

1. Participant Screening and Study Preparation

1.1. Screen all participants for inclusion and exclusion criteria by phone, and then ask them again about these criteria while in person during the approved informed consent process.

1.1.1. Ask the participants about their polio history and whether they have been diagnosed with PPS.

1.1.2. Ask if the participants have any concern about being able to bear weight through their lower extremities (LEs) at least 20 min in short blocks of time.

1.2. Review the exclusion criteria commonly reported in the literature and used in the previously published study²⁶ for appropriateness for a future study.

Note: Typical exclusion criteria are body weight more than 227 kg (vibration platform maximum), current medical conditions such as cancer, infection, wound, fracture, thrombosis/embolism, epilepsy, severe migraine headaches, severe vestibular conditions, discopathy, spondylolysis, or implanted devices, metal fixators or joints.

1.2.1. Check the weight limit of the selected or available vibration platform, and weigh the participants in person if they appear close to that body weight maximum.

Note: A typical bathroom scale may not measure weights high enough. Many vibration platforms have weight limits of 227 kg or 500 lbs.

1.3. Determine the inclusion criteria that will be specific to the target population and clinical or laboratory setting. If a physician is not on the study team, consider requiring the participants to seek medical approval from their personal physician, for reasons similar to someone starting a new exercise program.

Note: In the previously reported study²⁶, the participants were English speaking survivors of polio, with or without PPS, 40-85 years of age, and able to weight bear through their LEs up to 20 min. Note that their ability to walk was not an inclusion criterion.

1.4. Document the reasons why the individuals do not agree to participate in the study and why they withdraw, if that occurs. Determine if the withdrawals need to be reported to the Internal Review Board of the institution or facility.

1.5. Explain to the participants what the experimental design is (in lay terms), what to expect in terms of time commitment per session, how many sessions, and how many weeks or months the study will last.

1.6. Explain to the participants that they will be standing up or weight bearing on the vibration platform for several short bouts for each session and what the vibration will feel like.

1.6.1. Show the participants the vibration platform. If the participants seem hesitant during the informed consent process and baseline testing, offer to turn on the platform to show the participants how it works. Inquire if he or she wishes to try it briefly prior to starting the intervention in the next session.

2. Experimental Design

2.1. Use random assignment to groups, if possible. If a randomized controlled study is selected, consider offering a delayed-start intervention for the control group to help avoid disappointment or attrition due to the lack of interest.

Note: The previously conducted study used a random-order cross-over experimental design with two weeks for a wash-out period and a two-week follow-up period; therefore, all participants received the same dosage or amount of vibration intervention with treatment block order randomized (low intensity first/high intensity second or high intensity first/low intensity second). No control group was used due to anticipated small sample size²⁶.

2.2. If the participants will use any type of written log for the symptoms such as cramping or activity levels, provide the log for them in paper or electronic version, based upon their preference. Request their log each week to help them keep up with their documentation and to avoid recall bias issues.

2.3. Build in a follow-up period with repeat testing in the study design to see if any changes persist.

2.4. Be available by email or phone for questions or concerns that may arise during the study period.

3. WBV Interventions

3.1. Schedule the sessions with at least one day off in between sessions, two to three times per week to avoid undue fatigue or muscle soreness.

3.1.1. Select the number of weeks of intervention based upon the current literature and the outcome measures selected.

Note: Because the previous study was a feasibility study, two four-week intervention blocks with treatment order randomized were used to determine if the WBV would be a safe and tolerable method to achieve weight bearing exercise²⁶. Gains in bone mineral density will require months, whereas other gains can occur immediately or in days or weeks.

3.2. Instruct the participants to stand on the WBV platform with the knees slightly bent and weight as evenly distributed between their two LEs as possible^{27,28}. Allow the participants to support themselves with their arms, as needed, for balance during the vibration and when stepping up and down from the platform.

3.3. If the person is not able to stand at all or stand long enough, allow him or her to sit in a chair or wheelchair, with the feet on the platform, forearms resting on thighs, leaning forward to achieve as much weight bearing through the lower body as possible, still keeping the feet on the platform.

3.3.1. If the motorized wheelchair has a seat elevator, raise the seat as high as possible.

3.3.2. If the person will be sitting in a standard chair, ask him or her to sit on some folded blankets or pillows to raise the hips up into a higher position.

Note: The closer the LEs are to a standing position, the more weight bearing will occur through the LEs to enhance vibration stimulation into the body.

3.4. Ask the participants to remove their shoes and LE orthoses (braces) and wear socks (for cleanliness) during the vibration. Offer socks with rubber treads on them if the person is not wearing socks or is apprehensive of standing or stepping on the tile floor of the lab.

3.4.1. Explain to the participants that wearing only socks on their feet will enhance their nerve stimulation to their muscles, spinal cord, and brain for peak effectiveness.

3.4.2. Explain to the participants that the orthoses are to be removed because the vibration may cause rubbing between the orthosis and their skin, putting them at risk for skin breakdown. Inform the participants that it may also cause hardware loosening of metal components of the braces.

3.5. Encourage each person to move into his or her optimal alignment.

Note: For people with significant LE weakness on one or both sides, they may not be able to weight bear symmetrically or stand with knees flexed, especially without their orthoses.

3.6. Use a wedge or cuff weight under the heel to accommodate plantarflex or contractures,

if necessary, to allow weight bearing through as much of the plantar surface of the foot as possible.

3.7. Start with ten 1-min vibration bouts, with 1-min sitting rest periods between the vibration bouts. Increase to ten 2-min vibration bouts, and do not change the rest periods unless needed or requested by the participant. See **Table 1** for a typical intervention progression.

3.7.1. Take the person's vital signs.

3.7.2. Observe this person who had polio as a child step up onto the platform used for the low intensity protocol.

3.7.3. Instruct the person to stand with the knees slightly bent, weight as evenly spread between two feet as possible.

3.7.4. Tell the person that he/she can hold onto the handle if he/she needs to balance, but to hold it as lightly as possible.

3.7.5. Inform the person that he/she will feel the vibration in the soles of his/her feet, radiating up into his/her body.

3.7.6. Ask the person if he/she is ready to start, and when he/she says yes, turn on the machine.

3.7.7. Use a timer set for 1 min for the first session if the platform does not have a preset time choice.

3.7.8. After 1 min of vibration has ended, ask the person to sit down for 1 min to rest, setting a timer, offering to answer any questions heshe may have.

3.7.9. After the 1 min rest, ask the person if he/she is ready to step up and resume the vibration. If so, continue for total of 10 repetitions of vibration.

3.7.10. If the person is not ready to continue, ask why, and provide more rest, as needed. Document how much rest time is required, if different than the established protocol.

3.7.11. Repeat this sequence with the higher intensity protocol, but inform the person that the intensity is higher, and he/she may feel it spread higher into his or her body.

3.7.11.1. Conversely, if using a two-intensity protocol and the person starts with the higher intensity intervention first, warn him/her that the second lower intensity will be quite gentle compared to what he/she is used to.

Note: The platform used for higher intensity vibration actually elevated a little, and it gradually lowered during the preset time. As it reached the end of the time, returning to the starting

position, the vibration intensifies, so inform the participant that this phenomenon will occur beforehand to avoid undue concern.

3.8. WBV intensity and equipment

3.8.1. Use a medium to higher intensity (peak to peak vertical displacement 8.82 mm, 35 Hz, and g force 2.76) vibration to allow significant effects. Depending on the targeted population and selected outcome measures, choose more or less aggressive settings.

Note: The previously published study²⁶ used settings in an attempt to simulate that reported for bone mineral density studies, thinking ahead to possible future studies. The second platform listed in the **Table of Materials** was used for this intensity. It is a commercial grade platform that was available in the research lab of Texas Woman's University. The settings used were 35 Hz, "low amplitude", and desired time.

3.8.2. Because low intensity (peak to peak vertical displacement 4.53 mm, 24 Hz, g force 2.21) vibration did not cause any significant changes, do not use this intensity or less unless the targeted study population is very frail, fearful, or both. Then if the participants are able to tolerate this intensity, gradually increase it for each person.

Note: The first platform listed in the **Table of Materials** was used for this intensity. It was purchased for the original study through Post-Polio Health International. It is a unit designed for home or personal use. It has no handles, and a grab rail was installed on the laboratory wall to ensure safety. This platform had a rotating dial, and the lowest setting was selected. A separate timer was used for on and off times.

3.8.3. Ideally, select a vibration platform that the amplitude and frequency can be set independently for individualization to the unique needs of the study population. See **Table of Materials** for the vibration platforms used in the previously reported study²⁶.

Note: Both units used for the reported study had synchronous vertical displacement during vibration. Some units have asynchronous vertical displacement, rotating over a center; in these, the wider the person's base of support is [feet further from center], the more intense the vibration¹⁶.

3.8.4. Use an independent accelerometer to determine peak to peak displacement, frequency, and g forces.

Note: Manufacturers may not provide all required information, and it is needed for accurate reporting of methods and equipment used¹⁶. G forces reported above were measured by an accelerometer listed in **Table of Materials**.

3.8.5. When reporting the amplitude of vibration platforms, be careful with the terminology due to inconsistencies in the literature.

Note: The term amplitude is to be used for equilibrium point to peak displacement. Although the term peak to peak displacement is synonymous with peak to peak amplitude, and this value will be twice that of traditional amplitude, the use of peak to peak displacement is preferred to avoid confusion¹⁶.

4. Participant Testing

4.1. Take heart rate and blood pressure prior to all testing and intervention sessions, especially if working with people with neurologic conditions. Keep in mind that hypertension is considered a “silent killer” and that many people may have this problem and not be aware of it or fail to take prescribed medications for it.

4.1.1. Take the participants’ vital signs again, after at least the first few sessions to ensure safety, and during sessions if the participants experience symptoms that could indicate excessive exertion or abnormal responses to vibration.

4.2. Ask the participants to report perceived exertion with a standardized scale, after each intervention session.

Note: Borg’s rating of perceived exertion [RPE] is widely used and has sound psychometric properties²⁹.

4.3. To reduce testing bias, have a licensed physical therapist blinded to intervention administer selected outcome measures pre and post each intervention block and at the end of the follow-up period.

Note: A double blinded study with vibration interventions will be difficult to conduct because the participants will be able to feel the vibration.

4.4. Select reliable and valid outcome measures that address more than one domain (body functions and structure, activities, and participation) of the International Classification of Functioning, Disability and Health (ICF) of the targeted population³⁰.

4.4.1. When available, use measures that have sound psychometric properties for the population. For less common populations, use more generic measures or ones that have been tested for populations similar in some way. Consider the use of the following outcome measures for survivors of polio, although none were developed specifically for them.

4.4.2. Measure physical performance of gait speed by 10 m walk test (10mWT)³¹ and gait endurance by 2 min walk test³².

4.4.3. Administer surveys to determine pain severity and interference by the Brief Pain Inventory (BPI)^{33,34}, sleep quality by the Pittsburgh Sleep Quality Index³⁵, and fatigue by the

Fatigue Severity Scale^{36,37}.

4.4.4. Measure LE strength by manual muscle test and handheld dynamometry³⁸⁻⁴⁰.

4.4.5. Provide log forms or calendar for participants to document muscle cramping incidence.

Note: No sensory outcome measures were selected in the previously reported study due to the fact that sensory impairments in polio survivors occur only in the presence of other medical conditions or comorbidities. The original poliomyelitis infection affected the anterior horn cells of the spinal cord (lower motor neurons) impacting motor function while preserving sensory function^{17,18}.

REPRESENTATIVE RESULTS:

Fifteen of the 21 survivors of polio who consented completed the study, with 14 of those completing the study having PPS. The seven people who withdrew did so for nonstudy-related reasons (six work and/or time issues, one medical concern). See **Table 2** for demographic characteristics of the participants²⁶.

No significant differences were found between demographic characteristics and pre-intervention outcome measures between the two treatment groups using descriptive statistics and Mann-Whitney *U* tests. Mann-Whitney *U* tests were also conducted between the high and low intensity conditions on measures at pretest, posttest, and follow-up for between subject differences. Within subject changes between pretests and posttests were conducted by Wilcoxon signed-rank tests, and Friedman's analysis of variance test was used to study the change over time from pretest through follow-up due to total of five testing sessions to reduce the risk of committing a Type I error with multiple Wilcoxon signed-rank tests. Nonparametric tests were conducted due to the small sample size and lack of normality^{26,41}. Pain severity as measured by the BPI improved significantly after the higher intensity WBV intervention regardless of treatment order ($Z = -1.97$, $p = 0.049$, Cohen's $d = 0.60$), with BPI pain interference trending towards significant improvement ($Z = -1.92$, $p = 0.055$, Cohen's $d = 0.81$). Gait speed significantly improved after the higher intensity intervention, but only for the group that participated in the higher intervention block first, even though both groups experienced both intensities of vibration in intervention blocks ($Z = -2.38$, $p = 0.017$, Cohen's $d = -1.294$). See **Table 3** for the data and results for the 10 mWT, BPI pain severity, and BPI pain interference. No significant changes occurred in the other outcome measures and following the low intensity protocol, and no changes that occurred were maintained during the two week follow-up period²⁶.

No study-related adverse events occurred. Even though this population can be at risk of developing overuse-related muscle weakness, fatigue, or pain from excessive physical activity or exercise, the vibration intensity levels utilized did not cause significant muscle weakness, pain, fatigue, or sleep disturbances²⁶. The representative results from the prior study need to be interpreted cautiously because of the small study sample size and the huge variability of muscle weakness, pain, walking patterns, and other issues with which survivors of polio present.

FIGURE AND TABLE LEGENDS:

Table 1. Scheduled WBV protocol, assuming no adverse events or problems tolerating intervention.

Table 2. Demographic characteristics of the participants.

Table 3. Representative results from outcome measures.

DISCUSSION:

Two different vibration intensities were studied due to the author's clinical experience in working with survivors of polio and hearing their reports of increased muscle weakness, pain, fatigue, and other symptoms of PPS. Little evidence is present in the literature related to safe, effective weight bearing exercise parameters for people with PPS, and only one study¹² was published prior to the current study²⁶ with WBV and survivors of polio. To date, only two published articles have studied the use of WBV with polio survivors, and both used the platforms with synchronous vertical oscillations^{12,26}. Therefore, no scientific literature exists related to the use of asynchronous vertical (rotatory) or horizontal vibration in this population.

This author was concerned about possibly worsening their symptoms during the WBV protocols, thereby selecting one intensity fairly typical of that reported in the bone density studies and one much lower, in case the participants would have difficulty tolerating the higher one. The author was vigilant during each vibration session and regularly asked if new symptoms surfaced or typical ones worsened at the beginning of the next session. Group means of RPE after higher intensity sessions were 9.4 to 9.8, or "very light," although the range extended from "no exertion at all" to "extremely hard." Although significant changes only occurred for pain severity across both groups, polio survivors and those who provide care for them may be encouraged that gait endurance, muscle strength, and typical PPS symptoms of fatigue, sleep disturbances, and muscle cramping did not significantly worsen during participation in the protocols. Therefore, a strength of the presented methods includes the use of a standard protocol with intentional monitoring of symptoms by a consistent health care provider knowledgeable about their condition. Because no significant improvements occurred with the low intensity protocol, this intensity or less could possibly be considered for use in a sham intervention in a future randomized controlled study.

Although the knowledge of the effects of WBV with survivors of polio is limited, some critical steps may exist. Due to their typical "type A", hard-working, and eager-to-please personalities, one must ask them specific questions and give them explicit permission to voice concerns or potential problems, such as muscle or joint discomfort, excessive muscular fatigue, or sense of worsened balance, that may arise during the intervention period. Similarly, limiting the WBV total dosage (frequency and intensity)¹⁶ per session to no more than 20 min (in one to two min bouts with forced rests) and scheduling sessions with at least one day off in between can help reduce risk of muscle overuse-related fatigue, pain, and cramping^{17,18}.

The mechanisms behind how WBV affects a person's neurophysiological system is poorly understood. The literature has not reported how changes due to WBV in the Hoffman, stretch,

and tonic vibratory reflexes of polio survivors who have a depleted alpha motor neuron pool due to initial disease destruction are impacted⁴²⁻⁴⁷. Reported circulatory changes⁴², in addition to reports of vibration decreasing delayed-onset muscle soreness after exercise⁴⁸, may enhance the polio survivors' weaker muscles to work at safe levels for longer time periods during functional activities, such as walking. A limitation of the presented method in this article is that measurement of reflexes, muscular activity, and circulation did not occur.

A second limitation of the method may be that it was tested on such a small group of polio survivors. Recruitment of people to come to a university research lab in the Texas Medical Center twice a week for three months was challenging due to the time, energy, and transportation costs of commuting within a huge metropolitan city. However, a noteworthy amount of participant diversity in physical mobility existed in the 15 individuals who completed the previously reported study²⁶, which could be considered a strength. As seen in **Table 2**, they ranged from not being able to walk at all, to walking "5%" of total locomotion, to walking full-time, and using no devices at all to utilizing a pair of knee ankle foot orthoses with crutches. All participants except one were able to independently get onto and off the vibration platforms, using the provided handles or grab bar. Even though the participants were able to use their arms for support as needed or preferred, some were unable to shift their weight onto the weaker leg or stand without mechanically locking their knees because of their weakness. Standing with the knees locked could be a potential joint safety issue for some people, and this could be considered a limitation to this method; however, locking also allows more of the vibration g force to extend more cranially^{27,28}. However, this author was surprised by how positively the individuals responded to the vibration during the sessions, with smiles, early arrivals for appointments, requests to "turn it up", and few to no complaints, even with some them dependent on orthoses or assistive devices for safe walking or appearing physically frail. Participants were noted to spontaneously shift weight and change postural alignment, apparently seeking the body position that "felt best" to them. The vibration was truly "whole body" in that it could be heard in their voices while talking during vibration, and dangling earrings or bracelets could be seen or heard to vibrate.

The presented WBV method seems to be a safe, tolerable, and feasible form of weight-bearing exercise for people with weakness or paralysis from polio and PPS. Short lasting improvements in pain and walking speed for some individuals are encouraging for polio survivors who have limited methods to exercise, especially in weight bearing positions. Further research is necessary in order to study long term use and efficacy of WBV in people with PPS and other neurological conditions, particularly to address decrease of barriers to exercise participation and active pursuit of wellness activities. An application of the presented method is the safe use of a WBV platform designed for home use to reduce the barriers of energy, time, and transportation costs associated with going to a health club or therapy clinic. Use within the home can make a longer term intervention feasible, particularly for people with mobility difficulties. Longer intervention studies (at least eight months) will be needed to determine if WBV can significantly impact bone mineral density in female and male polio survivors who have a higher prevalence of osteopenia and osteoporosis than typically aging adults⁴⁹⁻⁵². Additionally, examining the application of WBV in aging adults with neurologic conditions other than post-polio and its effects on bone mineral density will be important, and the presented methods may serve as a starting point in these studies.

ACKNOWLEDGMENTS:

The original study was funded by Post-Polio Health International in 2013, with no cost extension to 2014. Funding for this article was through Texas Woman's University Libraries' Open Access Fund. The results were presented as a poster at the American Physical Therapy Association Combined Sections Meeting, Anaheim, California, US, 2016, and published as a manuscript in *Physiotherapy Theory and Practice* in 2018 (see reference #26).

The author wishes to acknowledge Elson "Randy" Robertson for his assistance during the filming of the video and participation in the original study. She also wishes to acknowledge the following people for their assistance during the original study: C Lauren Szot, PT, DPT, NCS and Natasha deSa, PT, DPT, NCS as blinded testers and co-authors for original manuscript; Arianne Stoker, PT, DPT, Kelly Hodges, PT, DPT, NCS, Mariana Sanjuan, PT, DPT, and Maggie Strange, PT, DPT for help as DPT students during vibration sessions; and Zoheb Allam, MS and Rene Paulson, PhD for statistical assistance.

DISCLOSURES:

The author has nothing to disclose.

REFERENCES:

1. del Pozo-Cruz, B., *et al.* Effects of whole body vibration therapy on main outcome measures for chronic non-specific low back pain: a single-blind randomized controlled trial. *Journal of Rehabilitation Medicine*. **43** (8), 689-694 (2011).
2. Rittweger, J., Karsten, J., Kautzsch, K., Reeg, P., Felsenberg, D. Treatment of chronic low back pain with lumbar extension and whole-body vibration exercise. *Spine*. **27** (17), 1829-1834 (2002).
3. Marin, P.J., Rhea, M.R. Effects of vibration training on muscle strength: a meta-analysis. *Journal of Strength and Conditioning Research*. **24** (2), 548-556 (2010).
4. von Stengel, S., Kemmler, W., Engelke, K., Kalender, W.A. Effects of whole body vibration on bone mineral density and falls: results of the randomized controlled ELVIS study with postmenopausal women. *Osteoporosis International*. **22** (1), 317-325 (2011).
5. Bautmans, I., Hees, E.V., Lemper, J., Mets, T. The feasibility of whole body vibration in institutionalized elderly persons and its influence on muscle performance, balance and mobility: a randomized controlled trial. *BMC Geriatrics*. **5**, 17 <https://doi.org/10.1186/1471-2318-5-17> (2005).
6. Bruyere, O., *et al.* Controlled whole body vibration to decrease fall risk and improve health-related quality of life of nursing home residents. *Archives of Physical Medicine and Rehabilitation*. **86** (2), 303-307 (2005).
7. Gusi, N., Raimundo, A., Leal, A. Low-frequency vibratory exercise reduces the risk of bone

- fracture more than walking: a randomized controlled trial. *BMC Musculoskeletal Disorders*. **7**, 92 (2006).
8. Gómez-Cabello, A., Ara, I., González-Agüero, A., Casajús, J.A., Vicente-Rodríguez, G. Effects of training on bone mass in older adults: a systematic review. *Sports Medicine*. **42** (4), 301-325 (2012).
9. Slatkowska, L., Alibhai, S.M.H., Beyene, J., Cheung, A.M. Effect of whole-body vibration on BMD: a systematic review and meta-analysis. *Osteoporosis International*. **21** (12), 1969-1980 (2010).
10. del Pozo-Cruz, B., *et al.* Using whole-body vibration training in patients affected with common neurological diseases: a systematic literature review. *Journal of Alternative and Complementary Medicine*. **18** (1), 29-41 (2012).
11. Ness, L.L., Field-Fote, E.C. Whole-body vibration improves walking function in individuals with spinal cord injury: a pilot study. *Gait & Posture*. **30** (4), 436-440 (2009).
12. Brogårdh, C., Flansbjer, U.B., Lexell, J. No effects of whole-body vibration training on muscle strength and gait performance in people with late effects of polio: a pilot study. *Archives of Physical Medicine and Rehabilitation*. **91** (9), 1474-1477 (2010).
13. Ma, C., Lie, A., Sun, M., Zhu, H., Wu, H. Effect of whole-body vibration on reduction of bone loss and fall prevention in postmenopausal women: a meta-analysis and systematic review. *Journal of Orthopaedic Surgery Research*. **11**, 24. doi: 10.1186/s13018-016-0357-2 (2016).
14. Sitjà-Rabert, M., *et al.* Efficacy of whole body vibration exercise in older people: a systematic review. *Disability and Rehabilitation*. **34** (11), 883-893 (2012).
15. Collado-Mateo, D., *et al.* Effects of whole-body vibration therapy in patients with fibromyalgia: a systematic literature review. *Evidence Based Complementary and Alternative Medicine*. 2015:719082. doi: 10.1155/2015/719082 (2015).
16. Rauch, F., *et al.* International Society of Musculoskeletal and Neuronal Interactions. Reporting whole-body vibration intervention studies: Recommendations of the International Society of Musculoskeletal and Neuronal Interactions. *Journal of Musculoskeletal and Neuronal Interactions*. **10** (3), 193-198 (2010).
17. Bruno, R. *The polio paradox*. Warner Books. New York City. (2002).
18. Quiben, M.U. Management of chronic impairments in individuals with nervous system conditions. In: Umphred, D.A., Lazaro, R.T., Roller, M.L., Burton, G.U. (editors) *Neurological rehabilitation, 6th edition*. Mosby. St. Louis. (2013).
19. Agre, J.C., Rodriquez, A.A., Franke, T.M. Strength, endurance, and work capacity after muscle

- strengthening exercise in postpolio subjects. *Archives of Physical Medicine and Rehabilitation*. **78** (7), 681-686 (1997).
20. Chan, K.M., Amirjani, N., Sumrain, M., Clarke, A., Strohschein, F.J. Randomized controlled trial of strength training in post-polio patients. *Muscle & Nerve*. **27** (3), 332-338 (2003).
21. Ernstoff, B., Wetterqvist, H., Kvist, H., Grimby, G. Endurance training effect on individuals with postpoliomyelitis. *Archives of Physical Medicine and Rehabilitation*. **77** (9), 843-848 (1996).
22. Klein, M.G., Whyte, J., Esquenazi, A., Keenan, M.A., Costello, R. A comparison of the effects of exercise and lifestyle modification on the resolution of overuse symptoms of the shoulder in polio survivors: a preliminary study. *Archives of Physical Medicine and Rehabilitation*. **83** (5), 708-713 (2002).
23. Koopman, F.S., Beelen, A., Gilhus, N.E., de Visser, M., Nollet, F. Treatment for postpolio syndrome (review). *Cochrane Database of Systematic Reviews*. May 18;(5):CD007818. doi: 10.1002/14651858.CD007818.pub3. (2015).
24. Kriz, J.L., *et al.* Cardiorespiratory responses to upper extremity aerobic training by postpolio subjects. *Archives of Physical Medicine and Rehabilitation*. **73** (1), 49-54 (1992).
25. Bowen, D.J., *et al.* How we design feasibility studies. *American Journal of Preventative Medicine*. **36** (5), 452-457 (2009).
26. Da Silva, C.P., Szot, C.L., deSa, N. Whole body vibration on people with sequelae of polio. *Physiotherapy Theory and Practice*. Mar 29:1-11. doi: 10.1080/09593985.2018.1454559. [Epub ahead of print] (2018).
27. Abercromby, A.F.J., *et al.* Vibration exposure and biodynamic responses during whole-body vibration training. *Medicine & Science in Sports & Exercise*. **39** (10), 1794-1800 (2007).
28. Di Giaminiani, R., Masedu, F., Tihanyi, J., Scrimaglio, R., Valenti, M. The interaction between body position and vibration frequency on acute response to whole body vibration. *Journal of Electromyography and Kinesiology*. **23** (1), 245-251 (2013).
29. Voorn, E.L., Gerrits, K.H., Koopman, F.S., Nollet, F., Beelen, A. Determining the anaerobic threshold in postpolio syndrome: comparison with current guidelines for training intensity prescription. *Archives of Physical Medicine and Rehabilitation*. **95** (5), 935-940 (2014).
30. World Health Organization. <http://www.who.int/classifications/icf/en/> Accessed May 31, 2018.
31. Flansbjer, U.B., Lexell, J. Reliability of gait performance tests in individuals with late effects of polio. *PM&R*. **2** (2), 125-131 (2010).

32. Stolwijk-Swüste, J.M., Beelen, A., Lankhorst, G.J., Nollet, F., and the CARPA Study Group, SF36 physical functioning scale and 2-minute walk test advocated as core qualifiers to evaluate physical functioning in patients with late-onset sequelae of poliomyelitis. *Journal of Rehabilitation Medicine*. **40** (5), 387-392 (2008).
33. McDowell, I. *Measuring health: a guide to rating scales and questionnaires*, 3rd edition. Oxford University Press. New York City. (2006).
34. Mendoza, T.R., et al. The utility and validity of the modified brief pain inventory in a multiple-dose postoperative analgesic trial. *Clinical Journal of Pain* **20** (5), 357-362 (2004).
35. Buysse, D.J., Reynolds, III, C.F., Monk, T.J., Berman, S.R., Kupfer, D.J. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Research*. **28** (2), 193-213 (1989).
36. Horemans, H.L., Nollet, F., Beelen, A., Lankhorst, G.J. A comparison of 4 questionnaires to measure fatigue in patients with postpoliomyelitis syndrome. *Archives of Physical Medicine and Rehabilitation*. **85** (3), 392-398 (2004).
37. Krupp, L.B., LaRocca, N.G., Muir-Nash, J., Steinberg, A.D. The Fatigue Severity Scale: application to patients with multiple sclerosis and systemic lupus erythematosus. *Archives of Neurology* **46** (10), 1121-1123 (1989).
38. Horemans, H.L., Beelen, A., Nollet, F., Jones, D.A., Lankhorst, G.J. Reproducibility of maximal quadriceps strength and its relationship to maximal voluntary activation in postpoliomyelitis syndrome. *Archives of Physical Medicine Rehabilitation*. **85** (8), 1273-1278 (2004).
39. Hislop, H., Avers, D., Brown, M. *Daniels and Worthingham's muscle testing: techniques of manual examination and performance testing*, 9th edition. Elsevier Health Sciences. Philadelphia. (2013).
40. Nollet, F., Beelen, A. Strength assessment in postpolio syndrome: validity of a hand-held dynamometer in detecting change. *Archives of Physical Medicine and Rehabilitation*. **80** (10), 1316-1323 (1999).
41. Portney, L.G., Watkins, M.P. *Foundations of clinical research: applications to practice*, 2nd edition. Prentice Hall Health. Upper Saddle River. (2000).
42. Games, K.E., Sefton, J.M. Whole-body vibration influences lower extremity circulatory and neurological function. *Scandinavian Journal of Medicine & Science in Sports*. **23** (4), 516-523 (2013).
43. Ritzmann, R., Kramer, A., Gollhofer, A., Taube, W. The effect of whole body vibration of the

- H-reflex, the stretch reflex, and the short-latency response during hopping. *Scandinavian Journal of Medicine & Science in Sports*. **23** (3), 331-339 (2013).
44. Trimble, M.H., Koceja, D.M. Modulation of the triceps surae H-reflex with training. *International Journal of Neuroscience*. **76** (3-4), 293-303 (1994).
45. Sefton, J.M., *et al.* Segmental spinal reflex adaptations associated with chronic ankle instability. *Archives of Physical Medicine and Rehabilitation*. **89** (10), 1991-1995 (2008).
46. Wolpaw, J.R. The complex structure of a simple memory. *Trends in Neuroscience*. **20** (12), 588-594 (1997).
47. Pollock, R.D., Woledge, R.C., Martin, F.C., Newham, D.J. Effects of whole body vibration on motor unit recruitment and threshold. *Journal of Applied Physiology*. **112** (3), 388-395 (2012).
48. Lau, W.Y., Nosaka, K. Effect of vibration treatment on symptoms associated with eccentric exercise-induced muscle damage. *American Journal of Physical Medicine and Rehabilitation*. **90** (8), 648-657 (2011).
49. Chang, K.H., *et al.* The relationship between body composition and femoral neck osteoporosis or osteopenia in adults with previous poliomyelitis. *Disability and Health Journal*. **8** (2), 284-289 (2015).
50. Haziza, M., Kremer, R., Benedetti, A., Trojan, D.A. Osteoporosis in a postpolio clinic population. *Archives of Physical Medicine and Rehabilitation*. **88** (8), 1030-1035 (2007).
51. Mohammad, A., Khan, K., Galvin, L., Hardiman, O., O'Connell, P. High incidence of osteoporosis and fractures in an aging post-polio population. *European Neurology*. **62** (6), 369-374 (2009).
52. Centers for Disease Control and Prevention (CDC), National Center for Health Statistics, Osteoporosis. <http://www.cdc.gov/nchs/fastats/osteoporosis.htm> Accessed May 25, 2016.

N/A

Table 1. Scheduled WBV protocol, assuming no adverse events or problems tolerating intervention

	Day 1	Day 2	Day 3	Day 4	Days 5-8
1 WBV on	1 min	1.25 min	1.5 min	1.75 min	2 min
1 Sitting rest off	1 min	1 min	1 min	1 min	1 min
2 WBV on	1 min	1.25 min	1.5 min	1.75 min	2 min
2 Sitting rest off	1 min	1 min	1 min	1 min	1 min
3 WBV on	1 min	1.25 min	1.5 min	1.75 min	2 min
3 Sitting rest off	1 min	1 min	1 min	1 min	1 min
4 WBV on	1 min	1.25 min	1.5 min	1.75 min	2 min
4 Sitting rest off	1 min	1 min	1 min	1 min	1 min
5 WBV on	1 min	1.25 min	1.5 min	1.75 min	2 min
5 Sitting rest off	1 min	1 min	1 min	1 min	1 min
6 WBV on	1 min	1.25 min	1.5 min	1.75 min	2 min
6 Sitting rest off	1 min	1 min	1 min	1 min	1 min
7 WBV on	1 min	1.25 min	1.5 min	1.75 min	2 min
7 Sitting rest off	1 min	1 min	1 min	1 min	1 min
8 WBV on	1 min	1.25 min	1.5 min	1.75 min	2 min
8 Sitting rest off	1 min	1 min	1 min	1 min	1 min
9 WBV on	1 min	1.25 min	1.5 min	1.75 min	2 min
9 Sitting rest off	1 min	1 min	1 min	1 min	1 min
10 WBV on	1 min	1.25 min	1.5 min	1.75 min	2 min
Total time on	10 min	12.5 min	15 min	17.5 min	20 min

	Total Sample (N=19*)	Completing Participants (N=15)
Age (yrs, mean & SD)	63.53 SD 8.32	63.80 SD 9.40
Age onset polio (yrs, mean & SD)	3.55 SD 4.03	3.70 SD 4.80
PPS diagnosis (yes/no)	18/1	14/1
Gender (male/female)	8/11	6/9
Race/ethnicity		
African American/Black	2 (1 mixed)	1
Asian/Pacific islander	1	1
Hispanic/Latino	2	2
Native American	1 (mixed)	0
White	13	11
Walking status		
Full-time	12	11
Part-time	6	3
Not able to walk	1	1
Use of orthoses		
None	13	9
1 or bilateral AFOs	3	3
1 or bilateral KAFOs	2	2
Not applicable	1	1
Use of assistive devices		
None	12	8
1 or bilateral canes/walking sticks	5	5
Bilateral Canadian crutches	1	1
Not applicable	1	1
Working status		
Full-time	6	5
Part-time	4	2
Retired	9	8

* 2 females withdrew from original 21 who consented, prior to data collection.

AFO: Ankle foot orthosis

KAFO: Knee ankle foot orthosis

Table modified and used with permission from original article: Copyright©2018, Taylor & Francis Group, *Physiotherapy Theory and Practice*, Da Silva C.P., et al.²⁶

Table 3. Representative results from outcome measures

Measure	Lo-Hi Group n=6 Mean (Mdn) & SD Min-Max	Hi-Lo Group n=9 Mean (Mdn) & SD Min-Max	Between Subject Differences $p(d)$	Within Subject Differences $p(dz \& \eta^2)$
10mWT (m/s)				
Pre-Hi	1.32(1.07) SD 0.51 0.86-2.11	1.11(1.03) SD 0.39 0.69-1.82	0.346(0.46)	
Post-Hi	1.24(1.09) SD 0.49 0.76-1.96	1.27(1.17) SD 0.47 0.82-2.08	0.698(0.06)	0.087(0.52)
BPI Interference				
Pre-Hi	3.14(2.50) SD 3.14 0.29-7.57	3.62(3.57) SD 2.50 0.00-7.57	0.796(0.17)	
Post-Hi	2.26(.00) SD 3.09 0.00-5.86	2.39(2.57) SD 2.40 0.00-5.14	0.862(0.05)	0.055(0.81)
BPI Severity				
Pre-Hi	3.29(2.75) SD 2.57 0.75-7.25	3.44(3.50) SD 1.74 0.00-5.50	0.795(0.07)	
Post-Hi	2.68(0.13) SD 3.63 0.00-7.25	2.61(2.50) SD 2.65 0.00-5.75	0.674(0.02)	0.049*(0.60)

*Significant difference

Lo-Hi group: low intensity intervention first, higher intensity intervention second

Hi-Lo group: higher intensity intervention first, low intensity intervention second

Hi: Higher intensity intervention

Table modified and used with permission from original article: Copyright©2018, Taylor & Francis Group,

Physiother Theory Pract, Da Silva C.P., et al.²⁶

Name of Material/ Equipment	Company	Catalog Number	Comments/Description
WBV Platform	Soloflex, Inc		Used for low intensity WBV intervention
Power Plate Pro 5	Performance Health Systems, LLC		Used for higher intensity WBV intervention
Trigno tri-axial accelerometer	Delsys, Inc		Used to measure vibration frequencies and g forces



1 Alewife Center #200
Cambridge, MA 02140
tel. 617.945.9051
www.jove.com

ARTICLE AND VIDEO LICENSE AGREEMENT

Title of Article:

Whole body vibration methods with survivors of polio

Author(s):

Carylyn P De Silva

Item 1: The Author elects to have the Materials be made available (as described at <http://www.jove.com/publish>) via:

☐ Standard Access

☒ Open Access

Item 2: Please select one of the following items:

☒ The Author is **NOT** a United States government employee.

☐ The Author is a United States government employee and the Materials were prepared in the course of his or her duties as a United States government employee.

☐ The Author is a United States government employee but the Materials were NOT prepared in the course of his or her duties as a United States government employee.

ARTICLE AND VIDEO LICENSE AGREEMENT

1. **Defined Terms.** As used in this Article and Video License Agreement, the following terms shall have the following meanings: “**Agreement**” means this Article and Video License Agreement; “**Article**” means the article specified on the last page of this Agreement, including any associated materials such as texts, figures, tables, artwork, abstracts, or summaries contained therein; “**Author**” means the author who is a signatory to this Agreement; “**Collective Work**” means a work, such as a periodical issue, anthology or encyclopedia, in which the Materials in their entirety in unmodified form, along with a number of other contributions, constituting separate and independent works in themselves, are assembled into a collective whole; “**CRC License**” means the Creative Commons Attribution-Non Commercial-No Derivs 3.0 Unported Agreement, the terms and conditions of which can be found at: <http://creativecommons.org/licenses/by-nc-nd/3.0/legalcode>; “**Derivative Work**” means a work based upon the Materials or upon the Materials and other pre-existing works, such as a translation, musical arrangement, dramatization, fictionalization, motion picture version, sound recording, art reproduction, abridgment, condensation, or any other form in which the Materials may be recast, transformed, or adapted; “**Institution**” means the institution, listed on the last page of this Agreement, by which the Author was employed at the time of the creation of the Materials; “**JoVE**” means MyJove Corporation, a Massachusetts corporation and the publisher of The Journal of Visualized Experiments; “**Materials**” means the Article and / or the Video; “**Parties**” means the Author and JoVE; “**Video**” means any video(s) made by the Author, alone or in conjunction with any other parties, or by JoVE or its affiliates or agents, individually or in collaboration with the Author or any other parties, incorporating all or any portion

of the Article, and in which the Author may or may not appear.

2. **Background.** The Author, who is the author of the Article, in order to ensure the dissemination and protection of the Article, desires to have the JoVE publish the Article and create and transmit videos based on the Article. In furtherance of such goals, the Parties desire to memorialize in this Agreement the respective rights of each Party in and to the Article and the Video.

3. **Grant of Rights in Article.** In consideration of JoVE agreeing to publish the Article, the Author hereby grants to JoVE, subject to **Sections 4 and 7** below, the exclusive, royalty-free, perpetual (for the full term of copyright in the Article, including any extensions thereto) license (a) to publish, reproduce, distribute, display and store the Article in all forms, formats and media whether now known or hereafter developed (including without limitation in print, digital and electronic form) throughout the world, (b) to translate the Article into other languages, create adaptations, summaries or extracts of the Article or other Derivative Works (including, without limitation, the Video) or Collective Works based on all or any portion of the Article and exercise all of the rights set forth in (a) above in such translations, adaptations, summaries, extracts, Derivative Works or Collective Works and (c) to license others to do any or all of the above. The foregoing rights may be exercised in all media and formats, whether now known or hereafter devised, and include the right to make such modifications as are technically necessary to exercise the rights in other media and formats. If the “Open Access” box has been checked in **Item 1** above, JoVE and the Author hereby grant to the public all such rights in the Article as provided in, but subject to all limitations and requirements set forth in, the CRC License.

ARTICLE AND VIDEO LICENSE AGREEMENT

4. **Retention of Rights in Article.** Notwithstanding the exclusive license granted to JoVE in **Section 3** above, the Author shall, with respect to the Article, retain the non-exclusive right to use all or part of the Article for the non-commercial purpose of giving lectures, presentations or teaching classes, and to post a copy of the Article on the Institution's website or the Author's personal website, in each case provided that a link to the Article on the JoVE website is provided and notice of JoVE's copyright in the Article is included. All non-copyright intellectual property rights in and to the Article, such as patent rights, shall remain with the Author.

5. **Grant of Rights in Video – Standard Access.** This **Section 5** applies if the "Standard Access" box has been checked in **Item 1** above or if no box has been checked in **Item 1** above. In consideration of JoVE agreeing to produce, display or otherwise assist with the Video, the Author hereby acknowledges and agrees that, Subject to **Section 7** below, JoVE is and shall be the sole and exclusive owner of all rights of any nature, including, without limitation, all copyrights, in and to the Video. To the extent that, by law, the Author is deemed, now or at any time in the future, to have any rights of any nature in or to the Video, the Author hereby disclaims all such rights and transfers all such rights to JoVE.

6. **Grant of Rights in Video – Open Access.** This **Section 6** applies only if the "Open Access" box has been checked in **Item 1** above. In consideration of JoVE agreeing to produce, display or otherwise assist with the Video, the Author hereby grants to JoVE, subject to **Section 7** below, the exclusive, royalty-free, perpetual (for the full term of copyright in the Article, including any extensions thereto) license (a) to publish, reproduce, distribute, display and store the Video in all forms, formats and media whether now known or hereafter developed (including without limitation in print, digital and electronic form) throughout the world, (b) to translate the Video into other languages, create adaptations, summaries or extracts of the Video or other Derivative Works or Collective Works based on all or any portion of the Video and exercise all of the rights set forth in (a) above in such translations, adaptations, summaries, extracts, Derivative Works or Collective Works and (c) to license others to do any or all of the above. The foregoing rights may be exercised in all media and formats, whether now known or hereafter devised, and include the right to make such modifications as are technically necessary to exercise the rights in other media and formats. For any Video to which this **Section 6** is applicable, JoVE and the Author hereby grant to the public all such rights in the Video as provided in, but subject to all limitations and requirements set forth in, the CRC License.

7. **Government Employees.** If the Author is a United States government employee and the Article was prepared in the course of his or her duties as a United States government employee, as indicated in **Item 2** above, and any of the licenses or grants granted by the Author hereunder exceed the scope of the 17 U.S.C. 403, then the rights granted hereunder shall be limited to the maximum

rights permitted under such statute. In such case, all provisions contained herein that are not in conflict with such statute shall remain in full force and effect, and all provisions contained herein that do so conflict shall be deemed to be amended so as to provide to JoVE the maximum rights permissible within such statute.

8. **Protection of the Work.** The Author(s) authorize JoVE to take steps in the Author(s) name and on their behalf if JoVE believes some third party could be infringing or might infringe the copyright of either the Author's Article and/or Video.

9. **Likeness, Privacy, Personality.** The Author hereby grants JoVE the right to use the Author's name, voice, likeness, picture, photograph, image, biography and performance in any way, commercial or otherwise, in connection with the Materials and the sale, promotion and distribution thereof. The Author hereby waives any and all rights he or she may have, relating to his or her appearance in the Video or otherwise relating to the Materials, under all applicable privacy, likeness, personality or similar laws.

10. **Author Warranties.** The Author represents and warrants that the Article is original, that it has not been published, that the copyright interest is owned by the Author (or, if more than one author is listed at the beginning of this Agreement, by such authors collectively) and has not been assigned, licensed, or otherwise transferred to any other party. The Author represents and warrants that the author(s) listed at the top of this Agreement are the only authors of the Materials. If more than one author is listed at the top of this Agreement and if any such author has not entered into a separate Article and Video License Agreement with JoVE relating to the Materials, the Author represents and warrants that the Author has been authorized by each of the other such authors to execute this Agreement on his or her behalf and to bind him or her with respect to the terms of this Agreement as if each of them had been a party hereto as an Author. The Author warrants that the use, reproduction, distribution, public or private performance or display, and/or modification of all or any portion of the Materials does not and will not violate, infringe and/or misappropriate the patent, trademark, intellectual property or other rights of any third party. The Author represents and warrants that it has and will continue to comply with all government, institutional and other regulations, including, without limitation all institutional, laboratory, hospital, ethical, human and animal treatment, privacy, and all other rules, regulations, laws, procedures or guidelines, applicable to the Materials, and that all research involving human and animal subjects has been approved by the Author's relevant institutional review board.

11. **JoVE Discretion.** If the Author requests the assistance of JoVE in producing the Video in the Author's facility, the Author shall ensure that the presence of JoVE employees, agents or independent contractors is in accordance with the relevant regulations of the Author's institution. If more than one author is listed at the beginning of this Agreement, JoVE may, in its sole

ARTICLE AND VIDEO LICENSE AGREEMENT

discretion, elect not take any action with respect to the Article until such time as it has received complete, executed Article and Video License Agreements from each such author. JoVE reserves the right, in its absolute and sole discretion and without giving any reason therefore, to accept or decline any work submitted to JoVE. JoVE and its employees, agents and independent contractors shall have full, unfettered access to the facilities of the Author or of the Author's institution as necessary to make the Video, whether actually published or not. JoVE has sole discretion as to the method of making and publishing the Materials, including, without limitation, to all decisions regarding editing, lighting, filming, timing of publication, if any, length, quality, content and the like.

12. **Indemnification.** The Author agrees to indemnify JoVE and/or its successors and assigns from and against any and all claims, costs, and expenses, including attorney's fees, arising out of any breach of any warranty or other representations contained herein. The Author further agrees to indemnify and hold harmless JoVE from and against any and all claims, costs, and expenses, including attorney's fees, resulting from the breach by the Author of any representation or warranty contained herein or from allegations or instances of violation of intellectual property rights, damage to the Author's or the Author's institution's facilities, fraud, libel, defamation, research, equipment, experiments, property damage, personal injury, violations of institutional, laboratory, hospital, ethical, human and animal treatment, privacy or other rules, regulations, laws, procedures or guidelines, liabilities and other losses or damages related in any way to the submission of work to JoVE, making of videos by JoVE, or publication in JoVE or elsewhere by JoVE. The Author shall be responsible for, and shall hold JoVE harmless from, damages caused by lack of sterilization, lack of cleanliness or by contamination due to

the making of a video by JoVE its employees, agents or independent contractors. All sterilization, cleanliness or decontamination procedures shall be solely the responsibility of the Author and shall be undertaken at the Author's expense. All indemnifications provided herein shall include JoVE's attorney's fees and costs related to said losses or damages. Such indemnification and holding harmless shall include such losses or damages incurred by, or in connection with, acts or omissions of JoVE, its employees, agents or independent contractors.

13. **Fees.** To cover the cost incurred for publication, JoVE must receive payment before production and publication of the Materials. Payment is due in 21 days of invoice. Should the Materials not be published due to an editorial or production decision, these funds will be returned to the Author. Withdrawal by the Author of any submitted Materials after final peer review approval will result in a US\$1,200 fee to cover pre-production expenses incurred by JoVE. If payment is not received by the completion of filming, production and publication of the Materials will be suspended until payment is received.

14. **Transfer, Governing Law.** This Agreement may be assigned by JoVE and shall inure to the benefits of any of JoVE's successors and assignees. This Agreement shall be governed and construed by the internal laws of the Commonwealth of Massachusetts without giving effect to any conflict of law provision thereunder. This Agreement may be executed in counterparts, each of which shall be deemed an original, but all of which together shall be deemed to be one and the same agreement. A signed copy of this Agreement delivered by facsimile, e-mail or other means of electronic transmission shall be deemed to have the same legal effect as delivery of an original signed copy of this Agreement.

A signed copy of this document must be sent with all new submissions. Only one Agreement is required per submission.

CORRESPONDING AUTHOR

Name:	Carolyn ^P Da Silva	
Department:	School of Physical Therapy	
Institution:	Texas women's University	
Title:	Professor	
Signature:	Carolyn Da Silva	Date: 7-27-18

Please submit a **signed** and **dated** copy of this license by one of the following three methods:

1. Upload an electronic version on the JoVE submission site
2. Fax the document to +1.866.381.2236
3. Mail the document to JoVE / Attn: JoVE Editorial / 1 Alewife Center #200 / Cambridge, MA 02140

7/28/18

Dear Dr. Wu,

Here are the editorial comments that were provided and my replies.

1. Please take this opportunity to thoroughly proofread the manuscript to ensure that there are no spelling or grammar issues. [I have done this now. I “accepted” all changes under track changes for ease of readability. The only track changes that I left in place were specific to the comments below.](#)
2. Please do not highlight notes for filming. [I have removed the highlights.](#)
3. Please split some long steps so that each step contains 2-3 actions and is less than 4 lines. [This has been done, and left in track change format so that you may see them easily.](#)
4. Please revise the text in Protocol to avoid the use of any personal pronouns (e.g., "we", "you", "our" etc.). [I only found one personal pronoun and it has been changed.](#)
5. Please use h, min, s for time units. [I am sorry about this, but I changed several minutes into min.](#)
6. Step 1.3: Please ensure that all text is written in imperative tense. [Yes, it has been done.](#)
7. 1.4: Please ensure that all text is written in imperative tense. [No changes were made because it already was in imperative tense.](#)
8. Steps 4.4.1-4.4.7: Please write each step as complete sentences and in imperative tense. [This was done.](#)
9. Please sign the new Author License Agreement, which is attached to this email. Please upload it to your Editorial Manager account when you submit your revision. [Done and submitted.](#)

Thank you again for the opportunity to work with you and the journal you represent. I look forward to hearing from you and JoVE as the process proceeds.

Sincerely,


Carolyn Da Silva

DaSilva, Carolyn

From: DaSilva, Carolyn
Sent: Tuesday, June 05, 2018 3:50 PM
To: 'US Journal Permissions'
Subject: RE: permission request for table in article

Thank you VERY much and for your quick reply.

CAROLYN DA SILVA, PT, DSc
713-794-2087
cdasilva@twu.edu



From: US Journal Permissions [mailto:USJournalPermissions@taylorandfrancis.com]
Sent: Tuesday, June 05, 2018 3:48 PM
To: DaSilva, Carolyn <cdasilva@twu.edu>
Subject: RE: permission request for table in article

Dear Ms. DaSilva,
We appreciate your author reuse query.

As author, you retain the right to reuse your original work, in whole or in part, in a new publication that you are primary author or editor of.

For entire articles, you may use your Accepted Author Manuscript, or a link to your Version of Record (published version) as stated in the Taylor & Francis Author Reuse Guide:
<https://authorservices.taylorandfrancis.com/sharing-your-work/>

Please provide the full acknowledgment of your original work in your new publication.
This retained right is for non-commercial use only, that is, you are not to receive a large remuneration other than a small royalty payment for the work.

Thank you for contacting us.

Mary Ann Muller – Permissions Coordinator, US Journals Division

My Work Schedule is Tuesday, Wednesday, and Friday.

Find digital versions of our articles on: www.tandfonline.com to use RightsLink, our online permissions web page, for immediate processing of your permission request.

Please Note: Permissions requests for US journals may take up to three weeks for processing due to demand.



Taylor & Francis Group
an **informa** business

530 Walnut Street – 8th floor
Philadelphia | PA | 19106 | United States of America
Direct line: 215-606-4334
Main Office: 215 625 8900, ext. 14334

maryann.muller@taylorandfrancis.com
www.taylorandfrancisgroup.com
www.tandfonline.com

This electronic message and all contents transmitted with it are confidential and may be privileged. They are intended solely for the addressee. If you are not the intended recipient, you are hereby notified that any disclosure, distribution, copying or use of this message or taking any action in reliance on the contents of it is strictly prohibited. If you have received this electronic message in error, please destroy it immediately, and notify the sender.

Informa Group plc | Registered in England & Wales No. 3099067 5 Howick Place, London, SW1P 1WG

From: DaSilva, Carolyn [<mailto:cdasilva@twu.edu>]
Sent: Tuesday, June 5, 2018 4:24 PM
To: US Journal Permissions <USJournalPermissions@taylorandfrancis.com>
Subject: FW: permission request for table in article

From: DaSilva, Carolyn
Sent: Tuesday, June 05, 2018 2:53 PM
To: 'permissionrequest@tandf.co.uk' <permissionrequest@tandf.co.uk>
Subject: permission request for table in article

I went through the Right Link website, but I did not see how to make the type of request I am desiring. I am the primary author of "Whole body vibration on people with sequelae of polio" published in *Physiotherapy Theory and Practice*, on line March 29, 2018. I am wishing to re-use ~ 1/3 of Table 2 from this article for a new work on methods with whole body vibration to display "representative results" in a new open access journal. I am hoping that me using only part of 1 table from my article in a new article by me will qualify for the \$0 license, of course crediting the original source of the table and stating that it was modified from the original.

When I went through the on-line site, \$472 came up as the quick price. Please advise. Thank you,

CAROLYN DA SILVA, PT, DSc
Board Certified Neurologic Clinical Specialist
Professor & Coordinator of Entry Level Studies
School of Physical Therapy
Texas Woman's University
6700 Fannin Street
Houston, TX 77030-2343
P: 713-794-2087 | F: 713-794-2071