

Journal of Visualized Experiments

Supramaximal intensity hypoxic exercise in mice: feasibility and vascular function assessment --Manuscript Draft--

Article Type:	Invited Methods Article - JoVE Produced Video
Manuscript Number:	JoVE58708R3
Full Title:	Supramaximal intensity hypoxic exercise in mice: feasibility and vascular function assessment
Keywords:	Supramaximal intensity, hypoxia, vascular function, mice, treadmill, running
Corresponding Author:	Gregoire P Millet, PhD Universite de Lausanne Faculte de biologie et medecine Lausanne, SWITZERLAND
Corresponding Author's Institution:	Universite de Lausanne Faculte de biologie et medecine
Corresponding Author E-Mail:	gregoire.millet@unil.ch
Order of Authors:	Jessica Lavier Manon Beaumann Steeve Menetrey Lucia Mazzolai Anne-Christine Peyter Maxime Pellegrin Gregoire P Millet, PhD
Additional Information:	
Question	Response
Please indicate whether this article will be Standard Access or Open Access.	Standard Access (US\$2,400)
Please indicate the city, state/province, and country where this article will be filmed . Please do not use abbreviations.	Lausanne University Hospital, Avenue Pierre Decker 5/Rue du Bugnon 27, CH-1011 Lausanne, Vaud, Switzerland

TITLE:**Supramaximal Intensity Hypoxic Exercise and Vascular Function Assessment in Mice****AUTHORS & AFFILIATIONS:**

Jessica Lavier^{1,2}, Manon Beaumann³, Steeve Menetrey³, Lucia Mazzolai¹, Anne-Christine Peyter³,
Maxime Pellegrin¹, Grégoire P. Millet²

¹Division of Angiology, Heart and Vessel Department, Lausanne University Hospital (CHUV),
Lausanne, Switzerland

²Institute of Sport Sciences, Faculty of Biology and Medicine, University of Lausanne, Lausanne,
Switzerland

³Neonatal Research Laboratory, Clinic of Neonatology, Department Woman-Mother-Child,
Lausanne University Hospital, Lausanne, Switzerland

Corresponding author:

Jessica Lavier (Jessica.Lavier@unil.ch)

Email Addresses of Co-Authors:

Manon Beaumann (manon.beaumann@chuv.ch)

Steeve Menetrey (steeve.menetrey@chuv.ch)

Lucia Mazzolai (Lucia.mazzolai@chuv.ch)

Anne-Christine Peyter (Anne-Christine.Peyter@chuv.ch)

Maxime Pellegrin (maxime.pellegrin@chuv.ch)

Grégoire P. Millet (gregoire.millet@unil.ch)

KEYWORDS:

Supramaximal intensity, hypoxia, vascular function, mice, treadmill, running

SUMMARY:

High-intensity training in hypoxia is a protocol that has been proven to induce vascular adaptations potentially beneficial in some patients and to improve athletes' repeated sprint ability. Here, we test the feasibility of training mice using that protocol and identify those vascular adaptations using ex vivo vascular function assessment.

ABSTRACT:

Exercise training is an important strategy for maintaining health and preventing many chronic diseases. It is the first line of treatment recommended by international guidelines for patients suffering from cardiovascular diseases, more specifically, lower extremity artery diseases, where the patients' walking capacity is considerably altered, affecting their quality of life.

Traditionally, both low continuous exercise and interval training have been used. Recently, supramaximal training has also been shown to improve athletes' performances via vascular adaptations, amongst other mechanisms. The combination of this type of training with hypoxia could bring an additional and/or synergic effect, which could be of interest for certain

pathologies. Here, we describe how to perform supramaximal intensity training sessions in hypoxia on healthy mice at 150% of their maximal speed, using a motorized treadmill and a hypoxic box. We also show how to dissect the mouse in order to retrieve organs of interest, particularly the pulmonary artery, the abdominal aorta, and the iliac artery. Finally, we show how to perform ex vivo vascular function assessment on the retrieved vessels, using isometric tension studies.

INTRODUCTION:

In hypoxia, the decreased inspired fraction of oxygen (O_2) leads to hypoxemia (lowered arterial pressure in hypoxia) and an altered O_2 transport capacity¹. Acute hypoxia induces an increased sympathetic vasoconstrictor activity directed toward skeletal muscle² and an opposed 'compensatory' vasodilatation.

At submaximal intensity in hypoxia, this 'compensatory' vasodilatation, relative to the same level of exercise under normoxic conditions, is well established³. This vasodilation is essential to ensure an augmented blood flow and maintenance (or limit the alteration) of oxygen delivery to the active muscles. Adenosine was shown to not have an independent role in this response, while nitric oxide (NO) seems the primary endothelial source since significant blunting of the augmented vasodilatation was reported with nitric oxide synthase (NOS) inhibition during hypoxic exercise⁴. Several other vasoactive substances are likely playing a role in the compensatory vasodilatation during a hypoxic exercise.

This enhanced hypoxic exercise hyperemia is proportional to the hypoxia-induced fall in arterial O_2 content and is larger as the exercise intensity increases, for example during intense incremental exercise in hypoxia.

The NO-mediated component of the compensatory vasodilatation is regulated through different pathways with increasing exercise intensity³: if β -adrenergic receptor-stimulated NO component appears paramount during low-intensity hypoxic exercise, the source of NO contributing to compensatory dilatation seems less dependent on β -adrenergic mechanisms as the exercise intensity increases. There are other candidates for stimulating NO release during higher-intensity hypoxic exercise, such as ATP released from erythrocytes and/or endothelial-derived prostaglandins.

Supramaximal exercise in hypoxia (named repeated sprint training in hypoxia [RSH] in the exercise physiology literature) is a recent training method⁵ providing performance enhancement in team- or racket-sport players. This method differs from interval training in hypoxia performed at or near maximal speed⁶ (V_{max}) since RSH performed at maximal intensity leads to a greater muscle perfusion and oxygenation⁷ and specific muscle transcriptional responses⁸. Several mechanisms have been proposed to explain the effectiveness of RSH: during sprints in hypoxia, the compensatory vasodilation and associated higher blood flow would benefit the fast-twitch fibers more than the slow-twitch fibers. Consequently, RSH efficiency is likely to be fiber-type selective and intensity dependent. We speculate that the improved responsiveness of the vascular system is paramount in RSH.

Exercise training has been extensively studied in mice, both in healthy individuals and in pathological mouse models^{9,10}. The most common way to train mice is using a rodent treadmill, and the traditionally used regimen is low-intensity training, at 40%–60% of V_{\max} (determined using an incremental treadmill test¹¹), for 30–60 min^{12–15}. Maximal intensity interval training and its impact on pathologies have been widely studied in mice^{16,17}; thus, interval training running protocols for mice have been developed. Those protocols usually consist of about 10 bouts of running at 80%–100% of V_{\max} on a rodent motorized treadmill, for 1–4 min, interspersed with active or passive rest^{16,18}.

The interest in mice exercising at supramaximal intensity (i.e., above the V_{\max}) in hypoxia comes from previous results that the microvascular vasodilatory compensation and the intermittent exercise performance are both more increased at supramaximal than at maximal or moderate intensities. However, to our knowledge, there is no previous report of a supramaximal training protocol in mice, either in normoxia or in hypoxia.

The first aim of the present study was to test the feasibility of supramaximal intensity training in mice and the determination of a tolerable and adequate protocol (intensity, sprint duration, recovery, etc.). The second aim was to assess the effects of different training regimen in normoxia and hypoxia on the vascular function. Therefore, we test the hypotheses that (1) mice tolerate well supramaximal exercise in hypoxia, and (2) that this protocol induces a larger improvement in vascular function than exercise in normoxia but also than exercise in hypoxia at lower intensities.

PROTOCOL:

The local state's animal care committee (Service de la Consommation et des Affaires Vétérinaires [SCAV], Lausanne, Switzerland) approved all experiments (authorization VD3224; 01.06.2017) and all experiments were carried out in accordance with the relevant guidelines and regulations.

1. Animal housing and preparation

1.1. House 6- to 8-week-old C57BL/6J male mice in the animal facility for at least 1 week prior to the beginning of the experiments in order for the mice to get used to their new housing conditions. For practical reasons, mice of the same experimental group are usually housed together.

1.2. Keep the mice in a temperature-controlled room (22 ± 1 °C) with a 12 h light/dark cycle with ad libitum access to food and water.

2. Determination of the maximal speed and standard assessment of performance improvement by treadmill incremental test

NOTE: The following steps are critical to completing the training protocols.

2.1. Use a motorized treadmill for mice where mice can be on multiple lanes alongside each other, with a 0° inclination and mounted with an electric grid set to 0.2 mA at the back of the lane, in order to encourage the mice to run.

2.2. Prior to the first test, submit the mice to 4 days of acclimatization to the treadmill, according to the following protocol.

2.2.1. On day 1, have mice run for 10 min at 4.8 m/min.

2.2.2. On day 2, have the mice run for 10 min at 6 m/min.

2.2.3. On day 3, have the mice run for 10 min at 7.2 m/min.

2.2.4. On day 4, have the mice run for 10 min at 8.4 m/min.

2.3. On day 5, submit the mice to an incremental test to exhaustion, according to the following protocol.

2.3.1. Let the mice warm up for 5 min at 4.8 m/min (at a 0° inclination).

2.3.2. Increase the speed by 1.2 m/min every 3 min (e.g., 5 min at 4.8 m/min, then 3 min at 6 m/min, 3 min at 7.2 m/min, 3 min at 8.4 m/min, etc.) until exhaustion, which is reached when the mouse either spends 3 consecutive seconds on the electric grid or receives 100 shocks (displayed by the apparatus).

2.3.3. Write down the achieved speed (considered as the V_{\max}), duration, distance, number of shocks, and total time spent on the grid.

NOTE: Typically, V_{\max} was 28.8 ± 3.7 m/min.

2.3.4. Mid-training, resubmit the mice to this test in order to readjust the speeds of training to the updated V_{\max} of the mice (e.g., if the training protocol lasts 8 weeks, then perform a mid-training incremental test at 4 weeks. In that case, replace one of the scheduled trainings by the test), and do so again at the end of the study in order to assess performance improvements.

2.3.5. Implement a 48 h rest period before and after this test.

NOTE: All the incremental tests were performed in the morning.

3. Hypoxic environment

3.1. For the training sessions in hypoxia, place the treadmill in the hypoxic box (**Figure 1**) linked to a gas mixer. Use a calibrated oximeter to regularly control the ambient fraction of oxygen (F_{iO_2} [i.e., the level of hypoxia]) in the box.

3.2. Set the gas mixer on 100% of nitrogen (N_2) and use the oximeter to verify the level of hypoxia. Once $F_iO_2 = 0.13$, change the parameter of the gas mixer from 100% N_2 to 13% O_2 .

3.3. In order to avoid prolonged passive exposure to hypoxia, place the mice in a temporary smaller cage with litter and enrichment, and quickly place it in the box once $F_iO_2 = 0.13$ has been reached. Verify that the environment is still at 13% O_2 after putting the cage in; if not, readjust it.

3.4. Regularly verify the level of O_2 over the course of a training session to make sure that it remains at $F_iO_2 = 0.13 \pm 0.002$.

4. Normoxic environment

4.1. For the training sessions in normoxia, keep the treadmill in the hypoxic box, but remove the gloves so that there is ambient air ($F_iO_2 = 0.21$). The aim is to recreate the same training environment as the mice in hypoxia.

5. Supramaximal intensity training

5.1. Place the mice on individual lanes in the treadmill (at a 0° inclination) and submit them to the following protocol.

5.1.1. Have the mice warm up for 5 min at 4.8 m/min, followed by 5 min at 9 m/min.

5.1.2. Set the speed of the sprints to 150% of the previously determined V_{max} .

NOTE: Typically, the sprint velocity was 42.1 ± 5.5 m/min.

5.1.3. Train the mice for four sets of 5x 10 s sprints with 20 s of rest between each sprint. The interset rest is 5 min (Figure 2).

NOTE: Add a cooldown period if the total workload of the training session needs to match that of another training group.

5.2. Perform this training 3x per week, with preferably 48 h between training sessions.

5.3. Use cotton swabs as a complementary method to electric shocks to encourage the mice to run. Place a cotton swab in a slit at the top of the lane, between the mouse and the electric grid, and gently nudge the mouse when it reaches the back of the treadmill. This will avoid the delivery of electric shocks and stimulate the mice to run in a softer way.

6. Low-intensity training

6.1. Place the mice on individual lanes in the treadmill (at a 0° inclination) and submit them to the following protocol.

6.1.1. Have the mice warm up for 5 min at 4.8 m/min, followed by 5 min at 7.2 m/min.

6.1.2. Set the speed of the continuous running session to 40% of the previously determined V_{\max} .

NOTE: Typically, the continuous running velocity was 9.9 m/min.

6.1.3. Train the mice for 40 min.

6.1.4. Perform this training 3x per week with preferably 48 h between training sessions.

6.1.5. Use cotton swabs as a complementary method to electric shocks to encourage the mice to run.

7. Mice euthanasia and organ extraction

7.1. At the end of the training protocol and at least 24 h after the last incremental test, anesthetize the mouse in an induction chamber using isoflurane (4%–5% in O₂ to induce anesthesia, and 1%–2% in O₂ to maintain anesthesia). Confirm proper anesthetization using the paw retraction reflex (firmly pinch the animal's paw; anesthesia is considered proper when the animal does not react to the stimulus).

7.2. Using a 25 G needle, perform a percutaneous cardiac puncture, to collect maximum blood volume as previously described¹⁹.

7.3. Perform a cervical dislocation and remove the skin of the mouse by cutting through the first layer of skin on the abdomen with round-tip scissors and pulling on the two sides of the incision (toward the head and the tail).

7.4. Cut through the peritoneum under the ribcage on the left side of the mouse with thin-point-tip scissors to reach the spleen and extract it if needed.

NOTE: Dissect out muscles if needed.

7.5. Dissect out the pulmonary artery.

7.5.1. Using both small scissors and forceps, remove the thoracic cage and clear the heart-lung area.

7.5.2. With “self-closing” tweezers, pinch the heart as close as possible to the apex and pull gently to stretch the base of the aortic arch and the pulmonary artery.

265 7.5.3. Using the right hand, insert curved tweezers under the pulmonary artery and the aorta,
266 and then move the tweezers back a little to hold only the pulmonary artery (**Figure 3**).
267

268 7.5.4. Use the left hand to insert another pair of tweezers to replace the one held with the right
269 hand.
270

271 7.5.5. Using sharp straight microscissors in the right hand, dissect the pulmonary artery as close
272 to the heart as possible on one side, and as far away as possible on the other side.
273

274 NOTE: It does not matter which hand holds which instrument, although we have found it easier
275 to cut with the right hand than with the left.
276

277 7.5.6. Put it in a 2 mL tube with cold phosphate-buffered saline (PBS) buffer and keep on ice.
278

279 7.6. Perform a whole-body perfusion.
280

281 7.6.1. At the top of the right lower limb of the mouse, use tweezers to clear out the external-
282 internal right iliac artery down to the right femoral artery (under the inguinal ligament). Using
283 sharp straight microscissors, make a full cut in the femoral artery.
284

285 7.6.2. Insert a 5 mL 25 G syringe filled with cold PBS in the left ventricle of the heart and gently
286 inject the cold PBS to remove the remaining blood from the vessels.
287

288 NOTE: Due to the extraction of the pulmonary artery, it is possible that PBS does not circulate all
289 the way to the incision.
290

291 7.7. Using tweezers, remove the soft tissue surrounding the aorta from the left and right inguinal
292 ligaments to the heart as thoroughly as possible.
293

294 NOTE: The heart can be extracted for further analysis if necessary.
295

296 7.8. Using both tweezers and microscissors, dissect out the heart up to the lowest point of the
297 external iliac artery (in both left and right limbs) and place the entirely dissected-out section in a
298 10 cm-diameter dish with cold PBS.
299

300 7.9. Using tweezers and/or microscissors, finish cleaning the remaining fat around the aorta and
301 arteries by gently pulling or cutting it away from the vessels.
302

303 7.10. Using microscissors, cut the left iliac artery at the left-right iliac artery bifurcation and store
304 it for further analysis.
305

306 7.11. Using microscissors, cut the abdominal aorta under the left renal artery, and place the
307 extracted vessel in cold PBS buffer on ice (**Figure 4**).
308

7.12. Keep the remaining cleaned vessel, from the aortic arch to right above the left renal artery, in storage for further analysis.

[Place **Figure 4** here]

8. Ex vivo vascular function assessment

NOTE: A wash corresponds to the emptying and refilling of the chambers with Krebs.

8.1. According to a previously described protocol²⁰, cut the isolated pulmonary artery, abdominal aorta, and right iliac artery segments into vascular rings of 1.5–2.0 mm long and 0.5–1.0 mm in diameter, and mount each ring on two 0.10 mm-diameter stirrups passed through the lumen.

8.2. Suspend the vessel rings in vertical organ chambers filled with 10 mL of modified Krebs-Ringer bicarbonate solution (118.3 mM NaCl, 4.7 mM KCl, 2.5 mM CaCl₂, 1.2 mM MgSO₄, 1.2 mM KH₂PO₄, 25.0 mM NaHCO₃, and 11.1 mM glucose) maintained at 37 °C and aerated with 95% O₂–5% CO₂ (pH 7.4). One stirrup is anchored to the bottom of the organ chamber and the other one is connected to a strain gauge for the measurement of isometric force in grams.

8.3. Bring the vessels to their optimal resting tension: stretch the rings to 0.5 g for the pulmonary artery, 1.5 g for the iliac artery, and 2 g for the abdominal aorta, and wash them after a 20 min period of equilibration. Repeat the stretch-equilibration-wash steps 1x.

8.4. To test the viability of the vessels, contract the rings with 235 µL of KCl (10⁻¹ M) for 10 min, wash them for another 10 min, and contract again with 235 µL of KCl (10⁻¹ M) for about 20 min until reaching a plateau.

8.5. Wash the vessels again for 10 min and add 58.4 µL of indomethacin (10⁻⁵ M) (an inhibitor of cyclooxygenase activity) for at least 20 min in order to avoid possible interference of endogenous prostanoids.

8.6. Add cumulative doses of phenylephrine (Phe) from 10⁻⁹ (10 µL) to 10⁻⁴ M (or 10⁻⁹ to 10⁻⁵ M for the pulmonary artery; 9 µL for all concentrations above 10⁻⁹ M) to contract the vessels.

8.7. After the last dose of Phe, wait for about 1 h until the vessels reach a relatively stable contraction state (plateau).

8.8. Add cumulative doses of the endothelium-dependent vasodilator acetylcholine (ACh), from 10⁻⁹ (58.5 µL) to 10⁻⁴ M (40 µL for all concentrations above 10⁻⁹ M), to induce NO-mediated relaxation.

8.9. At the end of the relaxation curve, wash the vessels for 10 min, and add 58.4 µL of indomethacin (10⁻⁵ M), as well as 184 µL of NG-nitro-L-arginine (NLA, 10⁻⁴ M), which is an inhibitor of the NOS, for at least 20 min.

8.10. Contract the vessels again with a unique dose of 10 μ L of Phe (10^{-5} and 10^{-4} M for the pulmonary artery and 10^{-4} M for the abdominal aorta and the iliac artery) for 1 h, to induce a relatively stable contraction.

8.11. Add a unique dose of 40 μ L of ACh (10^{-4} M) until reaching a plateau.

8.12. Wash the vessels again for 10 min, before adding 58.4 μ L of indomethacin (10^{-5} M) and 184 μ L of NLA (10^{-4} M) for 20 min.

8.13. Contract the vessels with 10 μ L of Phe (10^{-5} and 10^{-4} M) for 1 h.

8.14. Add cumulative doses (10^{-9} [58.4 μ L] to 10^{-4} M [40 μ L for all concentrations above 10^{-9} M]) of the NO donor diethylamine (DEA)/NO, in order to assess the endothelium-independent NO-induced relaxation.

8.15. At the end of the experiment, store the vessels in liquid nitrogen for future analyses if needed.

REPRESENTATIVE RESULTS:

To our knowledge, the present study is the first to describe a program of supramaximal intensity training in normoxia and in hypoxia for mice. In this protocol, mice ran four sets of five 10 s sprints with a 20 s recovery in between each sprint. The sets were interspersed with 5 min of recovery periods. It was unknown whether the mice would be capable of sustaining such a protocol and complete it properly. However, according to **Figure 5**, the body weight gain of the mice undergoing the supramaximal intensity training was similar to that of the mice undergoing the low-intensity training, both in normoxia and in hypoxia.

The wellness of the animals was monitored twice a week, using score sheets, based on the following criteria: appearance, natural behavior, and body weight. Each of those criteria was graded up to a score of 3, and a mouse with a score of 3 in any of those criteria was considered in pain and/or distress due to the sustained protocol and had to be euthanized. No mouse ever reached a score of 3 over the course of any of the training regimens (**Table 1**).

As presented in the introduction, it has been hypothesized that supramaximal training, in particular when combined with hypoxia, would induce a compensatory vasodilation. This phenomenon aims at providing sufficient O_2 to the contracting muscles, thereby compensating for the imbalance between O_2 supply and demand that is enhanced by the combination of supramaximal intensity training and hypoxia. In order to investigate this hypothesis, we used the second technique presented here, the ex vivo vascular function assessment, on the pulmonary artery, the abdominal aorta, and the right iliac artery. **Figure 6** shows the dose-response curves obtained at the end of the protocol, on the abdominal aorta of a mouse from the group training at supramaximal intensity in hypoxia. This graph shows the whole process of contraction-relaxation observed after the addition of different pharmacological agents (KCL, Phe, ACh, NLA,

and [DEA]/NO) in the organ baths.

Figure 7 shows the dose-response relaxation curve for the right iliac artery to increasing concentrations of ACh. The two represented groups are the supramaximal-intensity-in-normoxia group (SupraN) and the supramaximal-intensity-in-hypoxia group (SupraH). The preliminary results show that SupraH tended to improve ACh-induced relaxation compared to SupraN, with significant differences at 10^{-5} M and 10^{-4} M.

FIGURE AND TABLE LEGENDS:

Figure 1: Hypoxic setup. The treadmill is placed inside the homemade glovebox, which is linked to a gas mixer.

Figure 2: Description of a supramaximal intensity training session. The mice performed four sets of five 10 s sprints, interspersed with 20 s of rest. The interset rest was 5 min. This figure is adapted from Faiss et al.²¹.

Figure 3: Schematic representation of the technique to retrieve the pulmonary aorta. (1) Place the tweezers under both the pulmonary artery and the aorta. (2) Pull back the tweezers in the direction of number 2 in order to keep the tweezers under the pulmonary aorta only. (3) Final position of the tweezers.

Figure 4: Picture of the dissected vessels. Extracted vessel from the top of the abdominal aorta (underneath the left renal artery) to the end of the right iliac artery, ready to be placed in cold PBS buffer on ice. (1) Abdominal aorta. (2) Right common iliac artery. (3) External iliac artery. (4) Internal iliac artery. (5) Femoral artery.

Figure 5: Body weight evolution over the course of the experiment. In green, the low-intensity training groups; in red, the supramaximal intensity training groups. There was no significant difference between any of the groups at any of the time points ($n = 4$ mice per group; the data are presented as mean \pm SD). Statistical analysis was performed using a two-way repeated measure ANOVA).

Figure 6: Vascular function assessment curves. Succession of contraction and relaxation phases induced throughout the entire protocol, expressed in grams. Representative recording of the variations in vessel tension in response to the applied substances, in a ring of the abdominal aorta isolated from a mouse trained at supramaximal intensity in hypoxia.

Figure 7: Pharmacological responses of an isolated iliac artery preconstructed with phenylephrine (Phe) to acetylcholine (ACh). Cumulative dose-response relaxation curve of the right iliac artery to increasing concentrations of ACh (10^{-9} to 10^{-4} M). The results are expressed as mean \pm SD of the percentage of change in the tension induced by the vasodilator, with $n = 3$ in SupraN and $n = 4$ in SupraH. Statistical analysis was performed using a two-way ANOVA for repeated measures test. * $p < 0.05$ vs. SupraN.

Table 1: Typical score sheet of a mouse training at supramaximal intensity in hypoxia. We used score sheets to monitor the welfare of the mice. A score of 3 in any of the criteria indicated (appearance, natural behavior, and body weight) or a total score of 5 (by addition of the score of each category) meant the animal was suffering and had to be euthanized.

DISCUSSION:

The first objective of this study was to assess the feasibility of hypoxic high-intensity training in mice and to determine the adequate characteristics of the protocol that would be well tolerated by mice. Purposely, since there is no data using supramaximal (i.e., more than V_{\max}) intensity training in mice, we had to perform trials based on previous protocols developed with athletes, which consisted of four to five sets of five all-out sprints (about 200% of V_{\max}), interspersed with 20 s active recoveries, with an interset active recovery of 5 min^{21,22}. Therefore, the initial protocol consisted of six sets of six 10 s sprints at 200% of V_{\max} , interspersed with 20 s of passive recovery and with an interset passive recovery of 3 min, performed five times per week. After a few try-out runs at 200% of the V_{\max} , considering the mice had trouble sustaining such a high intensity, we decided to lower the speed to 150% of the V_{\max} . With that exercise intensity, we tried to run the mice over the length of a full protocol and adjusted both the number of sprints within each set and the number of sets per session. Finally, we increased the recovery time between sets and decreased the frequency of the training sessions. Following a trial-and-error method, we established a final optimal protocol that is very similar to the one used on athletes and made possible for mice to tolerate this supramaximal intensity test.

There is a slight possibility that the performance of the mice might be severely underestimated, as observed from large differences between previous studies utilizing animal exercise protocols^{23,24}. However, in the present study, based on pre-experiment values, it would have been impossible to impose a higher relative intensity on the animals considering the need to complete the entire repeated sprint session. Moreover, the V_{\max} values reported in this study (28.8 ± 3.7 m/min) seem to be in the range of values previously reported in the same C57BL/6J strain^{25–28}. For example, Lightfoot et al.²⁵ reported values of ~28 m/min and Muller et al.²⁷ values of 28.3 m/min. Therefore, we are confident that the supramaximal intensity corresponds to sprint training intensity in these mice.

Although critical speed (CS) has been shown (1) to be a valuable mean for prescribing exercise intensity in healthy humans and patients²⁹ and (2) to be perfectly determined in mice^{23,24,30}, the exercise intensity prescription based on the determination of V_{\max} remains relevant. It is known that, in mice, the determined $VO_{2\text{peak}}$ and $VO_{2\text{max}}$ depend on the protocol, and, as with humans, $VO_{2\text{max}}$ can be determined with a ramp exercise protocol¹¹. Since the aim of the present study was to determine the feasibility of supramaximal repeated sprint in mice, and despite the relevance of CS, we do not believe that using V_{\max} would be a flaw regarding the objectives of this study.

While observing mice behavior, it became clear that the electric grid at the rear of the treadmill admittedly encouraged mice to run; however, it also seemed to contribute to their fatigue.

Indeed, the grid being slightly shifted from the running band, the mice had to generate an extra effort to get back on the lane. We decided to complement this stimulation with another, softer, one, namely the cotton swab stimulation, which decreased the number of shocks received by the animals and prevented them from having to get back from the grid to the lane. Despite the recommendation by Kregel et al.³¹, it remains unclear whether stress is reduced using the air puff stimulation compared to the electric grid³².

As far as we know, only one study has used “sprint interval training”³³. However, since the highest intensity in that study corresponded to 75%–80% of V_{\max} and the sprint duration was 1.5 min, that protocol was very different from the present one (i.e., 150% of V_{\max} ; 10 s). It was unknown whether supramaximal intensity would be tolerated by the mice. In the present study, we provide results showing that the animals did perform very well in this supramaximal intensity training, both in hypoxia and normoxia. For instance, **Figure 5** shows an increase in body weight over the training period similar to that observed in the low-intensity groups. Similarly, **Table 1** reflects the level of welfare with a score lower than 3 in all groups. Altogether, those physiological parameters indicate that both hypoxia and supramaximal intensity training were very well tolerated by mice.

The second objective of the present study was to assess the vascular function of the pulmonary artery, the abdominal aorta, and the iliac artery, using isometric vessel tension studies²⁰. This technique allows determining whether the intervention of interest impacted the ability of the vessels to contract and relax in response to pharmacological drugs. As shown in **Figure 7**, the iliac artery was relaxed using increasing concentrations of ACh. The observed curves reflect a progressive increase in the relaxation of the vessels, more marked for the SupraH group. If any of the observed curves had been completely flat and around 0% of relaxation, it could mean that the drug was not delivered to the organ chamber, or that the vessels had been damaged during the dissection or the mounting on the stirrups, or that one of the drugs was not administered at the optimal dose or for long enough.

The supramaximal intensity training in hypoxia is now transferred to mice and could potentially be used on pathological models in order to improve various parameters, including vascular function, which can be assessed using isometric vessel tension studies.

ACKNOWLEDGMENTS:

The authors would like to thank Danilo Gubian and Stephane Altaus from the Lausanne University Hospital (CHUV) mechanical workshop for helping create the hypoxic setup. The authors would also like to thank Diane Macabrey and Melanie Sipion for their help with training the animals.

DISCLOSURES:

The authors have nothing to disclose.

REFERENCES:

1. Calbet, J.A. et al. Determinants of maximal oxygen uptake in severe acute hypoxia. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*. **284** (2), R291–R303

529 (2003).

530 2. Hanada, A., Sander, M., González-Alonso, J. Human skeletal muscle sympathetic nerve activity,
 531 heart rate and limb haemodynamics with reduced blood oxygenation and exercise. *The Journal*
 532 *of Physiology*. **551** (Pt 2), 635–647 (2003).

533 3. Casey, D.P., Joyner, M.J. Compensatory vasodilatation during hypoxic exercise: mechanisms
 534 responsible for matching oxygen supply to demand. *The Journal of Physiology*. **590** (24), 6321–
 535 6326 (2012).

536 4. Casey, D.P. et al. Nitric oxide contributes to the augmented vasodilatation during hypoxic
 537 exercise. *The Journal of Physiology*. **588** (Pt 2), 373–385 (2010).

538 5. Girard, O., Brocherie, F., Millet, G.P. Effects of Altitude/Hypoxia on Single- and Multiple-Sprint
 539 Performance: A Comprehensive Review. *Sports Medicine*. **47** (10), 1931–1949 (2017).

540 6. Faiss, R., Girard, O., Millet, G.P. Advancing hypoxic training in team sports: from intermittent
 541 hypoxic training to repeated sprint training in hypoxia. *British Journal of Sports Medicine*. **47**
 542 **Suppl 1**, i45-50 (2013).

543 7. Brocherie, F., Girard, O., Faiss, R., Millet, G.P. Effects of Repeated-Sprint Training in Hypoxia on
 544 Sea-Level Performance: A Meta-Analysis. *Sports Medicine* (Auckland, N.Z.). **47** (8), 1651–1660
 545 (2017).

546 8. Brocherie, F. et al. Repeated maximal-intensity hypoxic exercise superimposed to hypoxic
 547 residence boosts skeletal muscle transcriptional responses in elite team-sport athletes. *Acta*
 548 *Physiologica*. **222** (1), e12851 (2018).

549 9. Pellegrin, M. et al. New insights into the vascular mechanisms underlying the beneficial effect
 550 of swimming training on the endothelial vasodilator function in apolipoprotein E-deficient mice.
 551 *Atherosclerosis*. **190** (1), 35–42 (2007).

552 10. Picard, M. et al. Acute exercise remodels mitochondrial membrane interactions in mouse
 553 skeletal muscle. *Journal of Applied Physiology*. **115** (10), 1562–1571 (2013).

554 11. Ayachi, M., Niel, R., Momken, I., Billat, V.L., Mille-Hamard, L. Validation of a Ramp Running
 555 Protocol for Determination of the True VO₂max in Mice. *Frontiers in Physiology*. **7** (2016).

556 12. Pellegrin, M. et al. Running Exercise and Angiotensin II Type I Receptor Blocker Telmisartan
 557 Are Equally Effective in Preventing Angiotensin II-Mediated Vulnerable Atherosclerotic Lesions.
 558 *Journal of Cardiovascular Pharmacology and Therapeutics*. **22** (2) (2016).

559 13. Semin, I., Acikgöz, O., Gönenc, S. Antioxidant enzyme levels in intestinal and renal tissues
 560 after a 60-minute exercise in untrained mice. *Acta Physiologica Hungarica*. **88** (1), 55–62 (2001).

561 14. Cho, J. et al. Treadmill Running Reverses Cognitive Declines due to Alzheimer Disease:
 562 *Medicine & Science in Sports & Exercise*. **47** (9), 1814–1824 (2015).

563 15. Schill, K.E. et al. Muscle damage, metabolism, and oxidative stress in mdx mice: Impact of
 564 aerobic running. *Muscle & Nerve*. **54** (1), 110–117 (2016).

565 16. Cho, J., Kim, S., Lee, S., Kang, H. Effect of Training Intensity on Nonalcoholic Fatty Liver
 566 Disease. *Medicine & Science in Sports & Exercise*. **47** (8), 1624–1634 (2015).

567 17. Sabatier, M.J., Redmon, N., Schwartz, G., English, A.W. Treadmill training promotes axon
 568 regeneration in injured peripheral nerves. *Experimental Neurology*. **211** (2), 489–493 (2008).

569 18. Rolim, N. et al. Aerobic interval training reduces inducible ventricular arrhythmias in diabetic
 570 mice after myocardial infarction. *Basic Research in Cardiology*. **110** (4), 44 (2015).


571 19. JoVE Science Education Database. Lab Animal Research. Blood Withdrawal I.
 572 <https://www.jove.com/science-education/10246/blood-withdrawal-i> (2018).

20. Peyter, A.-C. et al. Muscarinic receptor M1 and phosphodiesterase 1 are key determinants in pulmonary vascular dysfunction following perinatal hypoxia in mice. *American Journal of Physiology-Lung Cellular and Molecular Physiology*. **295** (1), L201–L213 (2008).
21. Faiss, R. et al. Significant Molecular and Systemic Adaptations after Repeated Sprint Training in Hypoxia. *PLOS ONE*. **8** (2) (2013).
22. Faiss, R. et al. Repeated Double-Poling Sprint Training in Hypoxia by Competitive Cross-country Skiers. *Medicine & Science in Sports & Exercise*. **47** (4), 809–817 (2015).
23. Billat, V.L., Mouisel, E., Roblot, N., Melki, J. Inter- and intrastrain variation in mouse critical running speed. *Journal of Applied Physiology*. **98** (4), 1258–1263 (2005).
24. Ferguson, S.K. et al. Effects of living at moderate altitude on pulmonary vascular function and exercise capacity in mice with sickle cell anemia. *The Journal of Physiology* (2018).
25. Lightfoot, J.T., Turner, M.J., Debate, K.A., Kleeberger, S.R. Interstrain variation in murine aerobic capacity. *Medicine & Science in Sports & Exercise*. **33** (12) (2001).
26. Wojewoda, M. et al. Running Performance at High Running Velocities Is Impaired but V'O₂max and Peripheral Endothelial Function Are Preserved in IL-6–/– Mice. *PLOS ONE*. **9** (2) (2014).
27. Muller, C.R., Américo, A.L.V., Fiorino, P., Evangelista, F.S. Aerobic exercise training prevents kidney lipid deposition in mice fed a cafeteria diet. *Life Sciences*. **211**, 140–146 (2018).
28. Petrosino, J.M. et al. Graded Maximal Exercise Testing to Assess Mouse Cardio-Metabolic Phenotypes. *PLOS ONE*. **11** (2), e0148010 (2016).
29. Poole, D.C., Jones, A.M. Oxygen Uptake Kinetics. *Comprehensive Physiology* (2012).
30. Copp, S.W., Hirai, D.M., Musch, T.I., Poole, D.C. Critical speed in the rat: implications for hindlimb muscle blood flow distribution and fibre recruitment. *The Journal of Physiology*. **588** (Pt 24), 5077–5087 (2010).
31. Kregel, K. et al. *Resource Book for the Design of Animal Exercise Protocols*. American Physiological Society (2006).
32. Lamy, S. et al. Air puffs as refinement of electric shocks for stimulation during treadmill exercise test. *The FASEB Journal*. **30** (1 Supplement), 1014.5 (2016).
33. Koenen, K. et al. Sprint Interval Training Induces A Sexual Dimorphism but does not Improve Peak Bone Mass in Young and Healthy Mice. *Scientific Reports*. **7** (2017).





All-out 10 s Sprints

20 s recovery



Aorta
Pulmonary
Artery





1

2

3


4

5

Figure

[Click here to access/download;Figure;Figure 5-Mice Weights overtime .pdf](#)

Weight (g)



- LOWN
- LowH
- SupraN
- SupraH

Figure 6





Figure 7



Animal Welfare Monitoring Sheet

Authorization n°

Experimentators:

Baseline

	Date	24/07	31/07
Body weight baseline:	Day	01	08
ANIMAL ID:	Score	0	0
Types of interventions:		/	/
APPEARANCE			
Normal	0	X	X
General lack of grooming	1		
Piloerection	2		
Hunched posture	3		
NATURAL BEHAVIOUR			
Normal	0	X	X
Less mobile and/or isolated	2		
Vocalization, self mutilation, restless or still	3		
BODY WEIGHT			
% of body weight change compared to mean body weight of the control group			
0-5% weight loss	0	X	X
6-10% weight loss	1		
11-15% weight loss	2		
>15% weight loss	3		
TOTAL SCORE		0	0
Signature (initials)			

CRITERIA FOR MICE EUTHANASIAAny mice with a total combined score ≥ 5

Any mice with one or more individual maximum score of 3

MidExp					End of Exp				
03/08	07/08	10/08	14/08	17/08	21/08	24/08	28/08	31/08	04/09
12	15	19	22	25	29	32	36	39	43
0	0	0	1	0	1	1	1	1	1
/	(1)	/	/	/	/	/	/	/	(3)
									X
X	X	X	X	X	X	X	X	X	X
X	X	X	X	X	X	X	X	X	X
X	X	X		X					
			X		X	X	X	X	X
0	0	0	1	0	1	1	1	1	1

Type of interventions:

- (1) Tail vein blood collection
- (2) Forced treadmill running test
- (3) Cardiac puncture blood collection

Name of Material/ Equipment	Company	Catalog Number	Comments/Description
Cotton swab	Q-tip		
	LSI Swissgas, Geneva, Switzerland		Gas-flow: 10 L/min and 1 L/min for O ₂ and CO ₂ , respectively
Gas mixer Sonimix 7100			Made in Plexiglas
Hypoxic Box	Homemade		
Motorized rodents treadmill Panlab LE-871	Bioseb, France		
	GREISINGER electronic Gmbh, Regenstauf, Germany		
Oximeter Greisinger GOX 100			
Sedacom software	Bioseb, France PowerLab/8SP;		
Strain gauge	ADInstruments		

ARTICLE AND VIDEO LICENSE AGREEMENT

Title of Article:

High intensity hypoxic exercise in mice: feasibility and vascular function assessment.

Author(s):

Jessica Lavier, Manon Beaumann, Steeve Menetrey, Lucia Mazzolai,
Anne-Christine Peyter, Maxime Pellegrin, Grégoire P. Millet

Item 1: The Author elects to have the Materials be made available (as described at <http://www.jove.com/publish>) via:



Standard Access



Open Access

Item 2: Please select one of the following items:



The Author is **NOT** a United States government employee.



The Author is a United States government employee and the Materials were prepared in the course of his or her duties as a United States government employee.



The Author is a United States government employee but the Materials were NOT prepared in the course of his or her duties as a United States government employee.

ARTICLE AND VIDEO LICENSE AGREEMENT

1. **Defined Terms.** As used in this Article and Video License Agreement, the following terms shall have the following meanings: **"Agreement"** means this Article and Video License Agreement; **"Article"** means the article specified on the last page of this Agreement, including any associated materials such as texts, figures, tables, artwork, abstracts, or summaries contained therein; **"Author"** means the author who is a signatory to this Agreement; **"Collective Work"** means a work, such as a periodical issue, anthology or encyclopedia, in which the Materials in their entirety in unmodified form, along with a number of other contributions, constituting separate and independent works in themselves, are assembled into a collective whole; **"CRC License"** means the Creative Commons Attribution-Non Commercial-No Derivs 3.0 Unported Agreement, the terms and conditions of which can be found at: <http://creativecommons.org/licenses/by-nc-nd/3.0/legalcode>; **"Derivative Work"** means a work based upon the Materials or upon the Materials and other pre-existing works, such as a translation, musical arrangement, dramatization, fictionalization, motion picture version, sound recording, art reproduction, abridgment, condensation, or any other form in which the Materials may be recast, transformed, or adapted; **"Institution"** means the institution, listed on the last page of this Agreement, by which the Author was employed at the time of the creation of the Materials; **"JoVE"** means MyJoVE Corporation, a Massachusetts corporation and the publisher of The Journal of Visualized Experiments; **"Materials"** means the Article and / or the Video; **"Parties"** means the Author and JoVE; **"Video"** means any video(s) made by the Author, alone or in conjunction with any other parties, or by JoVE or its affiliates or agents, individually or in collaboration with the Author or any other parties, incorporating all or any portion

of the Article, and in which the Author may or may not appear.

2. **Background.** The Author, who is the author of the Article, in order to ensure the dissemination and protection of the Article, desires to have the JoVE publish the Article and create and transmit videos based on the Article. In furtherance of such goals, the Parties desire to memorialize in this Agreement the respective rights of each Party in and to the Article and the Video.

3. **Grant of Rights in Article.** In consideration of JoVE agreeing to publish the Article, the Author hereby grants to JoVE, subject to **Sections 4** and **7** below, the exclusive, royalty-free, perpetual (for the full term of copyright in the Article, including any extensions thereto) license (a) to publish, reproduce, distribute, display and store the Article in all forms, formats and media whether now known or hereafter developed (including without limitation in print, digital and electronic form) throughout the world, (b) to translate the Article into other languages, create adaptations, summaries or extracts of the Article or other Derivative Works (including, without limitation, the Video) or Collective Works based on all or any portion of the Article and exercise all of the rights set forth in (a) above in such translations, adaptations, summaries, extracts, Derivative Works or Collective Works and (c) to license others to do any or all of the above. The foregoing rights may be exercised in all media and formats, whether now known or hereafter devised, and include the right to make such modifications as are technically necessary to exercise the rights in other media and formats. If the "Open Access" box has been checked in **Item 1** above, JoVE and the Author hereby grant to the public all such rights in the Article as provided in, but subject to all limitations and requirements set forth in, the CRC License.

ARTICLE AND VIDEO LICENSE AGREEMENT

4. **Retention of Rights in Article.** Notwithstanding the exclusive license granted to JoVE in **Section 3** above, the Author shall, with respect to the Article, retain the non-exclusive right to use all or part of the Article for the non-commercial purpose of giving lectures, presentations or teaching classes, and to post a copy of the Article on the Institution's website or the Author's personal website, in each case provided that a link to the Article on the JoVE website is provided and notice of JoVE's copyright in the Article is included. All non-copyright intellectual property rights in and to the Article, such as patent rights, shall remain with the Author.

5. **Grant of Rights in Video – Standard Access.** This **Section 5** applies if the "Standard Access" box has been checked in **Item 1** above or if no box has been checked in **Item 1** above. In consideration of JoVE agreeing to produce, display or otherwise assist with the Video, the Author hereby acknowledges and agrees that, Subject to **Section 7** below, JoVE is and shall be the sole and exclusive owner of all rights of any nature, including, without limitation, all copyrights, in and to the Video. To the extent that, by law, the Author is deemed, now or at any time in the future, to have any rights of any nature in or to the Video, the Author hereby disclaims all such rights and transfers all such rights to JoVE.

6. **Grant of Rights in Video – Open Access.** This **Section 6** applies only if the "Open Access" box has been checked in **Item 1** above. In consideration of JoVE agreeing to produce, display or otherwise assist with the Video, the Author hereby grants to JoVE, subject to **Section 7** below, the exclusive, royalty-free, perpetual (for the full term of copyright in the Article, including any extensions thereto) license (a) to publish, reproduce, distribute, display and store the Video in all forms, formats and media whether now known or hereafter developed (including without limitation in print, digital and electronic form) throughout the world, (b) to translate the Video into other languages, create adaptations, summaries or extracts of the Video or other Derivative Works or Collective Works based on all or any portion of the Video and exercise all of the rights set forth in (a) above in such translations, adaptations, summaries, extracts, Derivative Works or Collective Works and (c) to license others to do any or all of the above. The foregoing rights may be exercised in all media and formats, whether now known or hereafter devised, and include the right to make such modifications as are technically necessary to exercise the rights in other media and formats. For any Video to which this **Section 6** is applicable, JoVE and the Author hereby grant to the public all such rights in the Video as provided in, but subject to all limitations and requirements set forth in, the CRC License.

7. **Government Employees.** If the Author is a United States government employee and the Article was prepared in the course of his or her duties as a United States government employee, as indicated in **Item 2** above, and any of the licenses or grants granted by the Author hereunder exceed the scope of the 17 U.S.C. 403, then the rights granted hereunder shall be limited to the maximum

rights permitted under such statute. In such case, all provisions contained herein that are not in conflict with such statute shall remain in full force and effect, and all provisions contained herein that do so conflict shall be deemed to be amended so as to provide to JoVE the maximum rights permissible within such statute.

8. **Protection of the Work.** The Author(s) authorize JoVE to take steps in the Author(s) name and on their behalf if JoVE believes some third party could be infringing or might infringe the copyright of either the Author's Article and/or Video.

9. **Likeness, Privacy, Personality.** The Author hereby grants JoVE the right to use the Author's name, voice, likeness, picture, photograph, image, biography and performance in any way, commercial or otherwise, in connection with the Materials and the sale, promotion and distribution thereof. The Author hereby waives any and all rights he or she may have, relating to his or her appearance in the Video or otherwise relating to the Materials, under all applicable privacy, likeness, personality or similar laws.

10. **Author Warranties.** The Author represents and warrants that the Article is original, that it has not been published, that the copyright interest is owned by the Author (or, if more than one author is listed at the beginning of this Agreement, by such authors collectively) and has not been assigned, licensed, or otherwise transferred to any other party. The Author represents and warrants that the author(s) listed at the top of this Agreement are the only authors of the Materials. If more than one author is listed at the top of this Agreement and if any such author has not entered into a separate Article and Video License Agreement with JoVE relating to the Materials, the Author represents and warrants that the Author has been authorized by each of the other such authors to execute this Agreement on his or her behalf and to bind him or her with respect to the terms of this Agreement as if each of them had been a party hereto as an Author. The Author warrants that the use, reproduction, distribution, public or private performance or display, and/or modification of all or any portion of the Materials does not and will not violate, infringe and/or misappropriate the patent, trademark, intellectual property or other rights of any third party. The Author represents and warrants that it has and will continue to comply with all government, institutional and other regulations, including, without limitation all institutional, laboratory, hospital, ethical, human and animal treatment, privacy, and all other rules, regulations, laws, procedures or guidelines, applicable to the Materials, and that all research involving human and animal subjects has been approved by the Author's relevant institutional review board.

11. **JoVE Discretion.** If the Author requests the assistance of JoVE in producing the Video in the Author's facility, the Author shall ensure that the presence of JoVE employees, agents or independent contractors is in accordance with the relevant regulations of the Author's institution. If more than one author is listed at the beginning of this Agreement, JoVE may, in its sole

ARTICLE AND VIDEO LICENSE AGREEMENT

discretion, elect not take any action with respect to the Article until such time as it has received complete, executed Article and Video License Agreements from each such author. JoVE reserves the right, in its absolute and sole discretion and without giving any reason therefore, to accept or decline any work submitted to JoVE. JoVE and its employees, agents and independent contractors shall have full, unfettered access to the facilities of the Author or of the Author's institution as necessary to make the Video, whether actually published or not. JoVE has sole discretion as to the method of making and publishing the Materials, including, without limitation, to all decisions regarding editing, lighting, filming, timing of publication, if any, length, quality, content and the like.

12. **Indemnification.** The Author agrees to indemnify JoVE and/or its successors and assigns from and against any and all claims, costs, and expenses, including attorney's fees, arising out of any breach of any warranty or other representations contained herein. The Author further agrees to indemnify and hold harmless JoVE from and against any and all claims, costs, and expenses, including attorney's fees, resulting from the breach by the Author of any representation or warranty contained herein or from allegations or instances of violation of intellectual property rights, damage to the Author's or the Author's institution's facilities, fraud, libel, defamation, research, equipment, experiments, property damage, personal injury, violations of institutional, laboratory, hospital, ethical, human and animal treatment, privacy or other rules, regulations, laws, procedures or guidelines, liabilities and other losses or damages related in any way to the submission of work to JoVE, making of videos by JoVE, or publication in JoVE or elsewhere by JoVE. The Author shall be responsible for, and shall hold JoVE harmless from, damages caused by lack of sterilization, lack of cleanliness or by contamination due to


the making of a video by JoVE its employees, agents or independent contractors. All sterilization, cleanliness or decontamination procedures shall be solely the responsibility of the Author and shall be undertaken at the Author's expense. All indemnifications provided herein shall include JoVE's attorney's fees and costs related to said losses or damages. Such indemnification and holding harmless shall include such losses or damages incurred by, or in connection with, acts or omissions of JoVE, its employees, agents or independent contractors.

13. **Fees.** To cover the cost incurred for publication, JoVE must receive payment before production and publication of the Materials. Payment is due in 21 days of invoice. Should the Materials not be published due to an editorial or production decision, these funds will be returned to the Author. Withdrawal by the Author of any submitted Materials after final peer review approval will result in a US\$1,200 fee to cover pre-production expenses incurred by JoVE. If payment is not received by the completion of filming, production and publication of the Materials will be suspended until payment is received.

14. **Transfer, Governing Law.** This Agreement may be assigned by JoVE and shall inure to the benefits of any of JoVE's successors and assignees. This Agreement shall be governed and construed by the internal laws of the Commonwealth of Massachusetts without giving effect to any conflict of law provision thereunder. This Agreement may be executed in counterparts, each of which shall be deemed an original, but all of which together shall be deemed to be one and the same agreement. A signed copy of this Agreement delivered by facsimile, e-mail or other means of electronic transmission shall be deemed to have the same legal effect as delivery of an original signed copy of this Agreement.

A signed copy of this document must be sent with all new submissions. Only one Agreement is required per submission.

CORRESPONDING AUTHOR

Name:	Grégoire Millet		
Department:	Sport Science Institute (ISSUL)		
Institution:	University of Lausanne		
Title:	Professor		
Signature:		Date:	27.08.2018

Please submit a **signed** and **dated** copy of this license by one of the following three methods:

1. Upload an electronic version on the JoVE submission site
2. Fax the document to +1.866.381.2236
3. Mail the document to JoVE / Attn: JoVE Editorial / 1 Alewife Center #200 / Cambridge, MA 02140

Editorial comments:

We thank you for reviewing our manuscript, and for providing supportive comments and suggestions that have helped us to improve the manuscript. We thank you for the consideration you have for our work. We have considered your comments and made amendments when necessary in the revised manuscript. We appreciate your further perusal of the revised manuscript. We have provided our responses to your comments that are in **bold**. *Amended sentences are in italic with the additional wordings in red.*

1. Please include typical numbers for V_{max} and sprint speed in the Protocol itself.

The following sentences were added to the manuscript:

"Note: Typically, V_{max} was 28.8±3.7 m/min." Page 5, line 190.

"Note: Typically, sprint velocity was 42.1±5.5 m/min." Page 4, line 147.

2. Please provide a protocol (or reference) for the low intensity training mentioned in the results.

The following steps were added to the protocol:

6. "Low intensity training"

6.1. Place the mice on individual lanes on the treadmill (0° inclination) and submit them to the following protocol:

6.1.1. Warm up mice for 5 min at 4.8 m/min followed by 5 min at 7.2 m/min.

6.1.2. Set the speed of the continuous running session to 40% of the previously determined V_{max}.

Note: Typically, continuous running velocity was 9.9 m/min.

6.1.3. Train mice for 40 min.

6.1.4. Perform this training 3 times per week with preferably 48 h between training sessions.

6.1.5. Use cotton swabs to encourage mice to run as a complementary method to electric shocks." Page 5, lines 205-220.

3. Figure 7: Please include a space between all values and units (e.g., KCl 100 mM).

Done.

4. Please ensure you have included all Figures in this revision (the last one only included the revised Figures 3 and 8).

Done.