

# Journal of Visualized Experiments

## Primary culture of rat adrenocortical cells and the assays of steroidogenic functions

--Manuscript Draft--

Article Type:	Invited Methods Article - JoVE Produced Video
Manuscript Number:	JoVE59016R3
Full Title:	Primary culture of rat adrenocortical cells and the assays of steroidogenic functions
Keywords:	adrenal gland, adrenocortical cell, steroidogenesis, endocrine, mineralocorticoid, glucocorticoid
Corresponding Author:	Yung-Chia Chen Kaohsiung Medical University Kaohsiung, -- select state -- TAIWAN
Corresponding Author's Institution:	Kaohsiung Medical University
Corresponding Author E-Mail:	i2485143@gmail.com
Order of Authors:	Yung-Chia Chen Bu-Miin Huang
Additional Information:	
Question	Response
Please indicate whether this article will be Standard Access or Open Access.	Standard Access (US\$2,400)
Please indicate the <b>city, state/province, and country</b> where this article will be <b>filmed</b> . Please do not use abbreviations.	Kaohsiung, Taiwan

**TITLE:**

Primary Culture of Rat Adrenocortical Cells and Assays of Steroidogenic Functions

**AUTHORS & AFFILIATIONS:**

Yung-Chia Chen<sup>1,2</sup>, Bu-Miin Huang<sup>3</sup>

<sup>1</sup>Graduate Institute of Medicine, College of Medicine, Kaohsiung Medical University, Kaohsiung, Taiwan

<sup>2</sup>Department of Anatomy, School of Medicine, Kaohsiung Medical University, Kaohsiung, Taiwan

<sup>3</sup>Department of Cell Biology and Anatomy, College of Medicine, National Cheng Kung University, Tainan, Taiwan

**Corresponding Author:**

Yung-Chia Chen (yungchia@kmu.edu.tw)

**KEYWORDS:**

adrenal gland, adrenocortical cell, steroidogenesis, endocrine, mineralocorticoid, glucocorticoid

**SUMMARY:**

The hormone secreted by the adrenal cortex is vital for animals against stress and diseases. Here, we present a protocol to culture the primary rat adrenal cells. It could be a good *in vitro* platform for investigating the mechanisms of the reagent of interest in adrenal steroidogenesis and lipid biosynthesis.

**ABSTRACT:**

The hormone which the adrenal cortex secretes is vital for animals against stress and diseases. The method described here is the procedure of primary cultured rat adrenal cells and related functional assays (immunofluorescence staining of lipid droplet surface protein, as well as corticosterone analysis). Unlike an *in vivo* model, the variation of interexperiments in adrenal monolayer cultures is less and the experimental condition is easy to control. Besides, the source of rats is also more stable than other animals, like bovine ones. There are also several human adrenal cell lines (NCI-H295, NCI-H295R, SW13, etc.) that can be used in adrenal studies. However, the steroid production of these lines will still be influenced by numerous factors, which include serum lot number, passage number, mutant/loss of distinct genes, etc. Except for lacking 17 $\alpha$ -hydroxylase, the primary culture of rat adrenocortical cells is a better and more convenient technique for studying adrenal physiology. In summary, primary rat adrenal cultures could be a good *in vitro* platform for researchers to investigate the mechanisms of the reagent of interest in the adrenal gland system.

**INTRODUCTION:**

The endocrine system is responsible for regulating physiological activities and homeostasis<sup>1</sup>. Adrenal glands located at the cranial pole of the kidneys are one of the major endocrine organs which secrete mineralocorticoids, glucocorticoids, and androgen<sup>2,3</sup>. There are two distinct parts of the adrenal gland: the cortex and the medulla. The adrenal cortex consists of three layers: the outer glomerulosa, the intermediate fasciculata, and the inner reticularis<sup>3</sup>. The zona glomerulosa

is the original secreting site of aldosterone, a mineralocorticoid, which aids in reabsorbing sodium ion and water in the kidneys<sup>3</sup>. The zona fasciculata primarily produces a basal level of glucocorticoids in normal physiological conditions<sup>3</sup>. Corticosteroids in rodents and cortisol in humans act to assist the body in coping with stress by regulating blood glucose<sup>3</sup>. To some extent, they can inhibit inflammatory responses and regulate the immune system<sup>4,5</sup>. In contrast to other mammals, mice and rats do not have a functional zona reticularis due to the lack of a 17 $\alpha$ -hydroxylase expression in the adrenal gland<sup>6,7</sup>. Thus, adrenals from mice and rats are devoid of the secretion of adrenal C-19 steroids (cortisol and adrenal androgens).

Primary cultured adrenocortical cells have been shown to be useful for investigating mechanisms controlling the adrenal physiology<sup>8</sup>. Like other steroidogenic tissues, each adrenal zone synthesizes its steroids from free cholesterol<sup>9</sup>. Upon adrenocorticotrophic hormone (ACTH) stimulation, the free cholesterol is released from breakdown lipid droplets and, thus, enhances steroid production in the fasciculata and reticularis<sup>10</sup>.

Many groups have attempted to establish stable cell lines from adrenocortical carcinomas. However, several concerns have limited the use of adrenocortical cell lines as in vitro models<sup>8</sup>. The steroid responses of cell lines may differ between passages, the lot number of serum, and the quality of the serum, etc. Moreover, commercial adrenal cell lines (Y1 and SW13) do not secrete corticosterone, making it more difficult to study adrenal steroidogenesis<sup>8</sup>. Using bovine and equine adrenals as ex vivo models may be a good choice, even though tissues from these markets may obscure some unknown risks. Compared to them, managing rat adrenal glands is easier and the source is relatively more stable and pathogen-free. Taken together, the present method described here could be used to investigate the related mechanism of corticosterone synthesis in rat adrenal fasciculata cells.

## **PROTOCOL:**

All the procedures including animal subjects have been approved by Institutional Animal Care and Use Committee of Kaohsiung Medical University.

### **1. Experimental Procedure**

1.1. Sacrifice the adult (8- to 12-week-old) female/male SD rats by CO<sub>2</sub> euthanasia.

1.2. Apply 70% ethanol and let it be absorbed into the skin, and then wipe the skin with a tissue paper.

1.3. Put the rat on a clean plastic plate in the supine position.

1.4. Lift the skin with “external” curved forceps at the midline and perform a blunt dissection by creating a “Y” shape (from the level of the thigh to the subcostal angle).

1.4.1. Rinse scissors with 70% ethanol and, then, cut the muscle, also creating a “Y” shape.

1.5. Find the left adrenal gland behind the stomach.

NOTE: The right adrenal can be found deeply behind the liver.

1.5.1. Use one pair of forceps to lift the stomach or liver and isolate the adrenals with another curved forceps.

1.6. After putting the adrenal glands into a sterile dish, carefully remove the adrenal capsule with fine forceps.

1.7. With a delicate pair of scissors, cut the adrenals into small pieces with a little bit of serum-free Dulbecco's modified Eagle's medium (DMEM)/F-12 medium.

1.8. Prepare three different pore-sized Pasteur pipets and use the first Pasteur pipet to aspirate all the adrenal pieces into a 50 mL conical tube containing medium with collagenase II (~0.5–0.8 mg/mL).

1.8.1. Gently triturate the tissue pieces about 10x. Use the rest of the pipets and repeat the above steps until all the pieces become smaller.

NOTE: The different pore sizes of the Pasteur pipets can be created by heating them with fire. The first pipet is the original size (about 1 mm), the pore size of the second one is about 0.8 mm, and the third Pasteur pipet is about 0.5 mm. During digestion, the pieces of adrenal will become smaller. Thus, via the mechanical procedure, the adrenals can be digested more completely. Incubate the tube in a water bath (at 37 °C) and shake it every 5 min, 4x in total.

1.9. After the 20 min incubation, add fresh and cool medium (sixfold volume; for example, 5 mL of tissue + 25 mL of cool DMEM/F-12 medium) containing 5% fetal bovine serum (FBS) and 2.5% horse serum to stop the enzyme effect.

1.10. Collect the cell pellets by centrifugation at 800 x g, 2x for 10 min.

1.11. Add an appropriate volume of medium to suspend the cells.

NOTE: Each gland can make about  $\sim 3 \times 10^5$  to  $\sim 4 \times 10^5$  cells.

1.12. Plate the cells in desired plates or dishes with growth medium (1:1 [v/v] mixture of DMEM and Ham's F-12 medium, supplemented with 25 mM HEPES and 1% penicillin and streptomycin) overnight (day 0) at 37 °C, in a humidified environment of 95% air and 5% CO<sub>2</sub>.

NOTE: Usually, we plate cells in 24-well plates (four glands/plate) for the corticosterone assay and in 35 mm dishes (one gland/approximately one to two dishes) for western blot analysis.

1.13. Carefully wash the cells 2x with serum-free warm DMEM/F-12 medium (day 1). Replace the



new growth medium.

1.14. Maintain the cells at 37 °C in a humidified environment of 95% air and 5% CO<sub>2</sub> until day 3.

#### REPRESENTATIVE RESULTS:

Using the procedure described here, primary cultured adrenal cells could be distinguished under a phase-contrast microscope (**Figure 1A**). To further confirm that the vesicles within the cytoplasm are lipid droplets, the immunofluorescence staining of adipose differentiation-related protein (ADRP) could be conducted on day-3 cultures (**Figure 1B**). Cells were grown on the glass coverslips. The cells were washed three times with phosphate-buffered saline (PBS) and fixed with 4% paraformaldehyde. After washing three times with PBS, the cells were blocked and permeabilized with 0.1% Triton X-100 in 5% nonfat milk. The primary ADRP antibody was diluted (1:50) in 5% nonfat milk and was incubated overnight. After three times of PBS washing, the cells were incubated with goat anti-rabbit Alexa Fluor 488 secondary antibodies (1:100). After repeating the washing steps, the cells were mounted with ProLong Gold Antifade mounting medium and were examined under the fluorescence microscope. Moreover, the hormone-producing ability of the rat adrenal cells could be determined by corticosterone assay. Stimulation of ACTH on rat adrenal cells could serve as a positive control (**Figure 1C**). The collected media were diluted and assayed for corticosterone content, according to the procedure described by Chen et al.<sup>11</sup>.

#### FIGURE AND TABLE LEGENDS:

**Figure 1: The characterization of adrenal cells and the corticosterone response to ACTH treatment.** (A) Phase-contrast picture of day-3 primary cultured rat adrenal cells. (B) ADRP staining is shown. (C) Day-3 cultured rat adrenal cells were treated with ACTH (1 pM and 1 nM) for 6 h. The media were collected and stored in a -80 °C refrigerator until use. Corticosterone levels were measured by enzyme-linked immunosorbent assay (ELISA).

#### DISCUSSION:

Adrenal glands play a key role in environmental adaptation<sup>3</sup>. The hormone by which the adrenal cortex secretes can regulate and adjust physiological functions and homeostasis<sup>3</sup>. The *in vivo* model reflects the real physiological effects in the body. However, it is still influenced by many complex factors, causing unstable effects in experiments. In contrast, the *in vitro* cell model has advantages which makes it incomparable to the animal model. First, the environmental condition is easy to control in the *in vitro* system. Hormones are likely to be affected by any stimulus. *In vivo* experiments usually result in a nonspecific reaction because animals are prone to distress. Second, the substances are not easy to monitor in the body but are liable in the cultured system. Third, the *in vitro* system reduces the variances between each independent experiment. Many scientific reports show a culture media of adrenocortical cells containing 10%–20% FBS<sup>12–15</sup>, whereas the protocol presented here uses 5% FBS and 2.5% horse serum to maintain the growth of adrenal cells for up to one week.

Steroidogenesis is started from the cleavage of free cholesterol by cytochrome P450 side-chain

cleavage enzymes. As shown in **Figure 1A,B**, adrenal cells exhibit numerous and different sizes of lipid droplets, suggesting it is a good tool for the investigation of steroid biosynthesis and/or lipid biology (fasciculata cells have prominent and larger lipid droplets)<sup>1,3</sup>. The ADRP, also known as perilipin 2, is a lipid droplet surface protein which assists in the storage of neutral lipids within the lipid droplets<sup>16</sup>. Due to adrenocortical cells exhibiting an amount of lipid, the ADRP (or other lipid droplet surface proteins, like perilipin) can be used to examine whether the tested agents affect hormone synthesis<sup>16</sup>.

The growth and function of adrenal glands are tightly controlled by hypothalamic corticotrophic-releasing hormone and pituitary ACTH, neuropeptides, neurotransmitters, and growth factors, etc.<sup>17</sup>. **Figure 1C** reveals that the corticosterone production was significantly induced by ACTH treatment in a concentration-dependent fashion. Moreover, ACTH at 1 pM and 1 nM stimulated the corticosterone production about 30- and 70-fold, respectively. It has been demonstrated that day-3 cultured primary adrenal cells exhibit a better response to ACTH stimulation<sup>15</sup>. This is also the reason why day-3 culture was used in the present study. In addition, the measurement of corticosterone by ELISA kit is also a simple and easy tool for research. Therefore, the primary cultured rat adrenal cells not only provide the platform for studying adrenal steroidogenesis but may also apply to the investigation of inflammatory-related diseases.

The critical steps in the present protocol are the digestion of the adrenals and control of contamination. Therefore, control of the digestion time, physical suction force, and observation of the tissue digestion level are very important for cell survival. Furthermore, the method of cutting the skin and isolating the adrenals is also significant. Always immerse all the instruments in 70% ethanol to prevent any contamination by animal fur. In order to obtain more cells, the suction frequency should be controlled less than ten times during each suction. Also, if the condition of the tissues is not good for further steps, the incubation in collagenase-containing medium should not be longer than 30 min.

Because the adrenal glands of rats are small, the glomerulosa layer might be lost during the peeling of the adrenal capsule. It has been demonstrated that decapsulation resulted in the removal of the connective tissue capsule and the entire zona glomerulosa<sup>18</sup>. The adrenal glands of male rats are smaller than those of female rats. Additionally, the lipid droplets appear to be larger in 11-week-old female or male mice<sup>6</sup>. Regarding the available cells, female rats may be a good choice for experiments.

#### **ACKNOWLEDGMENTS:**

This work was supported by a grant from the National Science Council Taiwan (NSC102-2320-B-037-008-MY3) and a National Cheng-Kung University Hospital Research Grant (NCKUH-10102045).

#### **DISCLOSURES:**

The authors have nothing to disclose.

#### **REFERENCES:**

1. Frith, C. H. Histology, Adrenal Gland, Mouse. In *Endocrine System. Monographs on Pathology of Laboratory Animals*. Edited by Jones, T. C., Mohr, U., Hunt, R. D., Capen, C. C., 8-12, Springer. Berlin, Heidelberg (1983).
2. Keegan, C. E., Hammer, G. D. Recent insights into organogenesis of the adrenal cortex. *Trends in Endocrinology Metabolism*. **13** (5), 200-208 (2002).
3. Marieb, E., Wilhelm, P. B., Mallatt, J. Chapter 17: The Endocrine System. 558-581 (2017).
4. Chrousos, G. P. The hypothalamic-pituitary-adrenal axis and immune-mediated inflammation. *New England Journal Medicine*. **332** (20), 1351-1362 (1995).
5. Bornstein, S. R., Rutkowski, H., Vrezas, I. Cytokines and steroidogenesis. *Molecular and Cellular Endocrinology*. **215** (1-2), 135-141 (2004).
6. Bielohuby, M. et al. Growth analysis of the mouse adrenal gland from weaning to adulthood: time- and gender-dependent alterations of cell size and number in the cortical compartment. *American Journal of Physiology-Endocrinology Metabolism*. **293** (1), E139-146 (2007).
7. van Weerden, W. M., Bierings, H. G., van Steenbrugge, G. J., de Jong, F. H., Schroder, F. H. Adrenal glands of mouse and rat do not synthesize androgens. *Life Sciences*. **50** (12), 857-861 (1992).
8. Wang, T., Rainey, W. E. Human adrenocortical carcinoma cell lines. *Molecular and Cellular Endocrinology*. **351** (1), 58-65 (2012).
9. Payne, A. H., Hales, D. B. Overview of steroidogenic enzymes in the pathway from cholesterol to active steroid hormones. *Endocrine Review*. **25** (6), 947-970 (2004).
10. Ruggiero, C., Lalli, E. Impact of ACTH Signaling on Transcriptional Regulation of Steroidogenic Genes. *Frontiers in Endocrinology*. **7**, 24 (2016).
11. Chen, Y. C., Liang, Y. L., Huang, Y. L., Huang, B. M. Mechanism of *Toona sinensis*-stimulated adrenal steroidogenesis in primary rat adrenal cells. *Journal of Functional Foods*. **14** (2015), 318-323 (2015).
12. Rybak, S. M., Ramachandran, J. Primary culture of normal rat adrenocortical cells. I. Culture conditions for optimal growth and function. *In Vitro*. **17** (7), 599-604 (1981).
13. Rybak, S. M., Ramachandran, J. Primary culture of normal rat adrenocortical cells. II. Quantitation of steroid dehydrogenase stain. *In Vitro*. **17** (7), 605-611 (1981).
14. O'Hare, M. J., Neville, A. M. Steroid metabolism by adult rat adrenocortical cells in monolayer culture. *Journal of Endocrinology*. **58** (3), 447-462 (1973).
15. Di Blasio, A. M., Fujii, D. K., Yamamoto, M., Martin, M. C., Jaffe, R. B. Maintenance of cell proliferation and steroidogenesis in cultured human fetal adrenal cells chronically exposed to adrenocorticotrophic hormone: rationalization of in vitro and in vivo findings. *Biology of Reproduction*. **42** (4), 683-691 (1990).
16. Brasaemle, D. L. Thematic review series: adipocyte biology. The perilipin family of structural lipid droplet proteins: stabilization of lipid droplets and control of lipolysis. *Journal of Lipid Research*. **48** (12), 2547-2559 (2007).
17. Hoeflich, A., Bielohuby, M. Mechanisms of adrenal gland growth: signal integration by extracellular signal regulated kinases1/2. *Journal of Molecular Endocrinology*. **42** (3), 191-203 (2009).
18. O'Hare, M. J., Neville, A. M. Morphological responses to corticotrophin and cyclic AMP by adult rat adrenocortical cells in monolayer culture. *Journal of Endocrinology*. **56** (3), 529-536 (1973).






Figure 1

Name of Material/ Equipment	Company
female/ male, SD/Wistar rats (8-12 wk)	BioLASCO
collagenase, type II	Sigma
DMEM	ThermoFisher
Ham's F12	ThermoFisher
HEPES	JT.baker
NaHCO <sub>3</sub>	JT.baker
Horse serum	Hyclone
Fetal bovine serum	SAFC
penicillin (10,000 U)-streptomycin (10,000 ug/ml)	Corning
curved forceps	BRAUN
forceps	BRAUN
scissor	BRAUN
delicate scissor	BRAUN
adipose-differentiation related protein	Abcam
corticosterone ELISA kit	cayman
inverted light microscopy	Leica
fluorescence microscope	Nikon
Triton X-100 goat anti-rabbit IgG Alexa 488 Fluor secondary antibody	JT.baker  ThermoFisher
anti-ADFP	Abcam

ProLong Gold antifade mountant

ThermoFisher

## Catalog Number

C-6885

12800-017

21700-075

4018

3506-1

16050014

12103C

30-002-CI

BD343R

BD331R

BC224R

BC101R

ab108323

500655

X198-07

A-11034

108323



P36935

### **Comments/Description**

male or female rats are suitable for experiments.  
Female's adrenal glands are larger than male rats,  
however, female rats own the oestrous cycles that may  
influence the experiments

isolation of adrenal cells

powder media for adrenal cells

powder media for adrenal cells

buffer (powder)

antibiotics

cut rat skin and muscle

cut adrenal glands

antibody for lipid droplet associated protein

assay for corticosterone synthesis

for immunofluorescence staining

for immunofluorescence staining

for immunofluorescence staining

for immunofluorescence staining



## ARTICLE AND VIDEO LICENSE AGREEMENT


Title of Article:

Primary culture of rat adrenal cells


Author(s):

Yung-chia Chen, Bu-Miin Huang

Item 1: The Author elects to have the Materials be made available (as described at <http://www.jove.com/publish>) via:




Standard Access




Open Access


Item 2: Please select one of the following items:



The Author is **NOT** a United States government employee.



The Author is a United States government employee and the Materials were prepared in the course of his or her duties as a United States government employee.



The Author is a United States government employee but the Materials were NOT prepared in the course of his or her duties as a United States government employee.

### ARTICLE AND VIDEO LICENSE AGREEMENT

1. **Defined Terms.** As used in this Article and Video License Agreement, the following terms shall have the following meanings: "**Agreement**" means this Article and Video License Agreement; "**Article**" means the article specified on the last page of this Agreement, including any associated materials such as texts, figures, tables, artwork, abstracts, or summaries contained therein; "**Author**" means the author who is a signatory to this Agreement; "**Collective Work**" means a work, such as a periodical issue, anthology or encyclopedia, in which the Materials in their entirety in unmodified form, along with a number of other contributions, constituting separate and independent works in themselves, are assembled into a collective whole; "**CRC License**" means the Creative Commons Attribution-Non Commercial-No Derivs 3.0 Unported Agreement, the terms and conditions of which can be found at: <http://creativecommons.org/licenses/by-nc-nd/3.0/legalcode>; "**Derivative Work**" means a work based upon the Materials or upon the Materials and other pre-existing works, such as a translation, musical arrangement, dramatization, fictionalization, motion picture version, sound recording, art reproduction, abridgment, condensation, or any other form in which the Materials may be recast, transformed, or adapted; "**Institution**" means the institution, listed on the last page of this Agreement, by which the Author was employed at the time of the creation of the Materials; "**JoVE**" means MyJove Corporation, a Massachusetts corporation and the publisher of The Journal of Visualized Experiments; "**Materials**" means the Article and / or the Video; "**Parties**" means the Author and JoVE; "**Video**" means any video(s) made by the Author, alone or in conjunction with any other parties, or by JoVE or its affiliates or agents, individually or in collaboration with the Author or any other parties, incorporating all or any portion

of the Article, and in which the Author may or may not appear.

2. **Background.** The Author, who is the author of the Article, in order to ensure the dissemination and protection of the Article, desires to have the JoVE publish the Article and create and transmit videos based on the Article. In furtherance of such goals, the Parties desire to memorialize in this Agreement the respective rights of each Party in and to the Article and the Video.

3. **Grant of Rights in Article.** In consideration of JoVE agreeing to publish the Article, the Author hereby grants to JoVE, subject to **Sections 4** and **7** below, the exclusive, royalty-free, perpetual (for the full term of copyright in the Article, including any extensions thereto) license (a) to publish, reproduce, distribute, display and store the Article in all forms, formats and media whether now known or hereafter developed (including without limitation in print, digital and electronic form) throughout the world, (b) to translate the Article into other languages, create adaptations, summaries or extracts of the Article or other Derivative Works (including, without limitation, the Video) or Collective Works based on all or any portion of the Article and exercise all of the rights set forth in (a) above in such translations, adaptations, summaries, extracts, Derivative Works or Collective Works and (c) to license others to do any or all of the above. The foregoing rights may be exercised in all media and formats, whether now known or hereafter devised, and include the right to make such modifications as are technically necessary to exercise the rights in other media and formats. If the "Open Access" box has been checked in **Item 1** above, JoVE and the Author hereby grant to the public all such rights in the Article as provided in, but subject to all limitations and requirements set forth in, the CRC License.



# ARTICLE AND VIDEO LICENSE AGREEMENT

discretion, elect not take any action with respect to the Article until such time as it has received complete, executed Article and Video License Agreements from each such author. JoVE reserves the right, in its absolute and sole discretion and without giving any reason therefore, to accept or decline any work submitted to JoVE. JoVE and its employees, agents and independent contractors shall have full, unfettered access to the facilities of the Author or of the Author's institution as necessary to make the Video, whether actually published or not. JoVE has sole discretion as to the method of making and publishing the Materials, including, without limitation, to all decisions regarding editing, lighting, filming, timing of publication, if any, length, quality, content and the like.

12. **Indemnification.** The Author agrees to indemnify JoVE and/or its successors and assigns from and against any and all claims, costs, and expenses, including attorney's fees, arising out of any breach of any warranty or other representations contained herein. The Author further agrees to indemnify and hold harmless JoVE from and against any and all claims, costs, and expenses, including attorney's fees, resulting from the breach by the Author of any representation or warranty contained herein or from allegations or instances of violation of intellectual property rights, damage to the Author's or the Author's institution's facilities, fraud, libel, defamation, research, equipment, experiments, property damage, personal injury, violations of institutional, laboratory, hospital, ethical, human and animal treatment, privacy or other rules, regulations, laws, procedures or guidelines, liabilities and other losses or damages related in any way to the submission of work to JoVE, making of videos by JoVE, or publication in JoVE or elsewhere by JoVE. The Author shall be responsible for, and shall hold JoVE harmless from, damages caused by lack of sterilization, lack of cleanliness or by contamination due to

the making of a video by JoVE its employees, agents or independent contractors. All sterilization, cleanliness or decontamination procedures shall be solely the responsibility of the Author and shall be undertaken at the Author's expense. All indemnifications provided herein shall include JoVE's attorney's fees and costs related to said losses or damages. Such indemnification and holding harmless shall include such losses or damages incurred by, or in connection with, acts or omissions of JoVE, its employees, agents or independent contractors.

13. **Fees.** To cover the cost incurred for publication, JoVE must receive payment before production and publication the Materials. Payment is due in 21 days of invoice. Should the Materials not be published due to an editorial or production decision, these funds will be returned to the Author. Withdrawal by the Author of any submitted Materials after final peer review approval will result in a US\$1,200 fee to cover pre-production expenses incurred by JoVE. If payment is not received by the completion of filming, production and publication of the Materials will be suspended until payment is received.

14. **Transfer, Governing Law.** This Agreement may be assigned by JoVE and shall inure to the benefits of any of JoVE's successors and assignees. This Agreement shall be governed and construed by the internal laws of the Commonwealth of Massachusetts without giving effect to any conflict of law provision thereunder. This Agreement may be executed in counterparts, each of which shall be deemed an original, but all of which together shall be deemed to me one and the same agreement. A signed copy of this Agreement delivered by facsimile, e-mail or other means of electronic transmission shall be deemed to have the same legal effect as delivery of an original signed copy of this Agreement.

A signed copy of this document must be sent with all new submissions. Only one Agreement is required per submission.

## CORRESPONDING AUTHOR

Name:

Yung-chia Chen

Department:

Department of Anatomy

Institution:

Kaohsiung Medical University

Title:

Associate Professor

Signature:

Yung chia Chen

Date:

08/30/2018

Please submit a **signed** and **dated** copy of this license by one of the following three methods:

1. Upload an electronic version on the JoVE submission site
2. Fax the document to +1.866.381.2236
3. Mail the document to JoVE / Attn: JoVE Editorial / 1 Alewife Center #200 / Cambridge, MA 02140



Dear editor:

We are grateful for the opportunity to revise our manuscript with the manuscript ID: JoVE59016. We deeply appreciate reviewers' time, efforts, and helpful remarks on our manuscript. We have revised the manuscript based on the reviewers' suggestions and editor's detailed comments, and include a separate document with point-by-point responses to the comments, questions, and concerns of the reviewers'. All changes in the manuscript are in red and underlined. We hope that you, the reviewers, and the editorial board now feel our paper is worthy of publication in Journal of Visualized Experiments.

Sincerely yours

Yung-Chia Chen  
Associate Professor  
Kaohsiung Medical University

#### Editorial comments

1. Please take this opportunity to thoroughly proofread the manuscript to ensure that there are no spelling or grammar issues.

A: A: Thanks for reviewer's suggestion! We have done our best to correct all spelling or grammar errors in the manuscript.

2. Please include at least one paragraph of text to explain the Representative Results in the context of the technique you have described, e.g., how do these results show the technique, suggestions about how to analyze the outcome, etc. The paragraph text should refer to all of the figures.

A: We have revised the section of representative results according to editor's comment. Please see the following text.

#### **REPRESENTATIVE RESULTS:**

Using the procedure described here, primary cultured adrenal cells could be distinguished under phase-contrast microscope (figure 1A). To further confirm the vesicles within the cytoplasm are lipid droplets, the immunofluorescence staining of adipose-differentiated related protein (ADRP) can be conducted on Day-3 cultures (figure 1B). Cells were grown on the glass coverslips. The cells were washed with

phosphate buffered saline (PBS) three times and fixed with 4% paraformaldehyde. After washing three times with PBS, cells were blocked and permeabilized with 0.1% Triton X-100 in 5% nonfat milk. The primary ADRP antibody was diluted (1:50) in 5% nonfat milk and was incubated overnight. After 3 times of PBS washing, cells were incubated with goat anti-rabbit Alexa Fluor 488 secondary antibodies (1:100). Repeat the washing steps, the cells were then mounted with Prolong Gold antifade mounting medium and were examined under the fluorescence microscope.

Moreover, the hormone-producing ability of the rat adrenal cells can be determined by corticosterone assay. Stimulation of ACTH on rat adrenal cells could be served as a positive control (figure 1C). The collected media were diluted and assayed for corticosterone content according to the procedure described by chen et al. (2015)<sup>11</sup>.

3. Please remove all headers from Representative Results and Discussion.

A: Thanks for the kindly suggestion! The headers from representative results and discussion have been removed.

4. Please section you protocol steps. Please adjust the numbering of the Protocol to follow the JoVE Instructions for Authors. For example, 1 should be followed by 1.1 and then 1.1.1 and 1.1.2 if necessary.

A: Thanks for the kindly suggestion! We have revised the protocol section according to the editor's comments.