

# Journal of Visualized Experiments

## Initial 3D cell cluster control in a hybrid gel cube device for repeatable pattern formations

--Manuscript Draft--

<b>Article Type:</b>	Invited Methods Article - JoVE Produced Video
<b>Manuscript Number:</b>	JoVE59214R2
<b>Full Title:</b>	Initial 3D cell cluster control in a hybrid gel cube device for repeatable pattern formations
<b>Keywords:</b>	3D cell control; 3D culture; in vitro experiment; Tissue Engineering; pattern formation; airway branches
<b>Corresponding Author:</b>	Masaya Hagiwara Osaka Daigaku Daigakuin Seimei Kino Kenkyuka Sakai, Osaka JAPAN
<b>Corresponding Author's Institution:</b>	Osaka Daigaku Daigakuin Seimei Kino Kenkyuka
<b>Corresponding Author E-Mail:</b>	m-hagiwara@21c.osakafu-u.ac.jp
<b>Order of Authors:</b>	Masaya Hagiwara Rina Nobata Tomohiro Kawahara
<b>Additional Information:</b>	
<b>Question</b>	<b>Response</b>
Please indicate whether this article will be Standard Access or Open Access.	Standard Access (US\$2,400)
Please indicate the <b>city, state/province, and country</b> where this article will be <b>filmed</b> . Please do not use abbreviations.	Sakai-shi, Osaka, Japan



December 3rd, 2018.  
Dr. Nandita Singh  
Senior Science Editor  
*Journal of Visualized Experiments*

Dear Dr. Singh:

I, along with my coauthors, would like to re-submit the revised manuscript entitled “***Initial 3D cell cluster control in the Hybrid gel cube device for repeatable pattern formations***”, for publication in *Journal of Visualized Experiments*. The manuscript has been carefully rechecked and appropriate changes have been made in accordance with the reviewers’ suggestions

This manuscript has not been published or presented elsewhere and is not under consideration by another journal. We have read and understood your journal’s policies, and we believe that neither the manuscript nor the study violates any of these.

Because some of the authors’ affiliation may be changed on March 2019, we would like to finish filming by the end of February, 2019 if possible.

Thank you for your consideration. I look forward to hearing from you.

Sincerely,

M. Hagiwara, Prof.  
Nanoscience and Nanotechnology Research Center  
Osaka Prefecture University  
1-2 Gakuen-cho, Naka-ku Sakai-shi, Osaka 599-8570, Japan  
E-mail: m-hagiwara@21c.osakafu-u.ac.jp  
Tel: +81-72-254-9829

**TITLE:**

**Initial 3D Cell Cluster Control in a Hybrid Gel Cube Device for Repeatable Pattern Formations**

**AUTHORS AND AFFILIATIONS:**

Masaya Hagiwara<sup>1</sup>, Rina Nobata<sup>2</sup>, Tomohiro Kawahara<sup>3</sup>

<sup>1</sup>NanoSquare Research Institute, Osaka Prefecture University, Osaka, Japan

<sup>2</sup>Department of Bioscience and Informatics, Osaka Prefecture University, Osaka, Japan

<sup>3</sup>Department of Biological Functions Engineering, Kyushu Institute of Technology, Kitakyushu, Japan

**Corresponding Author:**

Masaya Hagiwara (m-hagiwara@21c.osakafu-u.ac.jp)

**Email Addresses of Co-authors:**

Rina Nobata (sxc04074@edu.osakafu-u.ac.jp)

Tomohiro Kawahara (kawahara@lsse.kyutech.ac.jp)

**KEYWORDS:**

3D control, 3D culture, in vitro experiment, pattern formation, airway branches, multi-directional imaging

**SUMMARY:**

We present a procedure for controlling the initial cell cluster shape in a 3D extracellular matrix to obtain a repeatable pattern formation. A cubic device containing two different hydrogels is employed to achieve multi-directional imaging for tissue pattern formation.

**ABSTRACT:**

The importance of in vitro 3D cultures is considerably emphasized in cell/tissue culture. However, the lack of experimental repeatability is one of its restrictions. Producing few repeatable results of pattern formation deteriorates the analysis of the mechanisms underlying the self-organization. Reducing variation in initial culture conditions, such as the cell density and distribution in the extracellular matrix (ECM), is crucial to enhance the repeatability of a 3D culture. In this article, we demonstrate a simple but robust procedure for controlling the initial cell cluster shape in a 3D extracellular matrix to obtain highly repeatable pattern formations. A micromold with a desired shape was fabricated by using photolithography or a machining process, and it formed a 3D pocket in the ECM contained in a hybrid gel cube (HGC). Highly concentrated cells were then injected in the pocket so that the cell cluster shape matched with the fabricated mold shape. The employed HGC allowed multi-directional scanning by its rotation, which enabled high-resolution imaging and the capture of the entire tissue structure even though a low-magnification lens was used. Normal human bronchial epithelial cells were used to demonstrate the methodology.

**INTRODUCTION:**

The importance of a 3D culture, which better mimics biological environments than does a 2D culture, is considerably emphasized in cell/tissue culture<sup>1-3</sup>. The interaction between the cells and extracellular matrix (ECM) provides important cues regarding morphogenesis<sup>4,5</sup>. Many tissue formations can emerge only under 3D environments, such as the folding process<sup>6,7</sup>, invagination<sup>8</sup>, and tubular formation<sup>9,10</sup>. However, numerous difficulties prevent researchers from shifting to 3D experiments from 2D experiments on a dish. One of the major difficulties in 3D experiments is the issue of imaging 3D samples. Compared with planar experiments, acquisition of appropriate 3D images is still challenging in many cases. In particular, obtaining an appropriate 3D image is a difficult task when the sample size reaches the millimeter range owing to the large focal depth of low-magnification lenses. For example, the focal depth reaches more than 50  $\mu\text{m}$  when a 10x magnification lens is used while the size of the single cell is normally less than 10  $\mu\text{m}$ . To enhance the imaging quality, high-technology microscopy systems are being developed (e.g., two-photon microscopy<sup>11</sup> and light-sheet microscopy system<sup>12</sup>), but their availability is limited owing to their expensive price. As an alternative, we have previously developed a hybrid gel cube (HGC) device<sup>13</sup>. The device consists of two types of hydrogels: agarose as a support gel and an ECM such as collagen or Matrigel as a culture gel. The HGC allows us to collect the sample during culturing and rotate the cube to achieve multi-directional imaging, which addresses the focal depth problem<sup>14</sup>.

Another difficulty in 3D experiments is their low repeatability owing to the poor controllability of the 3D environments. Unlike a planar culture on a plastic dish, variations in the initial culture conditions easily occur in a 3D space surrounded by a soft material. A significant variation in the experimental results deteriorates the following analysis and masks the underlying mechanisms. Many engineering technologies have been developed to spatially align single cells, such as bioprinting<sup>15,16</sup>, fiber weaving<sup>17</sup>, and scaffolding<sup>18</sup>, but they require complex preprocessing or specifically designed equipment. In contrast, we have developed a methodology for achieving 3D cell alignment in an HGC<sup>19</sup>.

In this protocol, we illustrated a simple procedure with commonly used equipment for controlling the 3D initial cell cluster shape in an HGC. First, the fabrication process of the HGC was demonstrated. Then, micromolds fabricated by photolithography or a machining process were placed in the HGC to produce a pocket with an arbitrary shape in an ECM. Subsequently, highly dense cells after centrifugation were injected into the pocket to control the initial cell cluster shape in the HGC. The precisely controlled cell cluster could be imaged from many directions because of the HGC. Normal human bronchial epithelial (NHBE) cells were used to demonstrate the control of the initial cell cluster shape and imaging of the branches from multiple directions for enhancing the imaging quality.

## **PROTOCOL:**

### **1. Fabrication of hybrid gel cube device**

**1.1. Prepare a polycarbonate (PC) cubic frame either by using a machining process or a 3D printer. The size of the PC frame depends on the sample size. In this study, we used a frame of 5 mm on**

each side so that the branches had sufficient space to elongate during culturing.

1.2. Place the PC frame on a pre-cooled glass slide or another smooth surface in an ice box ( $<0^{\circ}\text{C}$ ) (**Figure 1A**).

1.3. Add 12  $\mu\text{L}$  of pre-heated 1.5% agarose (w/v) from the top face of the cubic frame to the bottom surface with a pipette. Stroke the pipette so that the agarose spreads to create a flat surface (**Figure 1B**).

1.4. Within minutes, the agarose will be cured. Pick up the PC frame by sliding it to the edge of the glass slide using tweezers. Note that vertically picking up the frame from the glass slide may result in detaching of the agarose from the cubic frame owing to the adhesive force between the agarose and the glass slide (**Figure 1C**).

1.5. Rotate the cubic frame to make the open face down, and then place on the glass slide again (**Figure 1D**).

1.6. Repeat steps 1.3–1.5 until three surfaces are filled with agarose (**Figure 1E**).

1.7. To form an agarose wall on the fourth and fifth faces, drop the agarose in from an open face (**Figure 1F**). Once an agarose wall is formed on the fifth face, the preparation of the HGC is completed.

## 2. Fabrication of micromolds by photolithography or a machining process

### 2.1. Fabrication by photolithography

2.1.1. Clean a silicon (Si) wafer via ultrasonic cleaning with acetone, isopropyl alcohol (IPA), then distilled (DI) water for 5 min each. After nitrogen blowing the Si wafer with nitrogen, dehydrate in an oven at  $140^{\circ}\text{C}$  for 20 min (**Figure 2A(i)**).

2.1.2. Spin-coat a sacrificial layer such as LOR and Az resists on the Si wafer (2,000 rpm, 30 s) by using a spin-coater under yellow light. Immediately after spin-coating, heat the wafer on a hot plate at  $90^{\circ}\text{C}$  for 3 min (**Figure 2A(ii)**). The desired thickness of the sacrificial layer is more than  $1\text{ }\mu\text{m}$  so that the lift off process can be carried out easily, but an accurate thickness is not necessary.

2.1.3. Place the wafer on a hot plate at  $70^{\circ}\text{C}$ , and then laminate a thick negative photoresist sheet with the desired thickness on the sacrificial layer using a hand roller (**Figure 2A(iii)**)

2.1.4. Five minutes after lamination, place the wafer on a mask aligner for UV exposure. Set a photomask with the desired pattern on the sheet resist and expose with UV light for an appropriate time duration. The exposure time is determined by the thickness of photoresist and the intensity of the UV light. Subsequently, place the wafer on a hot plate for a post-exposure

bake at 60 °C for 5 min, followed by baking at 90 °C for 10 min. (**Figure 2A(iv)**)

2.1.5. Develop a negative photoresist by immersing the wafer in propylene glycol monomethyl ether acetate (PGMEA) until the unexposed area is completely dissolved (approximately 20 min) (**Figure 2A(v)**).

2.1.6. Dissolve the sacrificial layer by using IPA for ultrasonic cleaning to lift off the microstructure. Once the microstructure is peeled off from the wafer, rinse with DI water by ultrasonic cleaning for 10 min (**Figure 2A(vi)**).

2.1.7. Immerse the mold in 30  $\mu$ L of the MPC polymer for 30 min at room temperature (approximately 20 °C) to completely dry it out. **Figure 2B** shows the fabricated prism mold.

## 2.2. Fabrication of a cylinder mold

2.2.1. Fabricate a steel cylinder mold with desired size by a machining process. Alternatively, use commercially available stainless steel. (In this study, a  $\phi$  600  $\mu$ m cylinder is used.)

2.2.2. Immerse the cylinder steel in 30  $\mu$ L of the MPC polymer for 30 min at room temperature (approximately 20 °C) to completely dry it out.

2.2.3. Prepare liquid polydimethylsiloxane (PDMS) by mixing base resin and its catalyst (see **Table of Materials**) in a 10:1 ratio followed by degassing with a vacuum degassing system for 20 min. Then pour the PDMS in a 12-well-plate.

2.2.4. Set the steel cylinder perpendicular to the datum surface and fixed to a linear z-stage so that it can be perpendicularly inserted in the uncured PDMS surface in a 12-well-plate placed on the same datum surface by moving the z-stage. Incubate the PDMS with the cylinder steel in an oven for 20 min at 90 °C for curing. Cut the PDMS to form a flank surface so that the pipet can be inserted in the following process (**Figure 2C**).

## 3. Controlling initial cell cluster shape in a hydrogel

3.1. Inject an appropriate ECM for the desired cell culture, such as collagen and artificial basement membrane, into the HGC to fill the culture space (**Figure 3A**).

3.2. Set the fabricated micromold on the HGC directly or indirectly with an appropriate holder, as shown in **Figure 3B**. Subsequently, place the HGC in a CO<sub>2</sub> incubator for 25 min at 37 °C for curing it.

3.3. Carefully lift out the micromold so that the ECM does not deteriorate (**Figure 3C**). The pocket, as the desired mold shape, will be fabricated in the ECM.

3.4. Harvest cells from the cultured dish by trypsin–EDTA or its equivalent depending on the cell

types. Subsequently, centrifuge the cells to obtain highly concentrated cells (for NHBE culture, apply 0.25% trypsin-EDTA and incubate cells for 3 min, then neutralize with FBS-containing medium and centrifuging at 300 x *g* for 4 min).

3.5. After centrifuging, remove the supernatant medium to condense the cells, and inject highly concentrated cells ( $3.0 \times 10^4$  cells/ $\mu\text{L}$  for NHBE cells) into the pocket in the ECM (**Figure 3D**).

3.6. Incubate the HGC with cells in a CO<sub>2</sub> incubator for 20 min at 37 °C so that the cells fall into the ECM pocket to fill the space created by the micromolds. If excessive medium or cells are present on the top surface, carefully remove using a pipette.

3.7. Place the HGC on a 24-well-plate and add 100  $\mu\text{L}$  of appropriate medium (**Figure 3E**). For NHBE cells, use a 1:1 mixture of commercially available specialized medium for NHBE and endothelial cells (see **Table of Materials**).

3.8. Inject additional ECM to close the pocket in the ECM, and then incubate at 37 °C for 25 min.

3.9. Cool down preheated 1.5% agarose to room temperature (approximately 20 °C). Drop approximately 10  $\mu\text{L}$  of the agarose onto the top surface of the HGC to close the surface and prevent the ECM from falling out of the HGC during culturing and imaging (**Figure 3F**). Then, incubate the HGC for another 20 min at 37 °C to cure the agarose.

3.10. Add medium to cover the entire HGC so that the osmotic pressure can help in providing nutrition to the cells inside the HGC.

## 4. Multi-directional imaging

### 4.1. Non-invasive 3D shape recognition by multi-directional observation

4.1.1. Place the culture dish or well plate containing the HGC on a microscope and orient the HGC to the camera frame. Then, obtain a sample image by bright field or phase contrast.

4.1.2. Pick up and rotate the HGC to place a different surface down.

4.1.3. Repeat steps 4.1.1 and 4.1.2 until images from all six sides are obtained.

### 4.2. Immuno-fluorescent imaging by multi-directional observation

4.2.1. For fixation, apply 4% paraformaldehyde to the sample over the HGC at room temperature (approximately 20 °C) for 20 min, followed by two rinses of PBS for 10 min each.

4.2.2. Permeabilize the HGC with PBS containing 0.5% Triton X-100 for 10 min at 4 °C, then wash 3x for 10 min with PBS.

4.2.3. Incubate the HGC with 10% goat serum in IF-buffer (0.2% Triton X-100, 0.1% bovine serum albumin, and 0.05% Tween-20 in PBS) for 60 min at room temperature for a primary blocking step.

4.2.4. For staining of a specific molecule, use the appropriate antibody. To observe the collective cell geometry, Alexa Fluor 488 Phalloidin can be used to stain actin.

4.2.5. Place the HGC on a glass bottom dish under a laser or fluorescent microscope and perform scanning from six sides to obtain the entire sample image.

#### REPRESENTATIVE RESULTS:

Normal human bronchial epithelial (NHBE) cells were used to demonstrate the illustrated methodology and control the initial collective cell geometry to achieve a cylinder shape and a prism shape, respectively in an ECM environment. The multi-directional imaging results obtained by phase contrast as well as phalloidin staining of a cylinder shape (**Figure 4A,B**) and the prism shape (**Figure 4C,D**) are presented. The cells are suitably aligned to yield the desired 3D shape in the ECM environment. Subsequently, the cells were cultured for four days to form a specific shape and then were analyzed by multi-directional imaging. The large sample size makes it quite difficult to capture the entire tissue image from one direction; however, the HGC allowed imaging from up to six sides, revealing the whole tissue shape. **Figure 5** shows the immunostaining results of the bronchial tree developed from the NHBE cells. The NHBE cells were initially controlled to a cylindrical shape in the HGC. Then, most of the branches were directed perpendicular to the cylindrical axis.

#### FIGURE AND TABLE LEGENDS:

**Figure 1: Fabrication process of hybrid gel cube device.** (A) The polycarbonate cubic frame is set on a precooled ice box. (B) Agarose (1.5%) is injected to cover the bottom surface. (C) The cube is slid to be collected. (D) The cube is rotated to make another surface down. (E) Agarose (1.5%) is injected to form another agarose wall. (F) After an agarose wall is produced for three faces, agarose is injected through an open face. Scale bar: 5 mm.

**Figure 2: Fabrication process of micromolds.** (A) Fabrication process of a micromold by photolithography. (B) Complete form of a fabricated prism shape micromold by photolithography. (C) Fabricated micromold with cylinder steel. Scale bar: (B) 2 mm, (C) 10 mm.

**Figure 3: Fabrication process of initial cell cluster control.** (A) ECM (e.g., collagen and Matrigel) is injected into the HGC. (B) The micromold is set on the cubic frame with or without a holder. (C) After the ECM is cured, the micromold is removed. (D) After centrifuging, highly dense cells are injected and incubated. (E) Medium is applied. (F) Additional ECM followed by agarose are injected to cover the top surface of the HGC. Scale bar: 5 mm.

**Figure 4: Results of initial cell control with NHBE cells.** (A) Phase-contrast images of the cylindrically controlled NHBE cells taken for the bottom and lateral surface. The entire shape could be recognized by multi-directional observation without requiring fluorescent imaging. (B)



Fluorescence images of the cylindrically controlled NHBE cells staining F-actin. (C) Phase-contrast images of the NHBE cells controlled in a triangular prism shape captured from the bottom and lateral surfaces. (D) Fluorescence images of the NHBE cells controlled in a triangular prism shape. Scale bar: 100  $\mu$ m. 10x magnification lens (0.3 numerical aperture) was used.

**Figure 5: Projected fluorescent images captured by multi-directional scanning.** The NHBE cells were initially controlled to produce a cylindrical shape and cultured in the HGC for 4 days. Subsequently, actin was stained by phalloidin. The branches were elongated from the initial cylinder along the x axis, and the length, size, and angle from the axis were equally developed. Lower right image shows the schematic of the x-y-z axis and the sample shape. Scale bar: 100  $\mu$ m. A 10x magnification lens (0.3 numerical aperture) was used.

## DISCUSSION:

The method presented in this paper is simple and can be performed without high-technology equipment. Concurrently, a precise cell cluster shape control result in the 3D space of hydrogel can be obtained. After the initial control, the cells can grow in the HGC as much as they are cultured on a dish. The multi-directional imaging is performed by rotating the sample with the HGC using any microscopy system, and it significantly enhances the imaging quality. The choice of the materials for the HGC frame and micromold is flexible as long as they are biocompatible. A 3D printer may be used to produce the HGC frame or micromold if the accuracy of the 3D printer is sufficient for their applications. The proposed method is compatible with many image enhancing technologies such as transparency reagents<sup>20,21</sup> and light-sheet microscopy systems. By employing these technologies, the image quality can be improved.

The volume of the agarose, ECM, and highly dense cells to be injected depends on the HGC and mold size. A careful manual operation that does not inject air bubbles is required, otherwise the bubbles formed will deteriorate the accuracy of the control, cell growth, and imaging quality.

The fabrication process of the micromolds will be determined by the desired shape to control. Photolithography enables fabrication of precise 2D shapes with a certain depth, such as a prism shape on the order of micrometers; however, it is not effective with 3D shapes, such as a cylinder. The machining process allows us to fabricate the 3D shape, but in general, the dimensional accuracy is lower than photolithography. After the fabrication of the micromolds, coating of the 2-methacryloyloxyethyl phosphorylcholine (MPC) polymer on the molds is required to prevent adhesion on the ECM.

By taking advantage of an HGC, microscopic imaging with or without laser exposure can be performed in multiple directions. A 3D sample shape can be approximately recognized by multi-directional observation without fluorescent labeling; this method does not cause any cell invasiveness during observation. In contrast, fluorescent imaging is required for obtaining a more accurate 3D sample shape and molecular expression.

The constraint of the presented method for initial cell control is that it cannot be applied to a mold with a complex 3D shape. The mold has to be removed by picking up without deteriorating

the ECM. Thus, the mold is limited to a simple straight or tapered shape. To obtain more complex shape control, further development of the protocol is needed.

The proposed methodology has broad utility and can be easily conducted in most of the laboratory environments. This simple procedure can overcome the limitations of 3D culture, imaging, and repeatability, and contribute to the further development required for 3D culture.

#### ACKNOWLEDGMENTS:

This work was financially supported by JSPS KAKENHI (18H04765) and the Program to Disseminate Tenure Track System, MEXT, Japan.


#### DISCLOSURES:


The authors and the Osaka Prefecture University and Kyushu Institute of Technology have filed a patent application for a hybrid gel cube device, and Nippon Medical and Chemical Instruments Co. Ltd, Japan has recently commercialized the cube. The company did not affect any of the design, process, and methods described.

#### REFERENCES:

1. Baker, B.M., Chen, C.S. Deconstructing the third dimension – how 3D culture microenvironments alter cellular cues. *Journal of Cell Science*. **125** (13), 3015–3024, doi: 10.1242/jcs.079509 (2012).
2. Sasai, Y. Cytosystems dynamics in self-organization of tissue architecture. *Nature*. **493** (7432), 318–326, doi: 10.1038/nature11859 (2013).
3. Shamir, E.R., Ewald, A.J. Three-dimensional organotypic culture: Experimental models of mammalian biology and disease. *Nature Reviews Molecular Cell Biology*. **15** (10), 647–664, doi: 10.1038/nrm3873 (2014).
4. Frantz, C., Stewart, K.M., Weaver, V.M. The extracellular matrix at a glance. *Journal of Cell Science*. **123** (24), 4195–4200, doi: 10.1242/jcs.023820 (2010).
5. Choquet, D., Felsenfeld, D.P., Sheetz, M.P. Extracellular matrix rigidity causes strengthening of integrin- cytoskeleton linkages. *Cell*. **88** (1), 39–48, doi: 10.1016/S0092-8674(00)81856-5 (1997).
6. Hannezo, E., Prost, J., Joanny, J.-F. Theory of epithelial sheet morphology in three dimensions. *Proceedings of the National Academy of Sciences of the United States of America*. **111** (1), 27–32, doi: 10.1073/pnas.1312076111 (2014).
7. Tawk, M. et al. A mirror-symmetric cell division that orchestrates neuroepithelial morphogenesis. *Nature*. **446** (7137), 797–800, doi: 10.1038/nature05722 (2007).
8. Eiraku, M. et al. Self-organizing optic-cup morphogenesis in three-dimensional culture. *Nature*. **472** (7341), 51–58, doi: 10.1038/nature09941 (2011).
9. Zegers, M.M.P., O'Brien, L.E., Yu, W., Datta, A., Mostov, K.E. Epithelial polarity and tubulogenesis in vitro. *Trends in Cell Biology*. **13** (4), 169–176, doi: 10.1016/S0962-8924(03)00036-9 (2003).
10. Affolter, M., Bellusci, S., Itoh, N., Shilo, B., Thiery, J.P., Werb, Z. Tube or not tube: Remodeling epithelial tissues by branching morphogenesis. *Developmental Cell*. **4** (1), 11–18, doi: 10.1016/S1534-5807(02)00410-0 (2003).

11. Denk, W., Strickler, J., Webb, W. Two-photon laser scanning fluorescence microscopy. *Science*. **248** (4951), 73–76, doi: 10.1126/science.2321027 (1990).
12. Huisken, J. et al. Optical Sectioning Deep Inside Live Embryos by Selective Plane Illumination Microscopy. **305**, 1007–1009, doi: 10.1126/science.1100035 (2004).
13. Hagiwara, M., Kawahara, T., Nobata, R. Tissue in Cube: In Vitro 3D Culturing Platform with Hybrid Gel Cubes for Multidirectional Observations. *Advanced Healthcare Materials*. **5** (13), 1566–1571, doi: 10.1002/adhm.201670069 (2016).
14. Hagiwara, M., Nobata, R., Kawahara, T. Large Scale Imaging by Fine Spatial Alignment of Multi-Scanning Data with Gel Cube Device. *Applied Sciences*. **8** (235), doi: 10.3390/app8020235 (2018).
15. Kolesky, D.B., Homan, K.A., Skylar-Scott, M.A., Lewis, J.A. Three-dimensional bioprinting of thick vascularized tissues. *Proceedings of the National Academy of Sciences of the United States of America*. **113** (12), 3179–3184, doi: 10.1073/pnas.1521342113 (2016).
16. Mironov, V., Visconti, R.P., Kasyanov, V., Forgacs, G., Drake, C.J., Markwald, R.R. Organ printing: Tissue spheroids as building blocks. *Biomaterials*. **30** (12), 2164–2174, doi: 10.1016/j.biomaterials.2008.12.084 (2009).
17. Onoe, H. et al. Metre-long cell-laden microfibres exhibit tissue morphologies and functions. *Nature Materials*. **12** (6), 584–590, doi: 10.1038/nmat3606 (2013).
18. Miller, J.S. et al. Rapid casting of patterned vascular networks for perfusable engineered three-dimensional tissues. *Nature Materials*. **11** (9), 768–774, doi: 10.1038/nmat3357 (2012).
19. Hagiwara, M., Nobata, R., Kawahara, T. High repeatability from 3D experimental platform for quantitative analysis of cellular branch pattern formations. *Integrative Biology*. **10**, 306–312, doi: 10.1039/C8IB00032H (2018).
20. Tainaka, K. et al. Whole-body imaging with single-cell resolution by tissue decolorization. *Cell*. **159** (4), 911–924, doi: 10.1016/j.cell.2014.10.034 (2014).
21. Murray, E. et al. Simple, Scalable Proteomic Imaging for High-Dimensional Profiling of Intact Systems. *Cell*. **163** (6), 1500–1514, doi: 10.1016/j.cell.2015.11.025 (2015).










Figure 4

[Click here to access/download;Figure;Fig4.tif](#)





Figure 5

[Click here to access/download;Figure;Fig5.tif](#)





Name of Reagent/ Equipment	Company	Catalog Number	Comments/Description
12-well-plate	Corning Inc.	3513	
2-Methoxy-1-methylethyl Acetate	Pure Chemical Co.	130-10505	PGMEA, CAS: 108-65-6
4% paraformaldehyde	FUJIFILM Wako		
Agarose, low gelling temperature	Pure Chemical Co.	161-20141	CAS: 30525-89-4
BioReagent	Sigma-Aldrich	A9414	
Alexa fluor 488 phalloidin	Thermo Fisher		
AZ1512	Scientific	A12379	
BEGM bullet kit	Merck		
	Lonza	CC-3170	Specialized medium for NHBE cells
Bovine Serum Albumin solution (10 %)	Sigma-Aldrich	A1595	
EGM-2 bullet kit	Lonza	CC-3162	Specialized medium for endothelial cells
Lipidure	NOF co.		MPC polymer
Matrigel growth factor reduced basement membrane matrix	Corning Inc.	354230	
	Thermo Fisher		
Normal Goat Serum (10%)	Scientific	50197Z	
Normal human bronchial epithelial cells	Lonza	CC-2541	
SILPOT 184 W/C	Dow Corning Co.	3255981	Base resin and catalyst for PDMS
SUEX D300		DJ MicroLaminates, Inc	Thick negative photoresist (thickness: 300 $\mu$ m)
Triton X-100 (1%)	Thermo Fisher		
	Scientific	HFH10	
Trypsin-EDTA (0.25%)	Thermo Fisher		
	Scientific	25200056	
TWEEN 20	Sigma-Aldrich	P9416	



1 Alewife Center #200  
Cambridge, MA 02140  
tel. 617.945.9051  
www.jove.com

## ARTICLE AND VIDEO LICENSE AGREEMENT

Title of Article:

Author(s):

Initial 3D cell cluster control in a hybrid gel cube device for repeatable pattern formation  
Masaya Hagiwara, Rina Nobata, Tomohiro Kawahara

Item 1: The Author elects to have the Materials be made available (as described at <http://www.jove.com/publish>) via:

☒ Standard Access

☐ Open Access

Item 2: Please select one of the following items:

☒ The Author is **NOT** a United States government employee.

☐ The Author is a United States government employee and the Materials were prepared in the course of his or her duties as a United States government employee.

☐ The Author is a United States government employee but the Materials were NOT prepared in the course of his or her duties as a United States government employee.

### ARTICLE AND VIDEO LICENSE AGREEMENT

1. **Defined Terms.** As used in this Article and Video License Agreement, the following terms shall have the following meanings: **"Agreement"** means this Article and Video License Agreement; **"Article"** means the article specified on the last page of this Agreement, including any associated materials such as texts, figures, tables, artwork, abstracts, or summaries contained therein; **"Author"** means the author who is a signatory to this Agreement; **"Collective Work"** means a work, such as a periodical issue, anthology or encyclopedia, in which the Materials in their entirety in unmodified form, along with a number of other contributions, constituting separate and independent works in themselves, are assembled into a collective whole; **"CRC License"** means the Creative Commons Attribution-Non Commercial-No Derivs 3.0 Unported Agreement, the terms and conditions of which can be found at: <http://creativecommons.org/licenses/by-nc-nd/3.0/legalcode>; **"Derivative Work"** means a work based upon the Materials or upon the Materials and other pre-existing works, such as a translation, musical arrangement, dramatization, fictionalization, motion picture version, sound recording, art reproduction, abridgment, condensation, or any other form in which the Materials may be recast, transformed, or adapted; **"Institution"** means the institution, listed on the last page of this Agreement, by which the Author was employed at the time of the creation of the Materials; **"JoVE"** means MyJoVE Corporation, a Massachusetts corporation and the publisher of The Journal of Visualized Experiments; **"Materials"** means the Article and / or the Video; **"Parties"** means the Author and JoVE; **"Video"** means any video(s) made by the Author, alone or in conjunction with any other parties, or by JoVE or its affiliates or agents, individually or in collaboration with the Author or any other parties, incorporating all or any portion

of the Article, and in which the Author may or may not appear.

2. **Background.** The Author, who is the author of the Article, in order to ensure the dissemination and protection of the Article, desires to have the JoVE publish the Article and create and transmit videos based on the Article. In furtherance of such goals, the Parties desire to memorialize in this Agreement the respective rights of each Party in and to the Article and the Video.

3. **Grant of Rights in Article.** In consideration of JoVE agreeing to publish the Article, the Author hereby grants to JoVE, subject to **Sections 4 and 7** below, the exclusive, royalty-free, perpetual (for the full term of copyright in the Article, including any extensions thereto) license (a) to publish, reproduce, distribute, display and store the Article in all forms, formats and media whether now known or hereafter developed (including without limitation in print, digital and electronic form) throughout the world, (b) to translate the Article into other languages, create adaptations, summaries or extracts of the Article or other Derivative Works (including, without limitation, the Video) or Collective Works based on all or any portion of the Article and exercise all of the rights set forth in (a) above in such translations, adaptations, summaries, extracts, Derivative Works or Collective Works and (c) to license others to do any or all of the above. The foregoing rights may be exercised in all media and formats, whether now known or hereafter devised, and include the right to make such modifications as are technically necessary to exercise the rights in other media and formats. If the "Open Access" box has been checked in **Item 1** above, JoVE and the Author hereby grant to the public all such rights in the Article as provided in, but subject to all limitations and requirements set forth in, the CRC License.



## ARTICLE AND VIDEO LICENSE AGREEMENT

4. **Retention of Rights in Article.** Notwithstanding the exclusive license granted to JoVE in **Section 3** above, the Author shall, with respect to the Article, retain the non-exclusive right to use all or part of the Article for the non-commercial purpose of giving lectures, presentations or teaching classes, and to post a copy of the Article on the Institution's website or the Author's personal website, in each case provided that a link to the Article on the JoVE website is provided and notice of JoVE's copyright in the Article is included. All non-copyright intellectual property rights in and to the Article, such as patent rights, shall remain with the Author.

5. **Grant of Rights in Video – Standard Access.** This **Section 5** applies if the "Standard Access" box has been checked in **Item 1** above or if no box has been checked in **Item 1** above. In consideration of JoVE agreeing to produce, display or otherwise assist with the Video, the Author hereby acknowledges and agrees that, Subject to **Section 7** below, JoVE is and shall be the sole and exclusive owner of all rights of any nature, including, without limitation, all copyrights, in and to the Video. To the extent that, by law, the Author is deemed, now or at any time in the future, to have any rights of any nature in or to the Video, the Author hereby disclaims all such rights and transfers all such rights to JoVE.

6. **Grant of Rights in Video – Open Access.** This **Section 6** applies only if the "Open Access" box has been checked in **Item 1** above. In consideration of JoVE agreeing to produce, display or otherwise assist with the Video, the Author hereby grants to JoVE, subject to **Section 7** below, the exclusive, royalty-free, perpetual (for the full term of copyright in the Article, including any extensions thereto) license (a) to publish, reproduce, distribute, display and store the Video in all forms, formats and media whether now known or hereafter developed (including without limitation in print, digital and electronic form) throughout the world, (b) to translate the Video into other languages, create adaptations, summaries or extracts of the Video or other Derivative Works or Collective Works based on all or any portion of the Video and exercise all of the rights set forth in (a) above in such translations, adaptations, summaries, extracts, Derivative Works or Collective Works and (c) to license others to do any or all of the above. The foregoing rights may be exercised in all media and formats, whether now known or hereafter devised, and include the right to make such modifications as are technically necessary to exercise the rights in other media and formats. For any Video to which this **Section 6** is applicable, JoVE and the Author hereby grant to the public all such rights in the Video as provided in, but subject to all limitations and requirements set forth in, the CRC License.

7. **Government Employees.** If the Author is a United States government employee and the Article was prepared in the course of his or her duties as a United States government employee, as indicated in **Item 2** above, and any of the licenses or grants granted by the Author hereunder exceed the scope of the 17 U.S.C. 403, then the rights granted hereunder shall be limited to the maximum

rights permitted under such statute. In such case, all provisions contained herein that are not in conflict with such statute shall remain in full force and effect, and all provisions contained herein that do so conflict shall be deemed to be amended so as to provide to JoVE the maximum rights permissible within such statute.

8. **Protection of the Work.** The Author(s) authorize JoVE to take steps in the Author(s) name and on their behalf if JoVE believes some third party could be infringing or might infringe the copyright of either the Author's Article and/or Video.

9. **Likeness, Privacy, Personality.** The Author hereby grants JoVE the right to use the Author's name, voice, likeness, picture, photograph, image, biography and performance in any way, commercial or otherwise, in connection with the Materials and the sale, promotion and distribution thereof. The Author hereby waives any and all rights he or she may have, relating to his or her appearance in the Video or otherwise relating to the Materials, under all applicable privacy, likeness, personality or similar laws.

10. **Author Warranties.** The Author represents and warrants that the Article is original, that it has not been published, that the copyright interest is owned by the Author (or, if more than one author is listed at the beginning of this Agreement, by such authors collectively) and has not been assigned, licensed, or otherwise transferred to any other party. The Author represents and warrants that the author(s) listed at the top of this Agreement are the only authors of the Materials. If more than one author is listed at the top of this Agreement and if any such author has not entered into a separate Article and Video License Agreement with JoVE relating to the Materials, the Author represents and warrants that the Author has been authorized by each of the other such authors to execute this Agreement on his or her behalf and to bind him or her with respect to the terms of this Agreement as if each of them had been a party hereto as an Author. The Author warrants that the use, reproduction, distribution, public or private performance or display, and/or modification of all or any portion of the Materials does not and will not violate, infringe and/or misappropriate the patent, trademark, intellectual property or other rights of any third party. The Author represents and warrants that it has and will continue to comply with all government, institutional and other regulations, including, without limitation all institutional, laboratory, hospital, ethical, human and animal treatment, privacy, and all other rules, regulations, laws, procedures or guidelines, applicable to the Materials, and that all research involving human and animal subjects has been approved by the Author's relevant institutional review board.

11. **JoVE Discretion.** If the Author requests the assistance of JoVE in producing the Video in the Author's facility, the Author shall ensure that the presence of JoVE employees, agents or independent contractors is in accordance with the relevant regulations of the Author's institution. If more than one author is listed at the beginning of this Agreement, JoVE may, in its sole



## ARTICLE AND VIDEO LICENSE AGREEMENT

discretion, elect not take any action with respect to the Article until such time as it has received complete, executed Article and Video License Agreements from each such author. JoVE reserves the right, in its absolute and sole discretion and without giving any reason therefore, to accept or decline any work submitted to JoVE. JoVE and its employees, agents and independent contractors shall have full, unfettered access to the facilities of the Author or of the Author's institution as necessary to make the Video, whether actually published or not. JoVE has sole discretion as to the method of making and publishing the Materials, including, without limitation, to all decisions regarding editing, lighting, filming, timing of publication, if any, length, quality, content and the like.

12. **Indemnification.** The Author agrees to indemnify JoVE and/or its successors and assigns from and against any and all claims, costs, and expenses, including attorney's fees, arising out of any breach of any warranty or other representations contained herein. The Author further agrees to indemnify and hold harmless JoVE from and against any and all claims, costs, and expenses, including attorney's fees, resulting from the breach by the Author of any representation or warranty contained herein or from allegations or instances of violation of intellectual property rights, damage to the Author's or the Author's institution's facilities, fraud, libel, defamation, research, equipment, experiments, property damage, personal injury, violations of institutional, laboratory, hospital, ethical, human and animal treatment, privacy or other rules, regulations, laws, procedures or guidelines, liabilities and other losses or damages related in any way to the submission of work to JoVE, making of videos by JoVE, or publication in JoVE or elsewhere by JoVE. The Author shall be responsible for, and shall hold JoVE harmless from, damages caused by lack of sterilization, lack of cleanliness or by contamination due to

the making of a video by JoVE its employees, agents or independent contractors. All sterilization, cleanliness or decontamination procedures shall be solely the responsibility of the Author and shall be undertaken at the Author's expense. All indemnifications provided herein shall include JoVE's attorney's fees and costs related to said losses or damages. Such indemnification and holding harmless shall include such losses or damages incurred by, or in connection with, acts or omissions of JoVE, its employees, agents or independent contractors.

13. **Fees.** To cover the cost incurred for publication, JoVE must receive payment before production and publication of the Materials. Payment is due in 21 days of invoice. Should the Materials not be published due to an editorial or production decision, these funds will be returned to the Author. Withdrawal by the Author of any submitted Materials after final peer review approval will result in a US\$1,200 fee to cover pre-production expenses incurred by JoVE. If payment is not received by the completion of filming, production and publication of the Materials will be suspended until payment is received.

14. **Transfer, Governing Law.** This Agreement may be assigned by JoVE and shall inure to the benefits of any of JoVE's successors and assignees. This Agreement shall be governed and construed by the internal laws of the Commonwealth of Massachusetts without giving effect to any conflict of law provision thereunder. This Agreement may be executed in counterparts, each of which shall be deemed an original, but all of which together shall be deemed to be one and the same agreement. A signed copy of this Agreement delivered by facsimile, e-mail or other means of electronic transmission shall be deemed to have the same legal effect as delivery of an original signed copy of this Agreement.

A signed copy of this document must be sent with all new submissions. Only one Agreement is required per submission.

### CORRESPONDING AUTHOR

Name:

Masaya Hagiwara

Department:

NanoSquare Research Institute

Institution:

Osaka Prefecture University

Title:

Initial 3D cell cluster control in a hybrid gel cube device for repeatable pattern formation

Signature:

*Masaya Hagiwara*

Date:

October 7th, 2018.

Please submit a **signed** and **dated** copy of this license by one of the following three methods:

1. Upload an electronic version on the JoVE submission site
2. Fax the document to +1.866.381.2236
3. Mail the document to JoVE / Attn: JoVE Editorial / 1 Alewife Center #200 / Cambridge, MA 02140

## Initial 3D cell cluster control in a hybrid gel cube device for repeatable pattern formations

by Masaya Hagiwara, Rina Nobata, Tomohiro Kawahara

Manuscript: JoVE59214

On behalf of all the authors, I would like to thank the editors for taking the time to carefully review our paper and for their insightful comments. The submitted manuscript (JoVE59214) has been revised based on the editor's comments.

The point-by-point responses are as follows:

### Responses to Editorial comments

1. There are still some spelling and grammatical errors (e.g., “planer” should be “planar”); please proofread further.

**Response 1:** Professional editorial service was used to correct all grammatical errors and tracked all changes in the document.

2. Please rewrite the protocol in the imperative tense; e.g., “1.1. Prepare a polycarbonate (PC) cubic frame either by a machining process or using a 3D printer.”

**Response 2:** Corrected.

3. There are still some details that need to be further explained in the Protocol; see comments in the manuscript.

**Response 3:** The detailed were added individually to address the comments in the manuscript.

4. Formatted per JoVE guidelines (US letter-sized pages, spaces between each step, all text aligned to the left margin; see attached manuscript), the protocol is currently over 3 pages long (~3.5). There is a 2.75 page limit for filmable content; please highlight 2.75 pages or less of the Protocol (including headers and spacing) that identifies the essential steps of the protocol for the video, i.e., the steps that should be visualized to tell the most cohesive story of the Protocol. Remember that non-highlighted Protocol steps will remain in the manuscript, and therefore will still be available to the reader.

**Response 4:** The essential steps of the protocol were highlighted in the manuscript and currently the length is less than 2.5 page.

5. Figure 1: “ $\mu\text{L}$ ”, not “ $\mu\text{l}$ ”. Also, it appears that “ $^{\circ}\text{C}$ ” is of different font from the rest of the Figure; can you correct that? Lastly, what is the thickness of the cubic frame?

**Response 5:** Corrected

6. Figure 2B: Can you provide schematics for the micromolds seen here?

**Response 6:** We do not consider that schematics is required for this micromolds. Besides, those are the examples of the mold.

7. Figure 3A: There appears to already be a micromold in the frame, in contrast to the protocol and legend which indicates

it is added later.

**Response 7:** Corrected.

8. Figure 3E: “Apply medium with lower than top surface” is unclear; perhaps “Add medium to a level lower than that of the top surface” or similar.’

**Response 8:** Corrected.

9. Figure 4/5: Please explain the cameras/lenses in the Legends.

**Response 9:** The objective lens information was added in the Legends.

10. Figure 5: What is the image in the lower right-a 3d reconstruction of what is seen in the other panels, or just a schematic? Also, what are the scale bars?

**Response 10:** It is just schematic which reviewer requested previously. The explanation and scale bar was added in the legend.

11. Please ensure that all items used in the protocol are in the Table of Materials.

**Response 11:** Done.