

Journal of Visualized Experiments

Fluorescence-based measurements of phosphatidylserine/phosphatidylinositol 4-phosphate exchange between membranes

--Manuscript Draft--

Article Type:	Invited Methods Collection - JoVE Produced Video
Manuscript Number:	JoVE62177R2
Full Title:	Fluorescence-based measurements of phosphatidylserine/phosphatidylinositol 4-phosphate exchange between membranes
Corresponding Author:	Guillaume Drin Centre National de la Recherche Scientifique Sophia-Antipolis, 06560 FRANCE
Corresponding Author's Institution:	Centre National de la Recherche Scientifique
Corresponding Author E-Mail:	drin@ipmc.cnrs.fr
Order of Authors:	Guillaume Drin Souade Ikhlef Nicolas-Frédéric Lipp Maud Magdeleine
Additional Information:	
Question	Response
Please specify the section of the submitted manuscript.	Biochemistry
Please indicate whether this article will be Standard Access or Open Access.	Standard Access (US\$2,400)
Please indicate the city, state/province, and country where this article will be filmed . Please do not use abbreviations.	Sophia-Antipolis, 06560, FRANCE
Please confirm that you have read and agree to the terms and conditions of the author license agreement that applies below:	I agree to the Author License Agreement
Please provide any comments to the journal here.	We'd like to integrate if possible the section 3 (liposome preparation) in the movie

TITLE:

Fluorescence-based Measurements of Phosphatidylserine/Phosphatidylinositol 4-Phosphate Exchange Between Membranes

AUTHORS AND AFFILIATIONS:

Souade Ikhlef*, Nicolas-Frédéric Lipp*, Maud Magdeleine, Guillaume Drin

Université Côte d'Azur, Centre National de la Recherche Scientifique, Institut de Pharmacologie Moléculaire et Cellulaire, 660 route des lucioles, 06560 Valbonne, France

*Equal contribution

Email addresses of co-authors:

Souade Ikhlef ikhlef@ipmc.cnrs.fr

Nicolas-Frédéric Lipp lipp@ipmc.cnrs.fr

Maud Magdeleine magdeleine@ipmc.cnrs.fr

Corresponding author:

Guillaume Drin drin@ipmc.cnrs.fr

KEYWORDS: ([Instructions](#))

lipid transfer protein, OSBP-related protein, phosphatidylserine, phosphatidylinositol 4-phosphate, fluorescence, recombinant protein, liposome

SUMMARY:

Here, we describe protocols using fluorescent lipid sensors and liposomes to determine whether a protein extracts and transports phosphatidylserine or phosphatidylinositol 4-phosphate *in vitro*.

ABSTRACT:

Several members of the evolutionarily conserved oxysterol-binding protein (OSBP)-related proteins (ORP)/OSBP homologs (Osh) family have recently been found to represent a novel lipid transfer protein (LTP) group in yeast and human cells. They transfer phosphatidylserine (PS) from the endoplasmic reticulum (ER) to the plasma membrane (PM) *via* PS/phosphatidylinositol 4-phosphate (PI(4)P) exchange cycles. This finding allows a better understanding of how PS, which is critical for signaling processes, is distributed throughout the cell and the investigation of the link between this process and phosphoinositide (PIP) metabolism. The development of new fluorescence-based protocols has been instrumental in the discovery and characterization of this new cellular mechanism *in vitro* at the molecular level. This paper describes the production and the use of two fluorescently labelled lipid sensors, NBD-C2_{Lact} and NBD-PH_{FAPP}, to measure the ability of a protein to extract PS or PI(4)P and to transfer these lipids between artificial membranes. First, the protocol describes how to produce, label, and obtain high-purity samples of these two constructs. Secondly, this paper explains how to use these sensors with a fluorescence microplate reader to determine whether a protein can extract PS or PI(4)P from

liposomes, using Osh6p as a case study. Finally, this protocol shows how to accurately measure the kinetics of PS/PI(4)P exchange between liposomes of defined lipid composition and to determine lipid transfer rates by fluorescence resonance energy transfer (FRET) using a standard fluorometer.

INTRODUCTION:

The precise distribution of lipids between different membranes and within the membranes of eukaryotic cells^{1,2} has profound biological implications. Decrypting how LTPs function is an important issue in cell biology³⁻⁶, and *in vitro* approaches are of great value in addressing this issue⁷⁻¹¹. Here, an *in vitro*, fluorescence-based strategy is presented that has been instrumental in establishing that several ORP/Osh proteins effect PS/PI(4)P exchange between cell membranes¹² and thereby constitute a new class of LTPs. PS is an anionic glycerophospholipid that represents 2–10% of total membrane lipids in eukaryotic cells¹³⁻¹⁶. It is distributed along a gradient between the ER and the PM, where it represents 5–7% and up to 30% of glycerophospholipids, respectively¹⁷⁻¹⁹. Moreover, PS is essentially concentrated in the cytosolic leaflet of the PM. This build-up and the uneven partition of PS in the PM are critical for cellular signaling processes¹⁹. Owing to the negative charge of PS molecules, the cytosolic leaflet of the PM is much more anionic than the cytosolic leaflet of other organelles^{1,2,19,20}. This enables the recruitment, *via* electrostatic forces, of signaling proteins such as myristoylated alanine-rich C-kinase substrate (MARCKS)²¹, sarcoma (Src)²², Kirsten-rat sarcoma viral oncogene (K-Ras)²³, and Ras-related C3 botulinum toxin substrate 1 (Rac1)²⁴ that contain a stretch of positively charged amino acids and a lipidic tail.

PS is also recognized by conventional protein kinase C in a stereoselective manner *via* a C2 domain²⁵. However, PS is synthesized in the ER²⁶, indicating that it must be exported to the PM before it can play its role. It was not known how this was accomplished¹⁹ until the finding that, in yeast, Osh6p and Osh7p transfer PS from the ER to the PM²⁷. These LTPs belong to an evolutionarily conserved family in eukaryotes whose founding member is OSBP and that contains proteins (ORPs in human, Osh proteins in yeast) containing an OSBP-related domain (ORD) with a pocket to host a lipid molecule. Osh6p and Osh7p contain only an ORD whose structural features are adapted to specifically bind PS and transfer it between membranes. Nevertheless, how these proteins directionally transferred PS from the ER to the PM was unclear. Osh6p and Osh7p can trap PI(4)P as an alternative lipid ligand¹². In yeast, PI(4)P is synthesized from phosphatidylinositol (PI) in the Golgi and PMs by PI 4-kinases, Pik1p and Stt4p, respectively. In contrast, there is no PI(4)P in the ER membrane, as this lipid is hydrolyzed to PI by the Sac1p phosphatase. Hence, a PI(4)P gradient exists at both the ER/Golgi and ER/PM interfaces. Osh6p and Osh7p transfer PS from the ER to the PM *via* PS/PI(4)P exchange cycles using the PI(4)P gradient that exists between these two membranes¹².

Within one cycle, Osh6p extracts PS from the ER, exchanges PS for PI(4)P at the PM and transfers PI(4)P back to the ER to extract another PS molecule. Osh6p/Osh7p interact with Ist2p²⁸, one of the few proteins that connect and bring the ER membrane and the PM into close proximity with each other to create ER-PM contact sites in yeast²⁹⁻³¹. In addition, the association of Osh6p with negatively charged membranes becomes weak as soon as the protein extracts one of its lipid

ligands due to a conformational change that modifies its electrostatic features³². This aids Osh6p by shortening its membrane dwell time, thereby maintaining the efficiency of its lipid transfer activity. Combined with the binding to Ist2p, this mechanism could allow Osh6p/7p to both quickly and accurately execute lipid exchange at the ER/PM interface. In human cells, ORP5 and ORP8 proteins execute PS/PI(4)P exchange at ER-PM contact sites *via* distinct mechanisms³³. They have a central ORD, akin to Osh6p, but are directly anchored to the ER *via* a C-terminal transmembrane segment³³ and dock into the PM *via* an N-terminal Pleckstrin homology (PH) domain that recognizes PI(4)P and PI(4,5)P₂³³⁻³⁵. ORP5/8 use PI(4)P to transfer PS, and it has been shown that ORP5/8 additionally regulate PM PI(4,5)P₂ levels and presumably modulate signaling pathways. In turn, a decrease in PI(4)P and PI(4,5)P₂ levels suppresses ORP5/ORP8 expression as these proteins associate with the PM in a PIP-dependent fashion. Abnormally high PS synthesis, which leads to Lenz-Majewski syndrome, impacts PI(4)P levels through ORP5/8³⁶. When the activity of both proteins is blocked, PS becomes less abundant at the PM, lowering the oncogenic capability of signaling proteins³⁷.

Conversely, ORP5 overexpression seems to promote cancer cell invasion and metastatic processes³⁸. Thus, alterations to ORP5/8 activity can severely modify cellular behavior through changes in lipid homeostasis. Further, ORP5 and ORP8 occupy ER-mitochondria contact sites and preserve some mitochondrial functions, possibly by supplying PS³⁹. Additionally, ORP5 localizes to ER-lipid droplet contact sites to deliver PS to lipid droplets by PS/PI(4)P exchange⁴⁰. The strategy described herein to measure (i) PS and PI(4)P extraction from liposomes and (ii) PS and PI(4)P transport between liposomes has been devised to establish and analyze the PS/PI(4)P exchange activity of Osh6p/Osh7p^{12,32} and used by other groups to analyze the activity of ORP5/ORP8³⁵ and other LTPs^{10,41}. It is based on the use of a fluorescence plate reader, a standard L-format spectrofluorometer, and two fluorescent sensors, NBD-C2_{Lact} and NBD-PH_{FAPP}, that can detect PS and PI(4)P, respectively. NBD-C2_{Lact} corresponds to the C2 domain of the glycoprotein, lactadherin, that was reengineered to include a unique solvent-exposed cysteine near the presumed PS binding site; a polarity-sensitive NBD (7-nitrobenz-2-oxa-1,3-diazol) fluorophore is covalently linked to this residue (**Figure 1A**)¹².

The C2 domain of lactadherin (*Bos taurus*, UniProt: Q95114, residues 270–427) was cloned into a pGEX-4T3 vector to be expressed in fusion with glutathione S-transferase (GST). The C2_{Lact} sequence was then mutated to substitute two solvent-accessible cysteine residues (C270, C427) with alanine residues and to introduce a cysteine residue into a region near the putative PS-binding site (H352C mutation)¹². A cleavage site for thrombin is present between the GST protein and the N-terminus of the C2 domain. A major advantage is that this domain selectively recognizes PS in a Ca²⁺-independent manner contrary to other known C2 domains or Annexin A5⁴². NBD-PH_{FAPP} is derived from the PH domain of the human PI(4)P adaptor protein, four-phosphate-adaptor protein 1 (FAPP1), which was reengineered to include a single solvent-exposed cysteine that can be labeled with an NBD group near the PI(4)P binding site (**Figure 1A**)⁴³. The nucleotide sequence of the PH domain of the human FAPP protein (UniProt: Q9HB20, segment [1-100]) has been cloned into a pGEX-4T3 vector to be expressed in tandem with a GST tag in *Escherichia coli*. The PH_{FAPP} sequence has been modified to insert a unique cysteine residue within the membrane-binding interface of the protein that can be subsequently labeled with

N,N'-dimethyl-*N*-(iodoacetyl)-*N'*-(7-nitrobenz-2-oxa-1,3-diazol-4-yl) ethylene diamine (IANBD)⁴³. Moreover, a nine-residue linker has been introduced between the thrombin cleavage site and the N-terminus of the PH domain to ensure accessibility to the protease.

To measure PS extraction from liposomes, NBD-C2_{Lact} is mixed with liposomes made of phosphatidylcholine (PC) containing trace amounts of PS. Owing to its affinity for PS, this construct binds to the liposomes, and the NBD fluorophore experiences a change in polarity as it comes into contact with the hydrophobic environment of the membrane, which elicits a blue-shift and an increase in fluorescence. If PS is extracted almost completely by a stoichiometric amount of LTP, the probe does not associate with liposomes, and the NBD signal is lower (**Figure 1B**)³². This difference in signal is used to determine whether an LTP (*e.g.*, Osh6p) extracts PS. A similar strategy is used with NBD-PH_{FAPP} to measure PI(4)P extraction (**Figure 1B**), as described previously^{12,32}. Two FRET-based assays were designed to (i) measure PS transport from L_A to L_B liposomes, which mimic the ER membrane and the PM, respectively, and (ii) PI(4)P transport in the reverse direction. These assays are performed under the same conditions (*i.e.*, same buffer, temperature, and lipid concentration) to measure PS/PI(4)P exchange. To measure PS transport, NBD-C2_{Lact} is mixed with L_A liposomes composed of PC and doped with 5 mol% PS and 2 mol% of a fluorescent lipid—rhodamine-phycoerythrin (Rhod-PE)—and L_B liposomes incorporating 5 mol% PI(4)P.

At time zero, FRET with Rhod-PE quenches the NBD fluorescence. If PS is transported from L_A to L_B liposomes (*e.g.*, upon injecting Osh6p), a fast dequenching occurs due to the translocation of NBD-C2_{Lact} molecules from L_A to L_B liposomes (**Figure 1C**). Given the amount of accessible PS, NBD-C2_{Lact} remains essentially in a membrane-bound state over the course of the experiment¹². Thus, the intensity of the NBD signal directly correlates with the distribution of NBD-C2_{Lact} between L_A and L_B liposomes and can be easily normalized to determine how much PS is transferred. To measure the transfer of PI(4)P in the opposite direction, NBD-PH_{FAPP} is mixed with L_A and L_B liposomes; given that it only binds to L_B liposomes that contain PI(4)P, but not Rhod-PE, its fluorescence is high. If PI(4)P is transferred to L_A liposomes, it translocates to these liposomes, and the signal decreases due to FRET with Rhod-PE (**Figure 1C**). The signal is normalized to determine how much PI(4)P is transferred⁴³.

PROTOCOL:

1. Purification of NBD-C2_{Lact}

NOTE: Although this protocol details the use of a cell disruptor to break bacteria, it can be modified to use other lysis strategies (*e.g.*, a French press). At the beginning of the purification, it is mandatory to use buffer that is freshly degassed, filtered, and supplemented with 2 mM dithiothreitol (DTT) to prevent the oxidation of cysteine. However, for the protein labelling step, it is crucial to completely remove DTT. Many steps must be carried out on ice or in a cold room to avoid any protein degradation. Samples of 30 µL volume must be collected at different steps of the protocol to perform an analysis on a 15% acrylamide sodium dodecylsulfate-polyacrylamide gel electrophoresis (SDS-PAGE) gel to check the progress of the purification. Mix

177 enough denaturing Laemmli sample buffer with each aliquot, and heat the mixture at 95 °C.
178 Freeze and store the tubes at -20 °C until analysis.

180 1.1 Expression of GST-C2_{Lact} in *Escherichia coli*

182 1.1.1. Mix 20 µL of BL21 Gold competent cells with 18 µL of sterilized water. Then, mix 2 µL of
183 pGEX-C2_{Lact} plasmid (at ~65 ng/µL) with the bacteria, and transform them by electroporation.
184 Resuspend the bacteria with 150 µL of autoclaved Lennox Luria-Bertani (LB) broth medium (10
185 g/L tryptone, 5 g/L yeast extract, 5 g/L NaCl in deionized water, glucose-free). Let the bacteria
186 grow at 37 °C for 1 h in a 2 mL snap-cap microcentrifuge tube.

188 1.1.2. Inoculate 25 mL of LB medium, supplemented with 50 µg/mL ampicillin, with 150 µL of
189 bacterial suspension in a 125 mL sterile Erlenmeyer flask. Place the flask in an orbital shaker at
190 37 °C, and let the bacteria grow overnight with agitation at 185 rpm.

192 1.1.3. Fill two sterile 2 L Erlenmeyer flasks with 500 mL of LB medium supplemented with 50
193 µg/mL ampicillin, and add 5 mL of preculture suspension. Let the bacteria grow at 37 °C with
194 agitation at 220 rpm.

196 1.1.4. Periodically measure the optical density (OD) of the suspension at a wavelength (λ) of 600
197 nm. When the OD reaches a value of ~0.6–0.7, add 500 µL of a stock solution of 1 M isopropyl β-
198 D-1-thiogalactopyranoside (IPTG) to each flask to initiate the expression of GST-C2_{Lact}. Shake the
199 flasks at 185 rpm for 4 h at 37 °C.

201 1.1.5. Transfer the contents of each flask to a polypropylene centrifuge bottle. Centrifuge the
202 two bottles for 30 min at 3000 × *g* at 4 °C to pellet the bacteria. Discard the supernatant, and
203 resuspend each pellet in 50 mL of cold phosphate-buffered saline.

205 1.1.6. Transfer the bacterial suspension contained in each bottle to a 50 mL conical centrifuge
206 tube. Centrifuge the two tubes for 30 min at 3500 × *g* at 4 °C. Remove the supernatant, and store
207 the tubes, each of which contains a bacterial pellet, at -20 °C.

209 1.2. Purification of C2_{Lact}

211 1.2.1. On ice, fill two 50 mL conical centrifugal tubes with 50 mL of buffer containing 50 mM
212 Tris-HCl, pH 7.4, and 150 mM NaCl (hereafter called TN buffer), previously filtered and degassed
213 by membrane vacuum filtration.

215 1.2.2. To prepare the lysis buffer in each tube, add dissolve an ethylenediamine tetraacetic acid-
216 free protease inhibitor cocktail in the TN buffer by mild sonication or vortexing. Add other
217 antiproteases, (10 µM bestatin, 1 µg/mL pepstatin A, and 10 µM mM phosphoramidon).
218 Importantly, supplement the buffer with 2 mM DTT.

220 1.2.3. Fill the two tubes that contain the bacterial pellets prepared in step 1.1.6, with lysis buffer

to obtain a final volume of 30 mL in each tube, and slowly defrost the pellets on ice for 10 min. Crush each pellet with a stainless steel spatula, and resuspend them by vortexing the tubes and/or by pipetting the suspension back and forth with a pipette controller and a 25 mL pipette until a homogeneous suspension is obtained.

1.2.4. Perform lysis using a precooled cell disruptor (see the **Table of Materials**) by loading 30 mL of the sample inside the reservoir and running a breaking cycle in continuous mode with a pressure of 1.6 bar. Collect the lysate in the same tube, keep the tube on ice, and immediately add 250 μ L from a stock solution of 200 mM phenylmethylsulfonyl fluoride (PMSF) prepared in isopropanol.

1.2.5. Lyse the other sample following the same procedure. Use the remainder of the lysis buffer to wash the cell disruptor, and collect the wash to adjust the volume of each lysate (~30 mL) to a final volume of 50 mL.

1.2.6. Supplement each lysate with 5 mM $MgCl_2$, and add 5 μ g/mL of DNase I to fragment the DNA and thus reduce the viscosity of the sample. Incubate on ice for 30 min. Collect a sample for gel analysis.

1.2.7. Transfer each 50 mL lysate to a prechilled polycarbonate ultracentrifuge tube (two in total, see the **Table of Materials**). Centrifuge at $186,000 \times g$ at 4 °C for at least 1 h using an ultracentrifuge.

1.2.8. In parallel to the centrifugation step, dispense 1.4 mL of a slurry containing glutathione coupled to 4% agarose into two 50 mL conical centrifugal tubes, add 20 mL of TN buffer supplemented with 1 mM DTT (TND buffer) to each tube, centrifuge at $500 \times g$ for 5 min, and discard the supernatant. Repeat this washing step twice.

1.2.9. After the centrifugation of the bacterial lysate, remove a 30 μ L sample from the supernatant, and transfer the supernatant from each ultracentrifuge tube to a corresponding 50 mL conical centrifuge tube that contains clean beads. For gel analysis, resuspend the debris pellet in one of the ultracentrifuge tubes with 50 mL of TND buffer, and collect a 30 μ L sample.

1.2.10. Place the tubes on a rotator for 3–4 h at 4 °C to obtain a homogeneous bead suspension. Pool the bead suspensions in an empty 25 mL chromatography column. Let the beads decant, and remove the buffer and unbound proteins by gravity flow. Take a sample from the eluate for analysis.

1.2.11. Resuspend the beads with 20 mL of TND buffer, and collect the eluate by gravity flow. Repeat this step twice to completely wash the beads. Pool the collected eluates, and retain a 30 μ L sample for further analysis.

NOTE: After a short decantation, a volume of ~2 mL of bead suspension, to which GST-C2_{Lact} is attached, sediments at the bottom of the column.

1.2.12. Add 1 mL of bead suspension to two 2 mL snap-cap microcentrifuge tubes. Fill each tube with TND buffer to a final volume of 1.970 mL. Take a 30 μ L sample from one tube for further analysis (B1 sample). Add 10 μ L of 10 mM CaCl_2 solution and 25 μ L from a stock solution of human thrombin protease solution at 0.02 U/ μ L.

1.2.13. Place the two tubes on a rotator at 4 $^{\circ}\text{C}$ overnight to allow thrombin to cleave off the GST tag from the C2_{Lact} domain. On the next day, in each tube, mix 10 μ L of 200 mM PMSF solution with the bead suspension to inhibit the thrombin action.

1.2.14. Centrifuge the tubes at $700 \times g$ for 5 min, and collect the supernatant, which contains soluble C2_{Lact} domain, from each tube, without taking the beads. Pool the supernatants into a 2 mL snap-cap microcentrifuge tube (E1 eluate) that is kept on ice.

1.2.15. Add 1 mL of TN buffer to each tube to resuspend the beads, and wash them; repeat step 1.2.14. Perform this step thrice more to recover a maximum amount of protein. Each time, pool the collected supernatants into a new 2 mL tube (E2, E3, E4, and E5 eluates), and take an aliquot for further analysis. At the end of washing steps, take an aliquot from the bead suspension (aliquot B2).

1.2.16. Analyze the 30 μ L samples that were collected at the different steps of the purification protocol by SDS-PAGE separation on a 15% acrylamide gel.

1.2.17. Remove potential contaminating beads by pooling all the supernatants (*i.e.*, ~ 10 mL) collected during steps 1.2.14 and 1.2.15 into a 10 mL chromatography column. Collect the eluate by gravity flow, and retain the beads at the bottom of the column.

1.2.18. Concentrate the C2_{Lact} sample using a centrifugal filter unit with a molecular weight cut-off (MWCO) of 3 kDa and a centrifugation speed of $1500 \times g$. Stop the concentration procedure when the volume of the protein sample is ~ 1 mL.

1.3. Preparation and purification of NBD- C2_{Lact}

1.3.1. Equilibrate a desalting column (see the **Table of Materials**) with TN buffer. Load the column with 1 mL of concentrated C2_{Lact} sample. Allow the sample to enter the gel bed completely, add 1.5 mL of freshly degassed DTT-free TN buffer to the column, and collect the eluate by gravity flow into a 2 mL snap-cap microcentrifuge tube.

1.3.2. Dilute 50 μ L of eluate in a final volume of 300 μ L TN buffer, and record an absorbance spectrum from 230 to 450 nm using pure TN buffer as a blank. Determine the C2_{Lact} concentration based on the absorbance measured at 280 nm, considering an extinction coefficient ϵ equal to $44,920 \text{ M}^{-1}\cdot\text{cm}^{-1}$.

1.3.3. To label the C2_{Lact} construct with a NBD fluorophore, mix the protein with a ten-fold molar

excess of *N,N'*-dimethyl-*N*-(iodoacetyl)-*N'*-(7-nitrobenz-2-oxa-1,3-diazol-4-yl)ethylenediamine (IANBD amide).

1.3.3.1. Dissolve 1 mg of IANBD in anhydrous dimethylformamide (DMF), keeping in mind that the final volume of DMF used for labeling the C2_{Lact} construct must not exceed 5% (v/v) of the protein sample volume.

1.3.3.2. To determine the volume of DMF (V_{DMF}) to dissolve IANBD, first calculate the required amount of IANBD (m , expressed in mg) to label the protein by using formula 1.

$$m = 10,000 \times C \times V \times MW_{IANBD} \quad (1)$$

where C is the concentration of C2_{Lact} measured in step 1.3.2, V is the volume of the C2_{Lact} sample, and MW_{IANBD} is the molecular weight of the fluorophore (420 g/mol).

1.3.3.3. Calculate V_{DMF} by using formulae 2 (with $m_0=1$) and 3.

$$V_{DMF} = (m_0/m) \times V_{IANBD} \quad (2)$$

$$V_{IANBD} = 0.05 \times V \quad (3)$$

Where m_0 is the quantity of IANBD powder in mg, and V_{IANBD} is the volume of IANBD solution to be added to the C2_{Lact} sample.

1.3.3.4. Add volume V_{IANBD} of the freshly-prepared IANBD solution to the C2_{Lact} sample, and shake the reaction mixture at 800–900 rpm for 30 min at 25 °C using a thermomixer protected from light. Let the reaction proceed for 90 min on ice. In the meantime, clean a centrifugal filter unit (MWCO= 3 kDa) with 10 mL of TN buffer.

1.3.4. Add L-cysteine in (10-fold molar excess to IANBD) to the reaction mixture to inactivate free IANBD. Add 15 mL of TN buffer to the NBD-C2_{Lact} solution, and transfer the NBD-C2_{Lact} solution to the centrifugal filter unit.

1.3.5. Concentrate the sample to 2 mL to separate most of the free NBD from the protein by centrifugation at $1500 \times g$. Repeat this washing step two times. Centrifuge the sample in a 2 mL snap-cap centrifuge tube for 10 min at $25,000 \times g$ at 4 °C to pellet potential aggregates, and collect the supernatant.

1.3.6. Dilute 50 μ L of eluate in a final volume of 300 μ L of TN buffer. Record the absorbance spectrum from 230 to 650 nm using the eluate collected during the concentration procedure as a blank. Determine the NBD-C2_{Lact} concentration using the maximal absorbance at $\lambda=280$ and 495 nm and extinction coefficients $\epsilon = 44,920 \text{ M}^{-1}.\text{cm}^{-1}$ (protein) and $25,000 \text{ M}^{-1}.\text{cm}^{-1}$ (NBD fluorophore).

NOTE: If the two concentration values are alike, this indicates that the C2_{Lact} construct is labeled at a 1:1 ratio with the NBD group.

1.3.7. If the NBD-C2_{Lact} concentration estimated from the measurement of NBD absorbance exceeds the concentration estimated from the absorbance of tryptophan (Trp) residues, repeat step 1.3.4 to further remove free NBD.

1.3.8. Add glycerol to the sample to obtain a final concentration of 10% (v/v) to cryo-protect the NBD-C2_{Lact} construct during flash-freezing. Measure the final protein concentration.

1.3.9. Prepare 50 µL aliquots of protein in 0.5 mL snap-cap microcentrifuge tubes. Flash-freeze the tubes in liquid nitrogen, and store them at -80 °C.

2. Purification of NBD-PH_{FAPP}

NOTE: The procedure to produce and label PH_{FAPP} is identical to that of NBD-C2_{Lact} until the transfer of NBD-C2_{Lact} solution to a centrifugal filter unit in step 1.3.4. From this step onwards, follow the protocol that is described below.

2.1. After the concentration step, keep 2 mL of NBD-PH_{FAPP} at 4 °C in the dark for not more than 1 day before performing size-exclusion chromatography. Prior to size-exclusion chromatography, verify that there is no orange deposit (aggregation during concentration) at the bottom of the tube. If this is the case, centrifuge the sample at 540,000 × *g* for 10 min at 4 °C, and purify the supernatant by size-exclusion chromatography.

NOTE: Size-exclusion chromatography is performed on a column packed with crosslinked dextran-acrylamide copolymer (see the **Table of Materials**), previously equilibrated with TND buffer, using a fast protein liquid chromatography system (see the **Table of Materials**). The column must be protected from light. A flow rate of 1 mL/min was used, and the elution of the NBD-PH_{FAPP} construct was followed by recording the absorbance at λ=280 (protein) and 480 nm (NBD) at the column exit.

2.1.1. Inject the NBD-PH_{FAPP} sample loaded in a 2 mL injection loop onto the column, and immediately collect 2.5 mL fractions of eluate.

2.2. Analyze all the fractions that correspond to the main peak detected at 280 and 480 nm on a 15% SDS-PAGE gel. Mix a 25 µL sample of each fraction with 15 µL of Laemmli sample buffer prior to heating and loading onto the gel.

NOTE: A main peak, which is simultaneously detected at λ= 280 and 480 nm, appears once a volume of ~150 mL buffer is passed through the column.

2.3. Pool the fractions that exclusively contain NBD-PH_{FAPP} protein (~12.2 kDa), and add glycerol at a final concentration of 10% (v/v). Concentrate the sample using a centrifugal filter

unit with a MWCO of 3 kDa to a final volume of 1 mL using a centrifugation speed of $1500 \times g$.

2.4. Prepare aliquots, and record an absorbance spectrum as described for NBD-C2_{Lact}. Use an extinction coefficient $\epsilon = 29,450 \text{ M}^{-1}\cdot\text{cm}^{-1}$ to determine the concentration of the protein based on the absorbance measured at $\lambda = 280 \text{ nm}$.

3. Preparation of liposomes for PS and PI(4)P extraction or transfer assays

NOTE: Perform all the steps at room temperature unless otherwise specified. Handle organic solvents, rotavapor, and liquid nitrogen with caution.

3.1. Prepare fresh, filtered, and degassed 50 mM 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES)-potassium hydroxide (KOH), pH 7.4, 120 mM potassium acetate (HK) buffer.

3.2. For each type of liposome, take precise amounts of different lipids from stock solutions, and mix them in a 25 mL pear-shaped glass flask (**Table 1**). Add pure chloroform to adjust the volume of each mixture to 1 mL. Label each flask with the liposome name. Wrap the flasks containing lipid mixture doped with Rhod-PE with aluminum foil.

3.3. Place the flask on a rotary evaporator. Dry the lipids under vacuum and at 25 °C for at least 30 min at a rotation speed of 500 rpm. For lipid films that contain PI(4)P, prewarm the flask at 32–34 °C for 5 min, under gentle rotation, to properly mix PI(4)P with the other lipids before creating vacuum in the flask to remove the solvent, which will leave behind a film of dry lipids on the flask wall.

3.4. Disconnect the flask from the evaporator, and place it in a vacuum chamber for 45 min to remove any remaining traces of solvent. Fill the flask with 2 mL of HK buffer, and add a few 4 mm-diameter glass beads to the solution. Gently vortex the flask for 2 min to resuspend the lipids and prepare multilamellar lipid vesicles (MLVs) with a lipid concentration of 4 mM. Prepare 0.5 mL aliquots of MLVs in 1.5 mL screw-cap microcentrifuge tubes.

3.5. Freeze-thaw the tubes 5x (using liquid nitrogen and a water bath at 37 °C, respectively). Extrude the liposomes or store them at -20 °C.

3.5.1. Use a mini extruder to prepare the liposomes (*i.e.*, large unilamellar vesicles) from the MLVs according to the manufacturer's guidelines. Use a polycarbonate filter with uniform cylindrical pores of 200 nm diameter.

3.5.2. To prepare each type of liposome, extrude at least 250 μL of the corresponding suspension of MLVs. Store the extruded liposomes at 4 °C and in the dark if they contain Rhod-PE. Use the liposomes within 2 days.

[Insert **Table 1** here]

4. Measurement of PS or PI(4)P extraction

NOTE: Measurements must be conducted using a black 96-well plate and a fluorescence plate reader equipped with monochromators: one for fluorescence excitation and one for emission, with a variable bandwidth.

4.1. Prepare fresh, filtered, and degassed HK buffer supplemented with 1 mM MgCl₂ (HKM buffer). Prepare pure PC liposomes and PC liposomes doped with 2 mol% PS or 2 mol% PI(4)P (4 mM final lipid concentration, see Table 1).

NOTE: Keep the tubes filled with the suspension of extruded liposomes at room temperature throughout the experiment, and keep the proteins on ice. Additionally, protect the lipid sensors from light.

4.2. For the PS extraction assay, in one well, mix liposomes containing 2 mol% PS (80 μM final lipid concentration, 0.8 μM accessible PS concentration) with NBD-C2_{Lact} (250 nM final concentration) in a final volume of 100 μL. Fill a second well with the same amount of liposome (80 μM, 2 mol% PS) and NBD-C2_{Lact} (250 nM) mixed with 3 μM LTP (Osh6p as a positive control or a protein of interest).

NOTE: An incubation time of 5 min is sufficient for Osh6p to achieve lipid extraction.

4.3. Fill a third well with NBD-C2_{Lact} (250 nM) mixed with pure PC liposomes (80 μM). Fill a fourth well only with pure PC liposomes (80 μM). Repeat steps 4.1–4.3 to prepare three additional series of four wells.

4.4. For each well, record an NBD spectrum from 505 to 650 nm (bandwidth 5 nm) upon excitation at 490 nm (bandwidth 5 nm) at 25 °C. For each series, subtract the spectrum recorded with only liposomes from the other spectra.

NOTE: F and F_{max} correspond to the intensity at 536 nm measured with PS-containing liposomes in the presence or absence of LTP, respectively, whereas F₀ is the intensity at the same wavelength with pure PC liposomes. For each series, the percentage of accessible PS that is extracted by the protein is given by using the following formula.

$$100 \times (1 - ((F - F_0) / (F_{\max} - F_0))) \quad (4)$$

4.5. For the PI(4)P extraction assay, prepare liposomes doped with 2 mol% PI(4)P, and carry out measurements with the NBD-PH_{FAPP} probe. Perform control experiments and determine the extraction percentage in the same way as described above.

NOTE: Liposome and protein concentration are identical to those used in the PS extraction assay.

5. Real-time measurement of PS transport

NOTE: A standard fluorimeter (90° format) equipped with a temperature-controlled cell holder and a magnetic stirrer is used to record lipid transfer kinetics. To accurately acquire data, it is key to permanently maintain the sample at the same temperature (set between 25 and 37 °C depending on the origin of the protein (*e.g.*, yeast or human)) and to constantly stir it. The protocol described below is for the measurement of lipid transport in a 600 µL sample contained in a cylindrical quartz cell.

5.1. Prepare freshly degassed and filtered HKM buffer. Keep the tubes containing extruded liposomes at room temperature. Wrap tubes containing liposomes with Rhod-PE in aluminum foil, and/or store them in an opaque box to prevent any photobleaching.

5.2. Adjust the excitation and emission monochromators at $\lambda = 460$ nm (with a short bandwidth (1-3 nm)) and at $\lambda = 530$ nm (with a large bandwidth (≥ 10 nm)), respectively. Set the acquisition time at 25 min with a time resolution ≤ 1 s.

5.3. In the quartz cuvette, dilute 30 µL of the L_A liposome suspension and a volume of NBD-C2_{Lact} stock solution in prewarmed HKM buffer to prepare a 570 µL sample that contains 200 µM total lipids and 250 nM NBD-C2_{Lact}. Add a small magnetic stirring bar, and position the cuvette in the fluorometer holder.

5.4. Once the sample is thermally equilibrated (after 3–5 min), trigger the measurement. After 1 min, add 30 µL of L_B liposome suspension (final concentration of 200 µM total lipids) to the sample. After 3 min, inject LTP into the sample so that the final concentration of the LTP is 200 nM, and acquire the signal for the remaining 21 min.

5.5. Carry out a parallel experiment to normalize the NBD signal. Mix 30 µL of L_{A-Eq} liposome suspension with 250 nM NBD-C2_{Lact} in HKM buffer (final volume of 570 µL). After 1 min, inject 30 µL of L_{B-Eq} liposome suspension.

NOTE: The lipid composition of L_{A-Eq} and L_{B-Eq} liposomes are similar to that of L_A and L_B liposomes used in the transfer assay, except that each of them contains 2.5 mol% PS and 2.5 mol% PI(4)P. As a result, the NBD signal that is measured, referred to as F_{Eq} , corresponds to the signal that should be measured if PS was fully equilibrated between L_A and L_B liposomes by a transfer process.

5.6. Convert the kinetic curves measured with an LTP of interest to determine the amount of PS (in µM) transferred from L_A to L_B liposomes over time. Normalize each data point (F) of the curve by using the following formula.

$$F_{Norm} = (F - F_0) / (F_{Eq} - F_0) \quad (5)$$

in which F_0 corresponds to the NBD signal just before the addition of an LTP, and F_{Eq} is the signal

measured in step 5.5.

NOTE: The amount of PS (in μM) transferred from L_A to L_B liposomes corresponds to $2.5 \times F_{\text{Norm}}$, considering that the equilibrium corresponds to a situation where one half of accessible PS molecules, contained in the outer leaflet of the L_A liposomes, (*i.e.*, corresponding to 5 mol% of $0.5 \times 200 \mu\text{M}$ total lipids) has been transferred into L_B liposomes.

6. Real-time measurement of PI(4)P transport

6.1. Set the fluorimeter (excitation and emission wavelength, bandwidth, acquisition time, time resolution) as done for the PS transfer assay. Likewise, use the same buffer, cuvette, and liposomes to perform the experiments under constant stirring at the same temperature.

6.2. In the cuvette, mix $30 \mu\text{L}$ of L_B liposome suspension and NBD- PH_{FAPP} with prewarmed HKM buffer to obtain a final volume of $570 \mu\text{L}$ ($200 \mu\text{M}$ total lipids, 250 nM NBD- PH_{FAPP}). Once the thermal equilibration of the sample is reached, start the measurement, and after 1 min, inject $30 \mu\text{L}$ of L_A liposome suspension. After 3 min, inject the LTP of interest (final concentration of 200 nM), and record the signal.

6.3. Perform a second experiment to normalize the NBD signal. Mix $30 \mu\text{L}$ of $L_{B\text{-Eq}}$ liposome suspension with 250 nM NBD- PH_{FAPP} in $570 \mu\text{L}$ of HKM buffer. After 1 min, inject $30 \mu\text{L}$ of $L_{A\text{-Eq}}$ liposome suspension.

NOTE: Here, the NBD signal, referred to as F_{Eq} , corresponds to the one which should be measured if PI(4)P was fully equilibrated between L_A and L_B liposomes.

6.4. Convert the kinetic curves to determine the amount of PI(4)P (in μM) transferred from L_B to L_A liposomes over time. Each data point (F) is normalized by using formula 5 in which F_0 corresponds to the NBD signal prior to the addition of an LTP, and F_{Eq} is the signal measured in step 6.3.

NOTE: The amount of PI(4)P (in μM) transferred from L_A to L_B liposomes corresponds to $2.5 \times F_{\text{Norm}}$, considering that the equilibrium corresponds to a situation where one half of the PI(4)P contained in the outer leaflet of the L_B liposomes (*i.e.*, $0.5 \times 5 \mu\text{M}$) has been transferred in L_A liposomes.

7. Analysis of kinetics curves

7.1. Quantify the extent to which an LTP is efficient by determining the speed at which this LTP transfers lipids from one liposome population to the other one in the first few seconds following its injection to the cuvette.

7.1.1. Perform a linear regression of the first data points of the transfer kinetics to obtain a slope. Divide the slope value by the LTP concentration in the reaction mixture to determine the

number of lipid molecules transferred per protein per time unit (min or s).

REPRESENTATIVE RESULTS

Figure 2A shows an SDS-PAGE analysis of the products of different steps leading to the purification of C2_{Lact}. Lane 1 shows the protein profile of the lysed bacteria expressing GST-C2_{Lact} (~44.8 kDa), whereas lanes 2 and 3 respectively show the protein profiles of the supernatant and bacterial debris after ultracentrifugation. The comparison of these lanes shows that GST-C2_{Lact} has been recovered in the supernatant and thus can be isolated using glutathione-linked agarose beads. Lanes 4 and 5 show the protein profiles of the supernatant after incubation with beads and washes recovered by gravity flow, whereas lane 6 shows the profile of proteins that have been retained on the beads. An analysis of these lanes indicates that almost all GST-C2_{Lact} has been recovered from the beads.

Lanes 8–12 show the presence of a major band corresponding to C2_{Lact} (~17.9 kDa) in the supernatants recovered through successive washes of the beads after thrombin treatment. Lane 13 indicates that non-cleaved GST-C2_{Lact}, along with GST (~26.9 kDa), remained bound to beads after this treatment. The comparison of these lanes indicates that the cleavage procedure, albeit not 100% efficient, did yield C2_{Lact} that was then fluorescently labelled. **Figure 2B** shows an ultraviolet (UV)-visible absorbance spectrum of C2_{Lact} labelled with NBD. As the construct is 100% pure, these results confirm that all C2_{Lact} molecules were labelled with an NBD group based on the optical density measured at 280 nm (Trp) and 495 nm (NBD). The purity of NBD-C2_{Lact} and its fluorescence were determined by SDS-PAGE analysis (**Figure 2C**).

[Insert Figure 2]

Figure 3 shows the results from PS and PI(4)P extraction assays using Osh6p. When only incubated with liposomes containing 2 mol% PS, the fluorescence of NBD-C2_{Lact} was maximal as the NBD fluorophore was inserted in the membrane (*i.e.*, a hydrophobic context), indicating that the sensor was membrane-bound. In the presence of Osh6p, the fluorescence was lower and comparable to that measured with pure PC liposomes (**Figure 3A**). The normalization of intensity values at 536 nm indicated that ~75% of accessible PS was extracted. In the second assay, NBD-PH_{FAPP} was mixed with liposomes containing 2 mol% PI(4)P. The NBD signal was high without Osh6p, but low when Osh6p was present and was similar to that measured with PI(4)P-free liposomes (**Figure 3B**). An analysis of the intensity revealed that ~100 % of accessible PI(4)P was extracted by the LTP.

[Insert Figure 3]

Figure 4A shows typical results from a PS transfer assay using Osh6p as an LTP. At time zero, NBD-C2_{Lact} was mixed with L_A liposomes containing 5 mol% 16:0/18:1-PS (POPS) and 2 mol% Rhod-PE in a volume of 570 µL of HKM buffer at 30 °C. As the probe bound to L_A liposomes, its signal was quenched due to FRET with Rhod-PE present in these liposomes. After one minute, Rhod-PE-free L_B liposomes (30 µL) were added; this was expected to only elicit a slight change in the signal due to light diffusion by this second liposome population and/or a dilution effect. The signal intensity after the addition of the L_B liposomes corresponds to F₀. The injection of a few µL of a stock solution

of Osh6p (typically 40 μ M) to dilute 200 nM of the protein in the reaction mix elicited a slow increase in the NBD signal due to the dequenching of the fluorophore as PS was transported from L_A to L_B liposomes, thereby promoting the translocation of NBD-C2_{Lact}.

When L_B liposomes contained 5 mol% PI(4)P, the dequenching was much faster, as PS was transferred more rapidly by Osh6p to L_B liposomes due to the counterexchange of PS with PI(4)P by the LTP (second curve). The third curve corresponds to an experiment in which NBD-C2_{Lact} was mixed with equal amounts of L_{A-Eq} and L_{B-Eq} liposomes. The signal was higher than F_0 and corresponded to a situation where the probe was evenly bound to L_A and L_B liposomes, thus reflecting a situation where PS was fully equilibrated between the two populations of liposomes. F_{Eq} was calculated by averaging the value of the signal measured in the last 5 min of the experiment.

[Insert Figure 4]

Figure 4B shows average kinetic curves of PS transfer from L_A liposomes to L_B liposomes, doped or not doped with PI(4)P after the normalization of F data using F_0 and F_{Eq} as reference values. The initial transport rate for each experiment was calculated by fitting the initial data points measured after the injection of the protein with a linear function. **Figure 4C** shows the mean initial PS transport rate determined from three distinct experiments using L_B liposomes with or without PI(4)P. When L_B liposomes contained 0 and 5 mol% PI(4)P, the rates were respectively equal to 1.4 and 15.8 PS.min⁻¹ per Osh6p molecule. **Figure 5A** shows typical results from a PI(4)P transfer assay using Osh6p as an LTP, which was carried out with the same materials and conditions as those used for the PS transport assay.

[Insert Figure 5]

At time zero, NBD-PH_{FAPP} was mixed with L_B liposomes containing 5 mol% PI(4)P in a volume of 570 μ L of HKM buffer. Because NBD-PH_{FAPP} was bound to L_B liposomes, its signal was high. After one minute, L_A liposomes (30 μ L) were added, which was expected to only elicit a slight change in the signal. The intensity then corresponded to F_0 . Injecting Osh6p (200 nM final concentration) into the reaction mix triggered a quenching of the NBD signal due to the translocation of NBD-PH_{FAPP} molecules to L_A liposomes as PI(4)P was transferred from L_B liposomes to L_A liposomes. When L_A liposomes contained 5 mol% POPS, the dequenching was much faster owing to a faster PI(4)P transfer resulting from PS/PI(4)P exchange. The third curve corresponds to an experiment in which NBD-PH_{FAPP} was mixed with equal amounts of L_{A-Eq} and L_{B-Eq} liposomes.

The signal was lower than F_0 as it corresponded to a situation where the probe was uniformly bound to L_A and L_B liposomes, thus indicative of a full equilibration of PI(4)P between the liposomes. F_{Eq} was calculated by averaging the value of the signal measured in the last 5 min of the experiment. **Figure 5B,C** show averaged kinetics curves obtained after signal normalization and mean PI(4)P initial transfer rates measured with L_A liposomes that were doped or not doped with 5 mol% PS. Here, it is worth noting that both lipid ligands were transported faster and with similar velocities when each lipid was initially present in L_A and L_B membranes, respectively, which indicates that Osh6p is a PS/PI(4)P exchanger.

FIGURE AND TABLE LEGENDS:

Figure 1: Description of the fluorescent lipid sensors and *in vitro* assays. (A) Three-dimensional models of NBD-C2_{Lact} and NBD-PH_{FAPP} based on the crystal structure of the C2 domain of bovine lactadherin (PDB ID: 3BN6⁴⁸) and the NMR structure of the PH domain of the human FAPP1 protein (PDB ID: 2KCJ⁴⁶). An *N,N'*-dimethyl-*N*-(thioacetyl)-*N'*-(7-nitrobenz-2-oxa-1,3-diazol-4-yl)ethylenediamine moiety, built manually and energetically minimized, was grafted onto the thiol function of C352 (NBD-C2_{Lact}) and C13 (NBD-PH_{FAPP}) residues (represented as spheres, with carbon in green, nitrogen in blue, oxygen in red, sulfur in yellow, and hydrogen in white). The surface of the lipid-binding site of each probe was colored in blue. (B) Extraction assays. In the PS extraction assay, PC/PS liposomes (98/2 mol/mol) were incubated with 250 nM NBD-C2_{Lact}. In the absence of extraction, the probe strongly bound to the liposomes, resulting in a blue shift of NBD fluorescence and an increase of its emission intensity. If PS extraction occurred in the presence of an LTP (*e.g.*, Osh6p), the probe dissociated from the liposomes, and its fluorescence was lower. In the PI(4)P extraction assay, liposomes were doped with 2 mol% PI(4)P, and 250 nM NBD-PH_{FAPP} was used. (C) FRET-based lipid transport assays. In the PS transport assay, PC/PS/Rhod-PE liposomes (93/5/2 mol/mol, L_A) were incubated with 250 nM NBD-C2_{Lact}. PC liposomes, doped or not with 5 mol% PI(4)P, and Osh6p were added at *t* = 1 min and *t* = 4 min, respectively. If PS transport occurs, this elicits a dequenching of the NBD signal corresponding to the translocation of NBD-C2_{Lact} from L_A to L_B liposomes. In the PI(4)P transport assay, L_B liposomes doped with 5 mol% PI(4)P were incubated with 250 nM NBD-PH_{FAPP}. PC liposomes (L_A) doped or not with 5 mol% PS were added. If PI(4)P transport occurs, this causes a quenching of the NBD signal due to the translocation of NBD-PH_{FAPP} from L_B to L_A liposomes. Abbreviations: NBD = 7-nitrobenz-2-oxa-1,3-diazol) fluorophore; NMR = nuclear magnetic resonance; FAPP1 = four-phosphate-adaptor protein 1; PDB = Protein Data Bank; PS= phosphatidylserine; PC = phosphatidylcholine; LTP = lipid transfer protein; Osh6p = oxysterol-binding protein (OSBP) homolog 6 protein; PI(4)P = phosphatidylinositol 4-phosphate; FRET = fluorescence resonance energy transfer; Rhod-PE = rhodamine-phycoerythrin; L_A liposomes = liposomes composed of PC and doped with 5 mol% PS and 2 mol% of Rhod-PE; L_B liposomes = liposomes incorporating 5 mol% PI(4)P.

Figure 2: NBD-C2_{Lact} purification. (A) SDS-PAGE analysis was used to check the presence of the protein at different steps of the purification procedure before labeling. The arrowheads indicate the position of the C2_{Lact} domain (red arrowhead), of the GST alone (grey arrowhead), and of the GST-C2_{Lact} construct (black arrowhead). (B) UV-visible absorbance spectrum of NBD-C2_{Lact}. (C) SDS-PAGE analysis of the purified NBD-C2_{Lact}. The first image was acquired under UV illumination without staining and reveals the presence of the NBD-C2_{Lact} construct as it emits fluorescence. The second image shows the same gel after a protein staining procedure (see the **Table of Materials**). Abbreviations: NBD = 7-nitrobenz-2-oxa-1,3-diazol) fluorophore; NBD-C2_{Lact} = *N,N'*-dimethyl-*N*-(thioacetyl)-*N'*-(7-nitrobenz-2-oxa-1,3-diazol-4-yl)ethylenediamine moiety linked to the thiol function of C352 (NBD-C2_{Lact}) C2 domain of bovine lactadherin (PDB ID: 3BN6⁴⁸); PDB = Protein Data Bank; SDS-PAGE = sodium dodecylsulfate polyacrylamide gel electrophoresis; GST = glutathione S-transferase; MW = molecular-weight size marker; UV = ultraviolet.

Figure 3: Extraction assays. (A) Fluorescence spectra of NBD-C2_{Lact} (250 nM) measured upon

excitation at 460 nm in the presence of liposomes (80 μ M, 2 mol% PS) in the absence or presence of 3 μ M Osh6p. Reference spectra were recorded with pure PC liposomes incubated or not with NBD-C2_{Lact} (left panel). Several spectra were recorded from different series of wells, corrected by subtracting the background scattering signal from DOPC liposomes alone and averaged ($n=4$, \pm SEM). The percentage of accessible PS that was extracted is indicated (right panel). (B) Fluorescence spectra of NBD-PH_{FAPP} (250 nM) mixed with liposomes (80 μ M, 2 mol% PI(4)P) in the absence or presence of 3 μ M Osh6p. Reference spectra recorded with PC liposomes in the presence and absence of the sensor are shown (left panel). Several spectra were recorded from different series of wells, corrected by subtracting the background scattering signal from DOPC liposomes alone and averaged ($n=4$, \pm SEM). The percentage of accessible PI(4)P that was extracted is indicated (right panel). Abbreviations: NBD = 7-nitrobenz-2-oxa-1,3-diazol fluorophore; NBD-C2_{Lact} = *N,N'*-dimethyl-*N*-(thioacetyl)-*N'*-(7-nitrobenz-2-oxa-1,3-diazol-4-yl)ethylenediamine moiety linked to the thiol function of C352 (NBD-C2_{Lact}) C2 domain of bovine lactadherin (PDB ID: 3BN6⁴⁸); PDB = Protein Data Bank; PS= phosphatidylserine; PC = phosphatidylcholine; DOPC = dioleoylphosphatidylcholine; Osh6p = oxysterol-binding protein (OSBP) homolog 6 protein; PI(4)P = phosphatidylinositol 4-phosphate; SEM = standard error of the mean; a.u. = arbitrary units.

Figure 4: Typical PS transport kinetics measured with Osh6p and reference curve. (A) L_B liposomes devoid of PI(4)P (200 μ M total lipids) and Osh6p (200 nM) were sequentially added to a cuvette containing NBD-C2_{Lact} (250 nM) and L_A liposomes (200 μ M) doped with 5 mol% PS and 2 mol% Rhod-PE (left curve). The same experiment was performed with L_B liposomes doped with 5 mol% PI(4)P (middle curve). To determine F_{Eq} , NBD-C2_{Lact} (250 nM) was premixed with L_{A-Eq} liposomes; then, L_{B-Eq} liposomes were added (right curve). (B) Averaged PS transport curves determined after the normalization of several measurements done using L_B liposomes with or without PI(4)P (mean \pm SEM, $n=3$). (E) Initial PS transfer rates (mean \pm SEM, $n=3$). Abbreviations: NBD = 7-nitrobenz-2-oxa-1,3-diazol fluorophore; NBD-C2_{Lact} = *N,N'*-dimethyl-*N*-(thioacetyl)-*N'*-(7-nitrobenz-2-oxa-1,3-diazol-4-yl)ethylenediamine moiety linked to the thiol function of C352 (NBD-C2_{Lact}) C2 domain of bovine lactadherin (PDB ID: 3BN6⁴⁸); PDB = Protein Data Bank; PS= phosphatidylserine; Osh6p = oxysterol-binding protein (OSBP) homolog 6 protein; PI(4)P = phosphatidylinositol 4-phosphate; Rhod-PE = rhodamine-phycoerythrin; L_A liposomes = liposomes composed of phosphatidylcholine and doped with 5 mol% PS and 2 mol% of Rhod-PE; L_B liposomes = liposomes incorporating 5 mol% PI(4)P; F = fluorescence; F_0 = fluorescence corresponding to NBD before addition of Osh6p; F_{Eq} = fluorescence signal if PS fully equilibrated between L_A and L_B liposomes by a transfer process; SEM = standard error of the mean.

Figure 5: Typical PI(4)P transport kinetics measured with Osh6p and reference curve. (A) PS-free liposomes containing 2 mol% Rhod-PE (L_A , 200 μ M total lipids) and Osh6p (200 nM) were sequentially added to a cuvette containing NBD-PH_{FAPP} (250 nM) and L_B liposomes (200 μ M) doped with 5 mol% PI(4)P (left curve). The same experiment was performed with L_A liposomes doped with 5 mol% PS (middle curve). To determine F_{Eq} , NBD-PH_{FAPP} (250 nM) was premixed with L_{B-Eq} liposomes; then, L_{A-Eq} liposomes were added (right curve). (B) PI(4)P transfer kinetics determined after normalization of several measurements of L_A liposomes with or without PS (mean \pm SEM, $n=3$). (C) Initial PI(4)P transfer rates (mean \pm SEM, $n=3$). Abbreviations: NBD = 7-

nitrobenz-2-oxa-1,3-diazol) fluorophore; NBD-PH_{FAPP} = NBD-labeled Pleckstrin homology domain of the human PI(4)P adapter protein, four-phosphate-adaptor protein 1 (FAPP1, UniProt: Q9HB20, segment [1-100]); PS= phosphatidylserine; Osh6p = oxysterol-binding protein (OSBP) homolog 6 protein; PI(4)P = phosphatidylinositol 4-phosphate; Rhod-PE = rhodamine-phycoerythrin; L_A liposomes = liposomes composed of phosphatidylcholine and doped with 5 mol% PS and 2 mol% of Rhod-PE; L_B liposomes = liposomes incorporating 5 mol% PI(4)P; F = fluorescence; F₀ = fluorescence corresponding to NBD before addition of Osh6p; F_{Eq} = fluorescence signal if PS fully equilibrated between L_A and L_B liposomes by a transfer process; SEM = standard error of the mean.

Table 1: Volumes of lipid stock solutions to be mixed for liposome preparation. Abbreviations: PS= phosphatidylserine; PC = phosphatidylcholine; PI(4)P = phosphatidylinositol 4-phosphate; Rhod-PE = rhodamine-phycoerythrin; DOPC = dioleoylphosphatidylcholine; 1-palmitoyl-2-oleoyl-sn-glycero-3-phospho-L-serine; 16:0 Liss Rhod-PE = 1,2-dipalmitoyl-sn-glycero-3-phosphoethanolamine-N-(lissamine rhodamine B sulfonyl) (ammonium salt).

DISCUSSION:

The outcomes of these assays directly rely on the signals of the fluorescent lipid sensors. Thus, the purification of these probes labelled at a 1:1 ratio with NBD and without free NBD fluorophore contamination is a critical step in this protocol. It is also mandatory to check whether the LTP under examination is properly folded and not aggregated. The amount of LTP tested in the extraction assays must be equal to or higher than that of accessible PS or PI(4)P molecules to properly measure whether this LTP efficiently extracts these lipids. Indeed, NBD-C2_{Lact} and NBD-PH_{FAPP} bind to PS and PI(4)P, respectively, following a classical saturation binding curve. Given their respective affinities for PS and PI(4)P, these probes remain largely bound to the liposomes even if the liposomes contain residual traces of these ligands. This can lead to the erroneous conclusion that an LTP does not efficiently extract these lipids. The protocol relies on standard and commercially available PC, PS, and PI(4)P subspecies that are 18:1/18:1-PC, 16:0/18:1-PS, and 16:0/16:0-PI(4)P, respectively. Performing experiments using lipid species with other acyl chains can give different results, as previously reported^{12,35}.

In the transfer assays, if the bulk lipid composition (not considering PS or PI(4)P) of L_A and L_B liposomes has to be modified, it is key to also perform the control experiments using L_{A-Eq} and L_{B-Eq} liposomes with a bulk composition similar to that of L_A and L_B liposomes, respectively. The same principle applies for the extraction assays. Genetically encoded fluorescent lipid-binding domains are broadly used to analyze lipid distribution inside cells⁴⁴. Caution is recommended when analyzing such experiments as the association of these lipid probes with the membrane, although primarily driven by the presence of the targeted lipid, can be influenced by other parameters (density of anionic lipids, lipid-packing) that differ between cell compartments. In these assays, the composition of liposomes is much simpler than that of cellular membranes: they are mostly made of PC, a zwitterionic lipid, and thereby expose a rather inert surface that does not impact how PS and PI(4)P are recognized by their corresponding sensor. PH_{FAPP} can be used in the presence of anionic lipids, such as PS, phosphatidic acid (PA), and PI³², and C2_{Lact} does not recognize PI(4)P¹²; both these observations allow unbiased measurements of PS/PI(4)P

exchange. NBD-PH_{FAPP} is not influenced by sterol⁴³.

However, it is not known whether these probes, in particular, NBD-C2_{Lact}, can be used with liposomes with extreme features (*e.g.*, very low or high lipid-packing, highly negatively charged surface, presence of lipid domains). Despite this potential limitation with respect to liposome composition, this method based on fluorescent sensors has tremendous advantages compared to other methods. First, it does not rely on fluorescently labeled PS and PI(4)P that bear an extra bulky group and are likely not properly accommodated by the binding pocket of LTPs. Second, these transport assays offer far better time resolution than methods based on liposome separation (radioactivity-based assays⁴⁵ or mass spectrometry³³), and it should be noted that radiolabeled PI(4)P and PS are not commercially available. The capacity of a protein to transfer PS or PI(4)P is not impacted by the association of the fluorescent sensors with liposomes. Indeed, if the amount of NBD-C2_{Lact} or NBD-PH_{FAPP} and that of PS and PI(4)P in the outer leaflet of liposomes that is accessible to the LTP is considered, only 5% of PS or PI(4)P are associated with probe during a kinetics measurement.

Moreover, only 0.55–0.86% of the membrane surface is covered by the probe, considering the total surface of liposomes (L_A+L_B liposomes; 5.1×10^{16} nm² with an area of 0.7 nm² per lipid)—the membrane surface that one individual C2 or PH molecule can occupy (≈ 3.1 and 4.8 nm², respectively, estimated from references⁴⁶⁻⁴⁸) and their concentration (250 nM). Thus, the principles underlying these assays are adapted to properly analyze the kinetic and mechanistic aspects of LTPs. As mentioned in the introduction, alterations in the activity of ORP5/8 can lead to cellular dysfunctions⁴⁹. For example, the invasiveness of pancreatic cancer cells seems to rely on the level of ORP5 expression. Moreover, a causality exists between high levels of ORP5 expression and the poor prognosis of human pancreatic cancer. A high level of ORP5 expression is also detected in lung tumor tissues, and more particularly, in metastatic cases. Interestingly, the ability of ORP5 to transfer lipids might explain why it promotes cell proliferation and migration.

ORP5 upregulates the mammalian target of rapamycin complex 1 (mTORC1) complex, which plays a central role in cell survival and proliferation, probably because it enhances the activity of Akt, an upstream activator of mTORC1, by supplying the PM with PS. Overall, these observations suggest that ORP5 might be an interesting pharmacological target, and that these *in vitro* assays might serve to screen molecules that are able to inhibit its activity. This assay can also help to better define whether other members of the ORP/Osh family are PS/PI(4)P exchangers. Notably, ORP10 was found to encapsulate PS *in vitro* and transfer PS in ER-Golgi contact sites^{27,50}, but it is still unknown whether it acts as a PS/PI(4)P exchanger. Furthermore, these protocols can serve to explore the ability of LTPs belonging to other families to transport PI(4)P, as was recently shown with steroidogenic acute regulatory transfer-like proteins¹⁰, or PS. Additionally, NBD-PH_{FAPP} has been used to follow the ability of LTP to transport PI(4,5)P₂ between liposomes, as it recognizes this PIP^{10,35}. Finally, this strategy could be adapted to measure other extraction or transport processes *in vitro* by using other lipid-binding domains (*e.g.*, a non-catalytic PI-PLC to detect PI⁵¹, sporulation-specific protein 20 fragment to detect PA⁵², domain 4 (D4) of perfringolysin O to detect sterol⁵³).

ACKNOWLEDGMENTS:

We are grateful to Dr. A. Cuttriss for her careful proofreading of the manuscript. This work is funded by the French National Research Agency grant ExCHANGE (ANR-16-CE13-0006) and by the CNRS.

DISCLOSURES:

The authors declare that there are no conflicts of interest.

REFERENCES

- 1 Drin, G. Topological regulation of lipid balance in cells. *Annual Review of Biochemistry*. **83**, 51–77 (2014).
- 2 Bigay, J., Antonny, B. Curvature, lipid packing, and electrostatics of membrane organelles: defining cellular territories in determining specificity. *Developmental Cell*. **23** (5), 886–895 (2012).
- 3 Prinz, W. A. Lipid trafficking sans vesicles: where, why, how? *Cell*. **143** (6), 870–874 (2010).
- 4 Holthuis, J. C., Menon, A. K. Lipid landscapes and pipelines in membrane homeostasis. *Nature*. **510** (7503), 48–57 (2014).
- 5 Wong, L. H., Copic, A., Levine, T. P. Advances on the transfer of lipids by lipid transfer proteins. *Trends in Biochemical Sciences*. **42** (7), 516–530 (2017).
- 6 Wong, L. H., Gatta, A. T., Levine, T. P. Lipid transfer proteins: the lipid commute via shuttles, bridges and tubes. *Nature Reviews Molecular Cell Biology*. **20** (2), 85–101 (2019).
- 7 Iaea, D. B., Dikiy, I., Kiburu, I., Eliezer, D., Maxfield, F. R. STARD4 membrane interactions and sterol binding. *Biochemistry*. **54** (30), 4623–4636 (2015).
- 8 Wilhelm, L. P. et al. STARD3 mediates endoplasmic reticulum-to-endosome cholesterol transport at membrane contact sites. *The EMBO Journal*. **36** (10), 1412–1433 (2017).
- 9 Bian, X., Saheki, Y., De Camilli, P. Ca(2+) releases E-Syt1 autoinhibition to couple ER-plasma membrane tethering with lipid transport. *The EMBO Journal*. **37** (2), 219–234 (2018).
- 10 Horenkamp, F. A., Valverde, D. P., Nunnari, J., Reinisch, K. M. Molecular basis for sterol transport by StART-like lipid transfer domains. *The EMBO Journal*. **37** (6), e98002 (2018).
- 11 Jentsch, J. A. et al. Structural basis of sterol binding and transport by a yeast StArkin domain. *The Journal of Biological Chemistry*. **293** (15), 5522–5531 (2018).
- 12 Moser von Filseck, J. et al. INTRACELLULAR TRANSPORT. Phosphatidylserine transport by ORP/Osh proteins is driven by phosphatidylinositol 4-phosphate. *Science*. **349** (6246), 432–436 (2015).
- 13 Daum, G. et al. Systematic analysis of yeast strains with possible defects in lipid metabolism. *Yeast*. **15** (7), 601–614 (1999).
- 14 Ejsing, C. S. et al. Global analysis of the yeast lipidome by quantitative shotgun mass spectrometry. *Proceedings of the National Academy of Sciences of the United States of America*. **106** (7), 2136–2141 (2009).
- 15 Leidl, K., Liebisch, G., Richter, D., Schmitz, G. Mass spectrometric analysis of lipid species of human circulating blood cells. *Biochimica et Biophysica Acta*. **1781** (10), 655–664 (2008).
- 16 Sampaio, J. L. et al. Membrane lipidome of an epithelial cell line. *Proceedings of the National Academy of Sciences of the United States of America*. **108** (5), 1903–1907 (2011).
- 17 Vance, J. E., Steenbergen, R. Metabolism and functions of phosphatidylserine. *Progress in*

845 *Lipid Research*. **44** (4), 207–234 (2005).

846 18 Zinser, E. et al. Phospholipid synthesis and lipid composition of subcellular membranes in
847 the unicellular eukaryote *Saccharomyces cerevisiae*. *Journal of Bacteriology*. **173** (6), 2026–2034
848 (1991).

849 19 Leventis, P. A., Grinstein, S. The distribution and function of phosphatidylserine in cellular
850 membranes. *Annual Review of Biophysics*. **39**, 407–427 (2010).

851 20 Yeung, T. et al. Membrane phosphatidylserine regulates surface charge and protein
852 localization. *Science*. **319** (5860), 210–213 (2008).

853 21 Kim, J., Shishido, T., Jiang, X., Aderem, A., McLaughlin, S. Phosphorylation, high ionic
854 strength, and calmodulin reverse the binding of MARCKS to phospholipid vesicles. *Journal of*
855 *Biological Chemistry*. **269** (45), 28214–28219 (1994).

856 22 Sigal, C. T., Zhou, W., Buser, C. A., McLaughlin, S., Resh, M. D. Amino-terminal basic
857 residues of Src mediate membrane binding through electrostatic interaction with acidic
858 phospholipids. *Proceedings of the National Academy of Sciences*. **91** (25), 12253–12257 (1994).

859 23 Bivona, T. G. et al. PKC regulates a farnesyl-electrostatic switch on K-Ras that promotes
860 its association with Bcl-XL on mitochondria and induces apoptosis. *Molecular Cell*. **21** (4), 481–
861 493 (2006).

862 24 Finkielstein, C. V., Overduin, M., Capelluto, D. G. Cell migration and signaling specificity is
863 determined by the phosphatidylserine recognition motif of Rac1. *The Journal of Biological*
864 *Chemistry*. **281** (37), 27317–27326 (2006).

865 25 Bolsover, S. R., Gomez-Fernandez, J. C., Corbalan-Garcia, S. Role of the
866 Ca²⁺/Phosphatidylserine Binding Region of the C2 Domain in the Translocation of Protein Kinase
867 Cα to the Plasma Membrane. *Journal of Biological Chemistry*. **278** (12), 10282–10290 (2003).

868 26 Vance, J. E., Tasseva, G. Formation and function of phosphatidylserine and
869 phosphatidylethanolamine in mammalian cells. *Biochimica et Biophysica Acta*. **1831** (3), 543–554
870 (2013).

871 27 Maeda, K. et al. Interactome map uncovers phosphatidylserine transport by oxysterol-
872 binding proteins. *Nature*. **501** (7466), 257–261 (2013).

873 28 D'Ambrosio, J. M. et al. Osh6 requires Ist2 for localization to ER-PM contacts and efficient
874 phosphatidylserine transport in budding yeast. *Journal of Cell Science*. **133** (11), jcs243733 (2020).

875 29 Manford, A. G., Stefan, C. J., Yuan, H. L., Macgurn, J. A., Emr, S. D. ER-to-plasma membrane
876 tethering proteins regulate cell signaling and ER morphology. *Developmental Cell*. **23** (6), 1129–
877 1140 (2012).

878 30 Collado, J. et al. Tricalbin-mediated contact sites control ER curvature to maintain plasma
879 membrane integrity. *Developmental Cell*. **51** (4), 476–487 e477 (2019).

880 31 Hoffmann, P. C. et al. Tricalbins contribute to cellular lipid flux and form curved ER-PM
881 contacts that are bridged by rod-shaped structures. *Developmental Cell*. **51** (4), 488–502 e488
882 (2019).

883 32 Lipp, N. F. et al. An electrostatic switching mechanism to control the lipid transfer activity
884 of Osh6p. *Nature Communications*. **10** (1), 3926 (2019).

885 33 Chung, J. et al. INTRACELLULAR TRANSPORT. PI4P/phosphatidylserine countertransport
886 at ORP5- and ORP8-mediated ER-plasma membrane contacts. *Science*. **349** (6246), 428–432
887 (2015).

888 34 Sohn, M. et al. PI(4,5)P₂ controls plasma membrane PI4P and PS levels via ORP5/8

recruitment to ER-PM contact sites. *The Journal of Cell Biology*. **217** (5), 1797–1813 (2018).

35 Ghai, R. et al. ORP5 and ORP8 bind phosphatidylinositol-4, 5-bisphosphate (PtdIns(4,5)P₂) and regulate its level at the plasma membrane. *Nature Communications*. **8** (1), 757 (2017).

36 Sohn, M. et al. Lenz-Majewski mutations in PTDSS1 affect phosphatidylinositol 4-phosphate metabolism at ER-PM and ER-Golgi junctions. *Proceedings of the National Academy of Sciences of the United States of America*. **113** (16), 4314–4319 (2016).

37 Kattan, W. E. et al. Targeting plasma membrane phosphatidylserine content to inhibit oncogenic KRAS function. *Life Science Alliance*. **2** (5), e201900431 (2019).

38 Du, X., Turner, N., Yang, H. The role of oxysterol-binding protein and its related proteins in cancer. *Seminars in Cell & Developmental Biology*. **81**, 149–153 (2018).

39 Galmes, R. et al. ORP5/ORP8 localize to endoplasmic reticulum-mitochondria contacts and are involved in mitochondrial function. *EMBO reports*. **17** (6), 800–810 (2016).

40 Du, X. et al. ORP5 localizes to ER-lipid droplet contacts and regulates the level of PI(4)P on lipid droplets. *The Journal of Cell Biology*. **219** (1), e201905162 (2020).

41 Wang, H. et al. ORP2 delivers cholesterol to the plasma membrane in exchange for phosphatidylinositol 4, 5-bisphosphate (PI(4,5)P₂). *Molecular Cell*. **73** (3), 458–473 e457 (2019).

42 Kay, J. G., Grinstein, S. Sensing phosphatidylserine in cellular membranes. *Sensors (Basel)*. **11** (2), 1744–1755 (2011).

43 Moser von Filseck, J., Vanni, S., Mesmin, B., Antonny, B., Drin, G. A phosphatidylinositol-4-phosphate powered exchange mechanism to create a lipid gradient between membranes. *Nature Communications*. **6**, 6671 (2015).

44 Wills, R. C., Goulden, B. D., Hammond, G. R. V. Genetically encoded lipid biosensors. *Molecular Biology of the Cell*. **29** (13), 1526–1532 (2018).

45 Raychaudhuri, S., Im, Y. J., Hurley, J. H., Prinz, W. A. Nonvesicular sterol movement from plasma membrane to ER requires oxysterol-binding protein-related proteins and phosphoinositides. *The Journal of Cell Biology*. **173** (1), 107–119 (2006).

46 Lenoir, M. et al. Structural basis of wedging the Golgi membrane by FAPP pleckstrin homology domains. *EMBO reports*. **11** (4), 279–284 (2010).

47 Liu, Y., Kahn, R. A., Prestegard, J. H. Interaction of Fapp1 with Arf1 and PI4P at a membrane surface: an example of coincidence detection. *Structure*. **22** (3), 421–430 (2014).

48 Shao, C., Novakovic, V. A., Head, J. F., Seaton, B. A., Gilbert, G. E. Crystal structure of lactadherin C2 domain at 1.7 Å resolution with mutational and computational analyses of its membrane-binding motif. *The Journal of Biological Chemistry*. **283** (11), 7230–7241 (2008).

49 Lipp, N. F., Ikhlef, S., Milanini, J., Drin, G. Lipid exchangers: cellular functions and mechanistic links with phosphoinositide metabolism. *Frontiers in Cell and Developmental Biology*. **8**, 663 (2020).


50 Venditti, R. et al. Molecular determinants of ER-Golgi contacts identified through a new FRET-FLIM system. *The Journal of Cell Biology*. **218** (3), 1055–1065 (2019).




51 Pemberton, J. G. et al. Defining the subcellular distribution and metabolic channeling of phosphatidylinositol. *The Journal of Cell Biology*. **219** (3), doi: 10.1083/jcb.201906130 (2020).


52 Nakanishi, H., de los Santos, P. & Neiman, A. M. Positive and negative regulation of a SNARE protein by control of intracellular localization. *Molecular Biology of the Cell*. **15** (4), 1802–1815 (2004).


53 Maekawa, M., Yang, Y., Fairn, G. D. Perfringolysin O theta toxin as a tool to monitor the

933 distribution and inhomogeneity of cholesterol in cellular membranes. *Toxins*. **8** (3), 67 (2016).
934
935












Table 1. Volumes of lipid stock solutions to be mixed for liposome preparation

		Lipid composition (mol/mol)	Lipid			
			DOPC (25 mg/mL)	POPS (10 mg/mL)	16:0 Liss Rhod-PE (1 mg/mL)	C16:0/C16:0-PI(4)P (1 mg/mL)
Extraction assays	Liposome name					
	Liposome 2 mol% PS	PC/PS 98/2	247 µL	12.5 µL		
	Liposome 2 mol% PI(4)P	PC/PI(4)P 98/2	247 µL			153 µL
	DOPC liposome	PC 100	252 µL			
	L _A	PC/PS/Rhod-PE 93/5/2	234 µL	31.4 µL	200 µL	
	L _A without PS	PC/Rhod-PE 98/2	247 µL		200 µL	
Transport assays	L _B	DOPC/PI(4)P 95/5	237 µL			383 µL
	L _B without PI(4)P	PC 100	252 µL			
	L _{A-Eq}	DOPC/PS/PI(4)P/Rhod-PE 93/2.5/2.5/2	234 µL	15.7 µL	200 µL	191 µL
	L _{B-Eq}	DOPC/PS/PI(4)P 95/2.5/2.5	239 µL	15.7 µL		191 µL

Name of Material/ Equipment	Company	Catalog Number	Comments/Description
L-cysteine ≥97 % (FG)	Sigma	W326305-100G	Prepare a 10 mM L-cysteine stock solution in water. Aliquots are stored at -20 °C
2 mL Amber Vial, PTFE/Rub Lnr, for lipids storage in CHCl ₃	Wheaton	W224681	
4 mm-diameter glass beads	Sigma	Z265934-1EA	FPLC
50 mL conical centrifuge tube	Falcon		
ÄKTA purifier	GE healthcare		
Aluminium foil			
Amicon Ultra-15 with a MWCO of 3 and 10 kDa	Merck	UFC900324, UFC901024	Prepare a 50 mg/mL stock solution with filtered and sterilized water and store it at -20 °C.
Amicon Ultra-4 with a MWCO of 3 and 10 kDa	Merck	UFC800324, UFC801024	
Ampicillin			
Bestatin	Sigma	B8385-10mg	
BL21 Gold Competent Cells	Agilent		Dissolve 1 mg of C16:0/C16:0-PI(4)P powder in 250 µL of MeOH and 250 µL of CHCl ₃ . Then complete with CHCl ₃ to 1 mL. The solution must become clear.
C16:0 Liss (Rhod-PE) in CHCl ₃ (1 mg/mL)	Avanti Polar Lipids	810158C-5MG	
C16:0/C16:0-PI(4)P			
	Echelon Lipids	P-4016-3	
C16:0/C18:1-PS (POPS) in CHCl ₃ (10 mg/mL)	Avanti Polar Lipids	840034C-25mg	Prepare 10 mM CaCl ₂ stock solution in water.
C18:1/C18:1-PC (DOPC) in CHCl ₃ (25 mg/mL)	Avanti Polar Lipids	850375C-500mg	
CaCl ₂	Sigma		
Cell Disruptor	Constant Dynamics		
Chloroform (CHCl ₃) RPE-ISO	Carlo Erba	438601	Prepare 1 M DTT stock solution in Milli-Q water. Prepare 1 mL aliquots and store them at -20 °C.
Complete EDTA-free protease inhibitor cocktail	Roche	5056489001	
Deionized (Milli-Q) water			
Dimethylformamide (DMF), anhydrous, >99% pure			
DNAse I Recombinant, RNase free, in powder	Roche	10104159001	Prepare 1 M IPTG stock solution in Milli-Qwater. Prepare 1 mL aliquots and store them at -20 °C.
DTT	Euromedex	EU0006-B	
Econo-Pac chromatography columns (1.5 × 12 cm).	Biorad	7321010	
Electroporation cuvette 2 mm	Ozyme	EP102	
Electroporator Eppendorf 2510	Eppendorf		Spinning the batcerial lysates
Fixed-Angle Rotor Ti45 and Ti45 tubes	Beckman		
Glass-syringes (10, 25, and 50 µL) for fluorecence experiment	Hamilton		
Glass-syringes (25 , 100, 250, 500, and 1000 µL) to handle lipid stock solutions	Hamilton	1702RNR, 1710RNR, 1725RNR, 1750RN type3, 1001RN	
Glutathione Sepharose 4B beads	GE Healthcare	17-0756-05	Prepare a 2 M MgCl ₂ solution. Filter the solution using a 0.45 µm filter.
Glycerol (99% pure)	Sigma	G5516-500ML	
Hemolysis tubes with a cap			
HEPES , >99 % pure	Sigma	H3375-500G	
Illustra NAP 10 desalting column	GE healthcare	GE17-0854-02	Prepared with milli-Q water and autoclaved.
Isopropyl β-D-1-thiogalactopyranoside (IPTG)	Euromedex	EU0008-B	
K-Acetate	Prolabo	26664.293	
Lennox LB Broth medium without glucose			
Liquid nitrogen	Linde		Prepare a 2 M MgCl ₂ solution. Filter the solution using a 0.45 µm filter.
Methanol (MeOH) ≥99.8%	VWR	20847.24	
MgCl ₂			
	Sigma		
Microplate 96 Well PS F-Botom Black Non-Binding	Greiner Bio-one	655900	
Mini-Extruder with two 1 mL gas-tight Hamilton syringes	Avanti Polar Lipids	610023	
Monochromator-based fluorescence plate reader	TECAN	M1000 Pro	

N,N'-Dimethyl-N-(iodoacetyl)-N'-(7-Nitrobenz-2-Oxa-1,3-Diazol-4-yl)Ethylenediamine) (IANBD Amide)

NaCl	Molecular Probes Sigma	S3014-1KG	
PBS			137 mM NaCl, 2.7 mM KCl, 10 mM NaH ₂ PO ₄ , 1.8 mM KH ₂ PO ₄ , autoclaved and stored at 4 °C.
Pear-shaped glass flasks (25 mL, 14/23, Duran glass)	Duran Group		
Pepstatin	Sigma	p5318-25mg	
pGEX-C2 _{LACT} plasmid			Available on request from our lab
pGEX-PH _{FAPP} plasmid			Available on request from our lab
Phenylmethylsulfonyl fluoride (PMSF) ≥98.5% (GC)	Sigma	P7626-25g	
Phosphoramidon	Sigma	R7385-10mg	Prepare a 200 mM PMSF stock solution in isopropanol
Polycarbonate filters (19 mm in diameter) with pore size of 0.2 μm	Avanti Polar Lipids	610006	
Poly-Prep chromatography column (with a 0-2 mL bed volume and a 10 mL reservoir)	Biorad	7311550	
Prefilters (10 mm in diameter).	Avanti Polar Lipids	610014	
PyMOL	http://pymol.org/		Construction of the 3D models of the proteins (Figure 1A)
Quartz cuvette for UV/visible fluorescence (minimum volume of 600 μL)	Hellma		
Quartz cuvettes	Hellma		
Refrigerated centrifuge	Eppendorf	5427R	
Rotary evaporator	Buchi	B-100	
Screw-cap microcentrifuge tubes (1.5 mL)	Sarsted		
Small magnetic PTFE stirring bar (5 × 2 mm)			
Snap-cap microcentrifuge tubes (0.5, 1, and 2 mL)	Eppendorf		fluorescent stain to detect protein in SDS-PAGE gel
SYPRO orange			
Thermomixer	Starlab		
THROMBIN, FROM HUMAN PLASMA	Sigma	10602400001	
Tris, ultra pure	MP	819623	
Ultracentrifuge L90K	Beckman		
UV/Visible absorbance spectrophotometer	SAFAS		
UV/visible spectrofluorometer with a temperature-controlled cell holder and stirring device	Jasco or Shimadzu	Jasco FP-8300 or Shimadzu RF-5301PC	
Vacuum chamber			
Water bath	Julabo		
XK 16/70 column packed with Sephacryl S200HR	GE healthcare		

Dissolve 25 mg of IANBD in 2.5 mL of dimethylsulfoxide (DMSO) and prepare 25 aliquot of 100 μL in 1.5 mL screw-cap tubes. Do not completely screw the cap. Then, remove DMSO in a freeze-dryer to obtain 1 mg of dry IANBD *per* tube. Tubes are closed and stored at -20 °C in the dark.

137 mM NaCl, 2.7 mM KCl, 10 mM NaH₂PO₄, 1.8 mM KH₂PO₄, autoclaved and stored at 4 °C.

Available on request from our lab

Available on request from our lab

Prepare a 200 mM PMSF stock solution in isopropanol

Construction of the 3D models of the proteins (Figure 1A)

fluorescent stain to detect protein in SDS-PAGE gel

Dissolve 20 units in 1 mL of milli-Q water and prepare 25 μL aliquots in 0.5 mL Eppendorf tubes. Then freeze and store at -80 °C.

Guillaume Drin
CNRS, Institut de Pharmacologie Moléculaire et Cellulaire
660 route des lucioles, 06560 Valbonne, France

Manuscript 62177_R2

JOVE,
Dr. Nam Nguyen
Manager of Review

Dr. Benjamin Werth
Editor

Valbonne, February 4, 2021

Dear editors,

Please find enclosed a second revised version of our manuscript entitled “Fluorescence-based measurements of phosphatidylserine/phosphatidylinositol 4-phosphate exchange between membranes”. We are delighted to read that both reviewers consider the manuscript suitable for publication. We have addressed their respective points by making textual changes in red in the manuscript and by revising Figure 4 and 5 as detailed in the point-by-point response letter.

Other textual changes: We have corrected some typos in the manuscript, added a few details, and corrected some inversions of terms in the Protocol section and Figure captions both referring to experiments shown in Figure 4A and Figure 5A (right panel). We have also rewritten the step 3 and 4 of Section 1.3 to better explain a calculation method. All these corrections are in blue. We have also reformatted the References by spelling out the journal names according to JOVE guidelines. We have modified the Protocol section to remove the notes.

We hope that the revised version of our manuscript is now suitable for publication in *JOVE*.

Sincerely yours,

Guillaume Drin



Point-by-point response to reviewers

Reviewer #1:

Manuscript Summary:

This very well written and explain manuscript describes in detail a protocol for the use of PS and PI4P recognizing protein domains (C2 and PH domains respectively) that have been engineered to contain one surface accessible cysteine residue to which has been conjugated a fluorescent 7-nitrobenoxadiazole group. This is used to detect the presence of the protein when bound to phospholipid vesicle membranes that contain the appropriate lipid. When a lipid transfer protein is added either PS or PI4P is removed from the membrane and the change in signal from the NBD group monitored to show lipid removal from the donor vesicle. In a similarly deigned FRET assay, a rhodamine-PE is used as the quenching fluorophore for the NBD group, and time-resolved transfer assays can be recorded. This is a cleverly designed assay system that should be a of general use to investigators interested in lipid transfer process driven by PI4P gradients. The text instructions are quite detailed, but easy to read. I have only minor cosmetic suggestions and word substitutions. We would like to thank the reviewer for these very nice comments on our manuscript.

Minor Concerns:

Line 57; Perhaps change "compose" to "comprise"

We think it is maybe better to change "compose" into "constitute".

Line 61; "repartition" works just as well as "partition"

Done.

Line 72; Maybe replace "encapsulates" with "binds"

Usually we prefer to use the verb "encapsulate" (or "trap" or "accommodate" instead of "bind" to clearly indicate that a LTP captures a lipid inside a pocket and is not a peripheral protein that only recognizes the polar head of a lipid. However, as we mention in the same paragraph what is the mode of binding between a LTP and its ligands, we agree that the verb "bind" is not ambiguous and can be used.

Line75; "We found that Osh6p and Osh7p can trap an alternative PI(4)P lipid ligand." Perhaps re-write to say "We found that Osh6p and Osh7p can trap PI(4)P as an alternative lipid ligand."

We agree. The sentence has been corrected

Line 85; "ligand" to "ligands"

Done.

Line 114; "We reengineered this domain to include a single solvent-exposed cysteine, labelled, with a NBD group, near the PI(4)P binding site." Perhaps change to: "We reengineered this domain to include a single solvent-exposed cysteine, which canbe labelled with a NBD group, near the PI(4)P binding site."

We agree, this is more precise. This correction has been made.

Line 120: "If PS is massively extracted by ..." to "If PS is nearly completely extracted by..."

We agree. The correction has been done.

Line 153: "Many steps must be carried OUT on ice ..."

The correction has been made.

Line 155: Omit "good".

Done.

Line 156: Omit second "sufficient"

Done.

Line 311: "Size-exclusion chromatography is performed on a Sephacryl S200HR column,.. " Size? Grams of gel? Dimensions of column?

We agree. These details were completely necessary. We now indicate the dimensions of the column and the volume of resin packed in the column according to supplier's guidelines.

Line 320: "Laemmli SAMPLE buffer"

We agree.

Line 358 to 426 was highlighted in yellow in my review copy. I'm not certain why.

A part of the manuscript was highlighted in yellow according to JOVE guidelines to mention that this is the one which must be converted into a script for the video.

Line 373 and 400: Change "bandwidth" to "slit width"

We prefer to keep the term "bandwidth" because this is the parameter that appears in the two software that pilot the fluorescence plate reader and the fluorimeter that are used for the experiments and will be shown in the video

References: Is the use of "et al." for when there are more than four authors?

We have formatted the references according to JOVE guidelines, using the provide EndNote style file

Ref 28 has now been published.

Absolutely correct. This is now corrected

Figure 1 was a bit blurry in my review copy.

Likely due to a loose conversion of the original Tiff file into a PDF file. We have increased the resolution of all the Figures.

Figure 4b: Lines in the legend do not seem to have the color of the lines in the graph

Figure 5b: Lines in the legend do not seem to have the color of the lines in the graph

GrapPad Prism automatically assigns the lines of the legend box with the color used to draw the curves. Maybe this change of color is due to a conversion problem. The legend lines are now thicker, also to respond to a Reviewer #2's comment. We now hope that the color correspondence between the lines of the legend box and the curves is easier to appreciate.

Page 35: Comments/Description It wasn't clear to me where these were mentioned in the text.

I also had several (apparently) blank pages with and a couple with vestigial text (p 38 and 40). I'm not certain what happened here.

Page 35-40 result from the conversion of a separated Excel file ,in which the different reagents used in our protocols are listed, and its merging with the manuscript file.

Reviewer #2:

Manuscript Summary:

The manuscript elegantly summarizes the protocol of in vitro PS/PI4P extraction as well as PS/PI4P counter transport between liposomes. The fluorescence probes that the authors developed are quite useful tools to monitor the PS/PI4P transport, and the similar strategy could be adapted to the other lipid molecules. The manuscript is written in a very clear manner and is useful for many readers, so it deserves to be published. I have very minor concerns to be addressed before publication.

We are grateful to Reviewer 2 for his/her very kind comments on our manuscript

Minor Concerns:

I was wondering how long we should incubate the sample when measuring the extraction of PS and PI4P? Do you measure the spectrum shown in Figure 3 right after adding the material? Or, do you measure several samples with different incubation times at the same time? I thought it would be more helpful to add this kind of information.

PS and PI(4)P extraction processes measured with Osh6p are extremely rapid and extraction is completed in less than 2 min. This is what we observed when we first devised this assay using a standard fluorimeter at that time (Moser et al., 2015). For the experiments with a fluorescence plate reader, as it takes a few minutes to load the different wells and for the plate to reach the thermal equilibrium once in the machine, the extraction is completed when the reading of the plate is launched. We now indicate in one sentence that after an incubation time of 5 minutes, the extraction is completed, at least in the case of Osh6p.

Are the spectra shown in right panels of Fig. 3 normalized by subtracting the liposome only signals?

This is entirely exact. We now indicate in the caption of Figure 3 that the spectra measured with each NBD-labelled probe are corrected by subtracting the background signal of liposomes alone.

Lines next to no PI4P or no PS, etc (Fig. 4B and 5B) are too thin and difficult for me to see the color. I thought it would be easier to read if they were a little bolder.

We have corrected the Figure 4 and 5 to increase the thickness of the lines in the legend box and to more clearly visualize the color of each of these lines.