

Journal of Visualized Experiments

Large-scale Multi-omics Genome-wide Association Studies (Mo-GWAS): Guidelines for sample preparation and normalization

--Manuscript Draft--

Article Type:	Methods Article - JoVE Produced Video
Manuscript Number:	JoVE62732R2
Full Title:	Large-scale Multi-omics Genome-wide Association Studies (Mo-GWAS): Guidelines for sample preparation and normalization
Corresponding Author:	Saleh Alseekh MPIMP Potsdam, GERMANY
Corresponding Author's Institution:	MPIMP
Corresponding Author E-Mail:	Alseekh@mpimp-golm.mpg.de
Order of Authors:	Mustafa Bulut Alisdair Fernie Saleh Alseekh
Additional Information:	
Question	Response
Please indicate whether this article will be Standard Access or Open Access.	Open Access (US\$4,200)
Please specify the section of the submitted manuscript.	Biochemistry
Please indicate the city, state/province, and country where this article will be filmed . Please do not use abbreviations.	Potsdam, Germany
Please confirm that you have read and agree to the terms and conditions of the author license agreement that applies below:	I agree to the Author License Agreement
Please provide any comments to the journal here.	
Please indicate whether this article will be Standard Access or Open Access.	Open Access (\$3900)
Please confirm that you have read and agree to the terms and conditions of the video release that applies below:	I agree to the Video Release

TITLE:

Large-scale Multi-omics Genome-wide Association Studies (Mo-GWAS): Guidelines for Sample Preparation and Normalization

AUTHORS AND AFFILIATIONS:

Mustafa Bulut¹, Alisdair R. Fernie^{1,2}, Saleh Alseekh^{1,2}

¹Max-Planck-Institute of Molecular Plant Physiology, Am Mühlenberg 1, 14476 Potsdam-Golm, Germany

²Center of Plant Systems Biology and Biotechnology, 4000 Plovdiv, Bulgaria

Email addresses of co-authors:

Mustafa Bulut (bulut@mpimp-golm.mpg.de)

Alisdair R. Fernie (fernie@mpimp-golm.mpg.de)

Corresponding author:

Saleh Alseekh (alseekh@mpimp-golm.mpg.de)

SUMMARY:

In this protocol, we present an optimized workflow, which combines an efficient and fast sample preparation of many samples. In addition, we provide a step-by-step guide to reduce analytical variations for high-throughput evaluation of metabolic GWAS studies.

ABSTRACT:

Both gas chromatography-mass spectrometry (GC-MS) and liquid chromatography-mass spectrometry (LC-MS) are widely used metabolomics approaches used to detect and quantify hundreds of thousands of metabolite features. However, the application of these techniques to a large number of samples is subject to more complex interactions, particularly for genome-wide association studies (GWAS). This paper describes an optimized metabolic workflow, which combines an efficient and fast sample preparation and the analysis of a large number of samples for legume crop species. This slightly modified extraction method was initially developed for the analysis of plant and animal tissues and based on extraction in methyl tert-butyl ether: methanol solvent to allow the capture of polar and lipid metabolites. In addition, this paper provides a step-by-step guide for reducing analytical variations, which are essential for the high-throughput evaluation of metabolic variance in GWAS studies.

INTRODUCTION:

Large-scale “omics” approaches have enabled the analysis of complex biological systems¹⁻³ and further understanding of the link between genotypes and the resulting phenotypes⁴. Metabolomics using ultra-high-performance liquid chromatography-mass spectrometry (UHPLC-MS) and GC-MS enabled the detection of a plethora of metabolite features, of which only some are annotated to a certain degree, resulting in a high proportion of unknown metabolites. Complex interactions can be explored by combining large-scale metabolomics with the underlying genotypic variation of a diverse population⁵. However, handling large sample sets is inherently associated with analytical variations, distorting the evaluation of metabolic variance for further downstream processes. Specifically, major issues leading to analytical variations are based on machine performance and instrumental drift over time⁶.

The integration of batch-to-batch variation is challenging and especially problematic when analyzing large-scale structured plant populations. Multiple normalization procedures were suggested to correct for non-biological variations, e.g., the usage of internal, external, and isotope-labeled internal standards to correct for analytical errors, which is inherently associated with known problems and pitfalls⁷⁻¹⁰.

In addition to analytical variation, the choice of extraction protocols generally varies depending on the analytical method. Ultimately, it is essential to reduce material and labor costs as well as the necessity of using several aliquots of the same sample for various analytical processes by performing phase separation-based extraction methods. These methods were first introduced using chloroform: methanol/water solvents to fractionate polar and hydrophobic compounds¹¹.

This protocol describes a fast and high-throughput protocol for a multi-omics platform to analyze legume metabolomes and lipidomes at the tissue level. Further, it shows how those datasets can be appropriately corrected for analytical variation and normalized before integrating genotypic information to detect metabolite quantitative trait loci (QTL) by performing GWAS.

PROTOCOL:

1. Experimental design and plant cultivation

NOTE: Set up the experiment depending on the experimental hypothesis, e.g., using a large-scale GWAS population decreases the necessity of multiple replicates, as statistical testing will be performed based on the haplotypes of all the individual SNPs instead of the accession. In contrast, multiple replicates are indispensable in other experimental approaches. The following points must be considered while preparing the experiment.

1.1. Include enough biological replicates, depending on the experimental hypothesis.

1.2. Randomize the biological replicates block-wise to reduce local environmental bias during cultivation, e.g., greenhouse, field.

1.3. Ensure proper maintenance of the plant during growth. Treat plants homogeneously to reduce bias.

2. Preparation of biological plant material

2.1. Harvest preparation

2.1.1. Label harvesting tubes (20 mL) containing two 5 mm and two 8 mm diameter metal beads for homogenizing. Fill up a dewar with liquid nitrogen.

NOTE: Plants should be in the vegetative stage for fresh leaf and root tissue harvesting.

2.2. Harvest biological samples by flash-freezing in liquid nitrogen. Harvest as quickly as possible to exclude the influence of circadian oscillation on metabolism during prolonged harvesting durations^{12,13}. Store the harvested fresh leaf and root tissues for further processing at -80 °C.

NOTE: Leaf-cutting to flash-freezing should not take longer than a few seconds as after leaf cleavage, active biological processes would alter metabolic profiles due to wounding. For roots, preclean the roots by washing with water before flash-freezing in liquid nitrogen. Excess water on the root surface should be soaked up with paper tissue. Dried seeds can be stored at room temperature; no freezing in liquid nitrogen is required.

2.3. Grind the tissue using a tissue mixer mill.

2.3.1. Precool the tube holders in liquid nitrogen for a couple of minutes to maintain a low temperature while grinding the tissue.

2.3.2. Transport the biological samples in a nitrogen-containing dewar after taking them out from the -80 °C freezer.

2.3.3. Grind the tissues to obtain homogeneous powder; use 25 Hz for 1 min and repeat after freezing in liquid nitrogen if the tissue is not homogeneously ground.

2.4. For grinding dried seeds, place the seeds in a grinding jar with a 15 mm diameter metal bead. Use the same frequency and time as mentioned in 2.3.3.

NOTE: Clean and precooled mortars and pestles can be used if a tissue mixer mill is unavailable.

2.5. Precool labeled 2 mL safe-lock microcentrifuge tubes. Weigh 50 mg with an error of ± 5 mg of fresh plant material by using an analytical scale. Precool the tools used for transferring plant material in liquid nitrogen. Ensure that plant material stays frozen during the weighing process.

NOTE: Do not expose fresh plant material too long to room temperature as biological processes are activated by increasing temperature, altering metabolic profiles¹⁴.

2.6. Generate additional quality control (QC) samples by pooling a proportion of each sample and weighing 50 mg with an error of ± 5 mg of pooled fresh plant material into precooled 2 mL safe-lock microcentrifuge tubes.

NOTE: At least three QC samples are advised for every 60 samples. The QC samples are essential for the downstream correction, normalization, and analyses.

3. Extraction reagents

3.1. Fresh tissue, e.g., leaves and roots

NOTE: Sample extraction is based on a previously described protocol¹⁵. This protocol has been modified based on present needs, e.g., multiple tissues, different internal standards, and large-scale experiments. Additionally, all volumes and instrument settings mentioned below are adjusted to in-house analytical units. Protocol users should adjust these according to their analytical unit and biological samples, based on test samples.

3.1.1. Extraction mixture 1 (EM1): methyl *tert*-butyl ether (MTBE)/methanol (MeOH) (3:1 v/v)

3.1.1.1. Prepare a mixture of MTBE/MeOH in a 3:1 ratio. For 100 mL of extraction solvent, mix 75 mL of MTBE with 25 mL of MeOH in a clean glass bottle.

NOTE: Solvents should be handled carefully in the fume hood with proper safety equipment.

3.1.1.2. Add 45 μ L of 1,2-diheptadecanoyl-sn-glycero-3-phosphocholine (1 mg/mL in chloroform) as an internal standard for the UHPLC-MS based lipid analysis, 400 μ L of ribitol (1 mg/mL in water) as internal standard for the GC-MS based analysis, and 125 μ L of isovitexin (1 mg/mL in MeOH/water (1:1 v/v)) for UHPLC-MS-based metabolite analysis.

NOTE: The addition of internal standards is necessary for the post-analysis normalization according to analytical needs. As 1 mL of EM1 is needed for each sample, prepare a stock solution according to the experimental sample size, which should be used for the entire experiment. EM1 must be stored at -20 °C. Check for the absence of the used internal standard and overlapping with other compounds in the investigated species. Several standards can be used; the selection of the internal standards in this protocol was based on previous tests using common bean extracts¹⁶.

3.1.2. Extraction mixture 2 (EM2) water/ methanol (MeOH) (3:1 v/v)

3.1.2.1. For 100 mL EM2, add 75 mL of double-distilled water and 25 mL of MeOH in a clean glass bottle.

3.1.2.2. Add 500 μ L of EM2 per sample, and prepare a stock solution according to the experimental sample size, which should be used for the entire experiment. Store EM2 at 4 °C.

3.2. Dried seeds

3.2.1. Extraction mixture 3 (EM3) methanol (MeOH)/ water (7:3 v/v)

3.2.1.1. For 100 mL of EM3, add 70 mL of MeOH and 30 mL of double-distilled water in a clean glass bottle. Prepare 1 mL of EM3 for each sample.

3.2.1.2. Add 400 μ L of ribitol (1 mg/mL in water) as internal standards for the GC-MS-based analysis and 125 μ L of Isoviteixin (1 mg/mL in MeOH/water (1:1 v/v)) for UHPLC-MS-based metabolite analysis.

NOTE: Prepare a stock solution according to the experimental sample size and use it for the entire experiment. Store EM3 at 4 °C.

4. Sample extraction

4.1. Fresh tissue, e.g., leaves and roots

4.1.1. Prepare three 1.5 mL safe-lock microcentrifuge tubes for each sample. Keep EM1 in a -20 °C liquid cooling system. Transfer the fresh samples from the -80 °C freezer to dry ice or liquid nitrogen for transportation. Add 1 mL of precooled EM1 to each 50 mg aliquot and vortex briefly before keeping on ice.

4.1.2. Incubate the samples on an orbital shaker at $800 \times g$ for 10 min at 4 °C.

4.1.3. Sonicate the samples in an ice-cooled sonication bath for 10 min.

4.1.4. Add 500 µL of EM2 using a multichannel pipette to avoid variation in added volumes.

4.1.5. Vortex the samples briefly to mix the extraction mixtures before centrifuging at $11,200 \times g$ for 5 min at 4 °C.

4.1.6. After phase separation occurs, transfer 500 µL of the upper lipid-containing phase to a prelabeled 1.5 mL safe-lock microcentrifuge tube. Remove the rest of the upper phase.

NOTE: Take care while transferring as this upper phase has a high vapor pressure and tends to leak out from the pipette.

4.1.7. Transfer 150 µL and 300 µL of the lower polar and semi-polar metabolite-containing phases in two 1.5 mL safe-lock microcentrifuge tubes used for GC-MS and UHPLC-MS analysis, respectively.

4.1.8. Concentrate all the extracted fractions by letting the solvents evaporate without heating using a vacuum concentrator and store at -20 °C.

4.2. Dried seeds

4.2.1. Prepare two 1.5 mL safe-lock microcentrifuge tubes for each sample. Keep EM3 on ice. Place a 5 mm diameter metal bead in the sample aliquots.

4.2.2. Add 1 mL of EM3 in each 50 mg aliquot and homogenize the samples at 25 Hz for 2–3 min before putting them on ice.

4.2.3. Sonicate the samples in an ice-cooled sonication bath for 10 min.

4.2.4. Vortex the samples briefly before centrifuging at $11,200 \times g$ for 5 min at 4 °C.

4.2.5. Transfer 150 μL and 300 μL of the supernatant in two 1.5 mL safe-lock microcentrifuge tubes used for GC-MS and UHPLC-MS analysis, respectively.

4.2.6. Concentrate all extracted fractions by letting the solvents evaporate at room temperature using a vacuum concentrator and store at $-20\text{ }^{\circ}\text{C}$.

NOTE: Based on experience, users are advised to perform step 4.2 for semi-polar metabolites and derivatized metabolite analysis in dried seeds. Perform extraction step 4.1 for dried seed lipid analysis.

5. Analysis of lipids using UHPLC-MS

5.1. Re-suspend the dried lipid fractions in 250 μL of acetonitrile:2-propanol (7:3, vol/vol).

5.2. Sonicate the lipid phase for 5 min, centrifuge at $11,200 \times g$ for 1 min.

5.3. Transfer 90 μL of the supernatant to a glass vial for LC-MS.

5.4. Inject 2 μL of the extracts into the LC-MS.

5.5. Perform lipid fractionation on a reversed-phase C_8 column held at $60\text{ }^{\circ}\text{C}$ running with a flow of 400 $\mu\text{L}/\text{min}$ with gradual changes of eluent A and B as shown in **Table 1**. Acquire the mass spectra in positive ionization mode with a mass range of 150–1,500 m/z .

5.6. Include several QC samples in all daily batches and a blank to ensure correction for analytical variation. Randomize samples properly block-wise in sequential order.

6. Analysis of polar and semi-polar metabolites using UHPLC-MS

6.1. Re-suspend the dried polar phase in 180 μL of UHPLC-grade methanol: water (1:1 v/v).

6.2. Sonicate the polar phase for 2 min, centrifuge at $11,200 \times g$ for 1 min.

6.3. Transfer 90 μL of the supernatant to a glass vial for LC-MS.

6.4. Inject 3 μL of the extracts into the LC-MS.

6.5. Perform metabolite fractionation on a reverse phase C_{18} column held at $40\text{ }^{\circ}\text{C}$ running with a flow of 400 $\mu\text{L}/\text{min}$ with gradual changes of eluent A and B as shown in **Table 1**. Acquire the mass spectra in a mass range of 100–1,500 m/z in a full MS scan and all ion fragmentation (AIF) induced by high-energy collisional dissociation (HCD) of 40 keV.

NOTE: Use both ionization modes. However, due to limited capacity while running large numbers of samples, run test samples in both ionization modes to determine the preferred ionization mode.

6.6. Include several QC samples in all daily batches and a blank to ensure correction for analytical variation. Randomize samples properly block-wise in sequential order.

6.7. Run a pooled QC in data-dependent MS² in both negative and positive ionization modes. Use the obtained mass spectra in a later step (8.5) for annotation.

7. Analysis of derivatized Metabolites using GC-MS^{17,18}

NOTE: The analysis of derivatized metabolites is based on a previously described protocol¹⁷. Handle all derivatization reagents in the fume hood. Ensure that *N*-methyl-*N*-(trimethylsilyl)trifluoroacetamide (MSTFA) does not get in contact with water and humidity.

7.1. Derivatization reagent 1 (DR1)

7.1.1. Dissolve methoxyamine hydrochloride in pyridine to obtain a concentration of 30 mg/mL of DR1. Use 60 µL of DR1 for each sample. Prepare a stock solution according to the sample size, and store at room temperature.

7.2. Derivatization reagent 2 (DR2)

7.2.1. Dissolve MSTFA with 20 µL of fatty acid methyl esters (FAMES) per 1 mL of MSTFA. Use 70 µL of DR2 for each sample. Prepare a stock solution according to the sample size. Store MSTFA at 4 °C and the FAMES at -20 °C.

NOTE: FAMES include methylcaprylate, methyl pelargonate, methylcaprate, methyl laurate, methylmyristate, methylpalmitate, methylstearate, methyleicosanoate, methyl docosanoate, lignoceric acid methyl ester, methylhexacosanoate, methyl octacosanoate, and triacontanoic acid methylester, which are dissolved in CHCl₃ at a concentration of 0.8 µL/mL or 0.4 mg/mL for liquid or solid standards, respectively.

7.3. Re-dry the pellet from the polar phase (stored at -80 °C) using a vacuum concentrator for 30 min to avoid any interference of H₂O originating during the storage with the solvents used for the downstream derivatization.

7.4. Add 40 µL of DR1.

7.5. Shake the samples at 950 × *g* for 2 h at 37 °C using an orbital shaker, followed by a short spin-down of the liquid.

7.6. Add 70 µL of DR2.

7.7. Shake again at 950 × *g* for 30 min at 37 °C using an orbital shaker.

7.8. Centrifuge briefly at room temperature before transferring 90 µL into glass vials for GC-MS analysis.

7.9. Inject 1 μL to GC-MS splitless mode, depending on the metabolite concentrations, with a constant helium carrier gas flow of 2 mL/min. Injection temperature is set to 230 °C using a 30-m MDN-35 capillary column.

NOTE: Additional information, e.g., temperature gradient, can be found in **Table 1**. The mass range is set to 70–600 m/z with 20 scans/min. Include split modes to enable the quantification of putative overloading compounds, saving costs and time for extract re-derivatization in such cases.

7.10. Include several QC samples in all daily batches and a blank to ensure correction for analytical variation. Randomize samples properly block-wise in sequential order.

8. Chromatogram processing and compound annotation

8.1. Filter chemical noise by defining intensity thresholds. Include all QC samples while processing the chromatograms.

NOTE: For large-scale data, noise filtering is crucial to decrease computing time and processing power.

8.2. Align the chromatograms by defining a retention time shift window. Check the chromatograms from each batch to assess the intra- and inter-batch variation.

8.3. Perform peak detection depending on the peak shape, e.g., height and width for full width at half-maximum (FWHM) calculations.

8.4. Cluster isotopes to reduce redundant signals and filter out singletons.

NOTE: See the **Table of Materials** for details on software used for chromatogram processing. In-depth protocols on how to process chromatograms using various freely available software tools, e.g., MS-DIAL, MetAlign, MzMine, and Xcalibur¹⁹⁻²¹, are provided.

8.5. Use the ddMS² data of a pooled QC sample for compound annotation. Assess the molecular structure by determining the monoisotopic mass and observing common neutral losses, known charged aglycones, and different types of cleavages, e.g., homolytic or heterolytic^{16,22}.

8.6. For reporting metabolite data, follow the recommendation described in Fernie et al. 2011²³.

NOTE: Different computational metabolomics approaches can be used to analyze metabolomics data²⁴⁻²⁶.

9. Normalization of large-scale metabolomics dataset

9.1. Check the distribution of the internal standard (s) and normalize by correcting for the response of single or multiple internal standards.

9.2. Correct the peak intensities obtained from the chromatogram over the exact sample weight by dividing the peak intensities by the aliquoted homogenized sample from step 2.5.

9.3. Correct for intensity drift across multi-batch series. Perform QC-based correction methods such as locally estimated scatterplot smoothing (LOESS)²⁷ using R.

NOTE: Several tools and packages are available to address the drift of the MS performance during the acquisition of the whole batches^{28,29}.

9.4. Ensure normal distribution of traits by data transformation, e.g., Box-Cox transformation³⁰ using the *boxcox ()* function from the R package **MASS** for carrying out GWAS.

9.5. Perform data scaling, e.g., Pareto scaling, for multivariate analysis to ensure proper weighing of low abundant compounds³¹.

NOTE: If possible, perform a recovery assay to avoid matrix effects, e.g., ion suppression¹⁴.

10. Genome-wide association studies (GWAS)³²

10.1. Call single nucleotide polymorphism (SNP) or structural variants (SV) from the genotype-by-sequencing (GBS) or whole-genome sequencing (WGS) data^{33,34}.

10.2. Before GWAS, filter genotypic data for minor allele frequency (MAF) < 5% and missing rate of >10% to avoid low-frequency bias using Tassel³⁵.

10.3. Calculate best linear unbiased predictions (BLUPs) for each normalized feature over the experimental repetitions to eliminate bias originating from environmental factors (random effects) using the R package **lme4**³⁶.

10.4. Use BLUPs of each feature individually to perform GWAS using the **rMVP** package in R³⁷.

NOTE: Each metabolomics feature is viewed here as an individual stand-alone phenotype.

10.5. While performing GWAS, correct for population structure using principal component analysis (PCA) and identity by state (IBS) or vanRaden to minimize confounding effects. Furthermore, consider using a mixed linear model (MLM) or a multi-locus mixed model (MLMM), as mixed models contain fixed and random effects.

11. QTL detection

11.1. Check the SNPs showing significant association, taking the Manhattan plots into consideration, for linkage disequilibrium (LD) calculations to determine the underlying genetic region. Perform LD calculations using the R package **LD heatmap** or Tassel 5.

11.2. Check the associated SNPs for the effect size over the trait by examining the trait levels for statistical changes between haplotypes to find potential causal SNPs, e.g., SNPs leading to an amino acid change in the protein-coding sequence, which could explain the phenotypic variation.

NOTE: As SNP-trait associations do not necessarily yield causal association, it is crucial to determine the genomic region. Compound identity by feature annotation can help immensely in finding the right candidate genes in a specific genomic region.

11.3. Combine all detected QTL associated with certain compounds in a pleiotropic map to underline the genetic regions associated with several traits³⁸, as shown in **Figure 4**.

NOTE: For validation of candidate genes, several approaches can be performed (see the discussion).

REPRESENTATIVE RESULTS:

Successful metabolomics GWAS experiments should begin with a proper experimental design, followed by sample collection, extraction, data acquisition, and processing, as illustrated in **Figure 1**. In this protocol, the MTBE method¹⁵ was used to extract and analyze hundreds of metabolites belong to several compound classes. Presented here are example lipidomic data. Chromatography depends highly on the properties of the utilized column as well as elution buffer mixtures.

Figure 2 shows a chromatogram of a QC sample, indicating the elution pattern of some major lipid classes in this analytical system. The applied gradients for each platform are given in **Table 1**. Strong emphasis was placed on handling systemic errors in large-scale experiments. Performing large-scale metabolomics is inherently associated with systemic errors. For demonstration, we analyzed lipidomic data across several common bean species. **Supplemental Table 1** provides the extracted raw lipidomics data obtained after chromatogram processing using the software indicated in the **Table of Materials**. Following this protocol enabled the circumventing of major issues in dealing with omics data, especially while handling large sample sets. The normalization procedure yields accurate correction of batch-wise analytical errors, as demonstrated in **Figure 3**. Although increasing the numbers of QC samples will increase the power of the normalization, this is not always feasible due to cost and time constraints. For high-throughput metabolomics GWAS with non-targeted metabolic features, it is essential to illustrate higher numbers of trait-genotype association properly. A pleiotropic map³⁸ combining multiple GWAS results could be used to highlight the genomic regions to which several traits are linked (**Figure 4**).

FIGURE AND TABLE LEGENDS:

Figure 1: Flowchart of the metabolomics-based GWAS in plants. Several steps starting from the experimental design up to the detection of QTL are shown in the left panel. In the right panel, multiple figures are shown to support several steps mentioned in the left panel. Starting from the right top, **(1)** a suggested sequence of samples is shown for LC-MS, **(2)** pre- and post-normalized score plots of PCA, including a representative feature distribution pre- and post-processing, with red indicating QC sample intensities, and **(3)** a Manhattan plot with significant associations to which LD and haplotypes were created. Abbreviations: GWAS =

genome-wide association studies; QTL = quantitative trait loci; PCA = principal component analysis; QC = quality control; LD = linkage disequilibrium; MS = mass spectrometry; LC-MS = liquid chromatography-mass spectrometry; GC-MS = gas chromatography-mass spectrometry; LOESS = locally estimated scatterplot smoothing; MLM/MLMM = mixed linear model/multi-locus mixed-model.

Figure 2: Chromatogram processing. Two QC chromatograms (base peak; lipid data) from different batches demonstrate the batch-wise variation for certain lipid classes in the pooled QC samples. Four major lipid classes are indicated with their respective elution windows in the in-house LC-MS system. The chromatograms were exported from MzMine²¹. Abbreviations: QC = quality control; LC-MS = liquid chromatography-mass spectrometry.

Figure 3: Correction of systematic error. Principal component analysis of acquired lipidomic data, pre- (left, raw data) and post-correction for systemic errors (right, batch loess). The lower panels illustrate the feature (Cluster_00005) distribution over the samples ($n=650$) and batches ($n=10$) pre- (left) and post (right)-correction for analytical variation. Abbreviations: PCA = principal component analysis; QC = quality control; LOESS = locally estimated scatterplot smoothing.

Figure 4: Pleiotropic map illustrating the combined GWAS results. The pleiotropic map highlights regions in the whole genome that are associated with several traits. The numbers on the outer rings indicate the corresponding chromosomes. Each circlet represents an individual trait with its significantly associated SNPs. The colors represent different compound classes (grey = compound class 1; green = compound class 2; purple = compound class 3; yellow = compound class 4). In the case of inter-compound class associations with the same genomic region, genes are highlighted. The inner grey circle shows the sum of all significant SNPs associated with a specific genomic position. The associations shown in this figure are artificially generated only for illustration. Abbreviations: GWAS = genome-wide association studies; SNPs = single-nucleotide polymorphisms.

Table 1: Gradient settings for each of the analytical platforms⁷. Abbreviations: lg = linear gradient; cw = column washing; eq = equilibrate; UHPLC-MS = ultra-high-performance liquid chromatography-mass spectrometry; UHPLC-MS/MS = ultra-high-performance liquid chromatography-tandem mass spectrometry; GC-MS = gas chromatography-mass spectrometry. * = percentage value corresponds to eluent A.

Supplemental Table 1: Raw lipidomics data. Indicates the peak intensities for each of the detected clusters over each sample.

DISCUSSION:

Both GC-MS and LC-MS are widely used tools for profiling complex mixtures of various metabolite classes. Handling large datasets with these tools is inherently associated with a non-biological variation, e.g., analytical variation, which interferes and biases the interpretation of the results. This protocol presents a robust and high-throughput extraction pipeline for comprehensive metabolomic and lipidomic profiling to eliminate variation of non-biological origin and conduct large-scale “omics” studies. The volumes and concentrations used in this protocol were adjusted for legume species in different tissues at the vegetative

stage for fresh leaf and root tissue. However, this protocol can be slightly modified and used for large-scale metabolic samples from other plant species as well.

The previously¹⁵ described MTBE-based extractions can be used to analyze derivatized metabolites, semi-polar metabolites, and lipids. This can be expanded for protein and plant hormone extractions³⁹, which were out of the scope of this protocol. Other extraction protocols rely on dichloromethane:ethanol mixtures^{40,41}. Of these extraction protocols, the MTBE:methanol extraction protocol provides a favorable and less hazardous alternative to the existing chloroform-based extraction protocols⁴² and does not result in a protein pellet as an interphase between the polar and lipid phases. Furthermore, MTBE methods have already been used in several studies for various biological samples⁴³⁻⁴⁵.

This protocol discusses several crucial steps that might lead to potential variation for a large sample size, e.g., in prolonged harvesting if not handled properly^{12,13}, extraction¹⁴, and randomization⁴⁶. In addition, there are additional issues that have not been discussed in this protocol that must be considered to ensure a high-quality metabolic data set, e.g., matrix effect and ion suppression¹⁴.

The power of QC-based normalization methods inherently depends on the number of QC samples in each batch. As mentioned earlier, although increasing the number would increase the power, the intra-batch variation of the QCs is relatively marginal compared to inter-batch variation in these analytical systems, as illustrated in **Figure 3**. Overall, there are other QC-based normalization methods, such as systemic error removal using random forest (SERRF), which have been shown to outperform most of the other normalization methods such as batch-wise-ratio, normalization using an optimal selection of multiple internal standards (NOMIS), and probabilistic quotient normalization (PQN)⁴⁷. However, SERRF relies on multiple QC samples in each batch, e.g., every tenth sample, which is not feasible while handling large numbers of samples. The main advantage of QC-based normalization over other data-driven or internal standard-based methods is that it retains the essential biological variation while accommodating unwanted technical variation²⁸. Readers may refer to this review on the handling of variation²⁸.

One main issue in GWAS is the rate of false positives, which originate mostly due to the linkage of causal and non-causal sites^{48,49}. Second, the conservative statistical correction approaches, e.g., Bonferroni and FDR, correct for the number of independent tests, which is not equal to the number of assayed SNPs in GWAS due to the linkage between proximate SNPs^{50,51}. Therefore, the actual number of independent tests is often lower. Another way to reduce the conservative statistical threshold would be to reduce the number of tested SNPs used for GWAS based on linkage decay over certain genomic regions⁵². The GWAS-integrated high-throughput metabolomics platform described in this protocol has a wide range of applications. In particular, it will facilitate improved crop breeding by changing the metabolite/lipid composition for industrially and nutritionally desired levels. Overall, metabolomics has provided an in-depth insight into the genetic architecture of a plethora of metabolites and metabolic diversification that occurred during crop domestication over the last decades, indicating the vast potential of metabolomics-associated breeding⁵³. Several molecular biological approaches for downstream validation include the generation of CRISPR/Cas9 mutant lines⁵⁴, T-DNA insertion lines⁵⁵, stable and/or transient overexpression

lines⁵⁶, VIGS, *ex vivo* metabolomics approaches⁵⁷ next to the conventional approaches in generating F₂ cross population.

Generally, by performing the necessary correction for the analytical variations as described above, several integrated approaches can be performed in addition to GWAS, such as metabolite-metabolite, metabolite-lipid correlation analysis, correlation analysis to other broad phenotypic data to shed light on more complex traits, and/or co-expression analysis to further unravel the basis of biological systems⁵⁸.

ACKNOWLEDGMENTS:

M.B. is supported by the IMPRS-PMPG 'Primary Metabolism and Plant Growth'. A.R.F. and S.A. acknowledge the financial support of the EU Horizon 2020 Research and Innovation Programme, project PlantaSYST (SGA-CSA No. 739582 under FPA No. 664620), and project INCREASE (GA 862862).

DISCLOSURES:

The authors have no conflicts of interest to declare.


REFERENCES:


- 1 Doerr, A. Global metabolomics. *Nature Methods*. **14** (1), 32–32 (2017).
- 2 Fessenden, M. Metabolomics: Small molecules, single cells. *Nature*. **540** (7631), 153–155 (2016).
- 3 Oliver, S. G., Winson, M. K., Kell, D. B., Baganz, F. Systematic functional analysis of the yeast genome. *Trends in Biotechnology*. **16** (9), 373–378 (1998).
- 4 Fiehn, O. Metabolomics—the link between genotypes and phenotypes. *Plant Molecular Biology*. **48** (1), 155–171 (2002).
- 5 Wu, S. et al. Mapping the *Arabidopsis* metabolic landscape by untargeted metabolomics at different environmental conditions. *Molecular Plant*. **11** (1), 118–134 (2018).
- 6 Sysi-Aho, M., Katajamaa, M., Yetukuri, L., Orešič, M. Normalization method for metabolomics data using optimal selection of multiple internal standards. *BMC Bioinformatics*. **8** (1), 93 (2007).
- 7 Chen, M., Rao, R. S. P., Zhang, Y., Zhong, C. X., Thelen, J. J. A modified data normalization method for GC-MS-based metabolomics to minimize batch variation. *SpringerPlus*. **3** (1), 439 (2014).
- 8 Dunn, W. B. et al. Metabolic profiling of serum using Ultra Performance Liquid Chromatography and the LTQ-Orbitrap mass spectrometry system. *Journal of Chromatography B*. **871** (2), 288–298 (2008).
- 9 Fiehn, O. et al. Metabolite profiling for plant functional genomics. *Nature Biotechnology*. **18** (11), 1157–1161 (2000).
- 10 van der Kloet, F. M., Bobeldijk, I., Verheij, E. R., Jellema, R. H. Analytical error reduction using single point calibration for accurate and precise metabolomic phenotyping. *Journal of Proteome Research*. **8** (11), 5132–5141 (2009).
- 11 Folch, J., Lees, M., Stanley, G. H. S. A simple method for the isolation and purification of total lipides from animal tissues. *Journal of Biological Chemistry*. **226** (1), 497–509 (1957).
- 12 Fukushima, A. et al. Impact of clock-associated *Arabidopsis* pseudo-response regulators in metabolic coordination. *Proceedings of the National Academy of Sciences of the United States of America*. **106** (17), 7251–7256 (2009).



- 13 Kerwin, R. E. et al. Network quantitative trait loci mapping of circadian clock outputs identifies metabolic pathway-to-clock linkages in *Arabidopsis*. *The Plant Cell*. **23** (2), 471–485 (2011).
- 14 Tohge, T. et al. From models to crop species: Caveats and solutions for translational metabolomics. *Frontiers in Plant Sciences*. **2**, 61 (2011).
- 15 Salem, M., Bernach, M., Bajdzienko, K., Giavalisco, P. A simple fractionated extraction method for the comprehensive analysis of metabolites, lipids, and proteins from a single sample. *Journal of Visualized Experiments: JoVE*. (124), e55802 (2017).
- 16 Tohge, T., Fernie, A. R. Combining genetic diversity, informatics and metabolomics to facilitate annotation of plant gene function. *Nature Protocols*. **5** (6), 1210–1227 (2010).
- 17 Lisec, J., Schauer, N., Kopka, J., Willmitzer, L., Fernie, A. R. Gas chromatography mass spectrometry-based metabolite profiling in plants. *Nature Protocols*. **1** (1), 387–396 (2006).
- 18 Osorio, S., Do, P. T., Fernie, A. R. in *Plant Metabolomics: Methods and Protocols*. Hardy, N. W., Hall, R. D. (Eds), Humana Press, 101–109 (2012).
- 19 De Vos, R. C. H. et al. Untargeted large-scale plant metabolomics using liquid chromatography coupled to mass spectrometry. *Nature Protocols*. **2** (4), 778–791 (2007).
- 20 Perez de Souza, L., Alseekh, S., Naake, T., Fernie, A. Mass spectrometry-based untargeted plant metabolomics. *Current Protocols in Plant Biology*. **4** (4), e20100 (2019).
- 21 Pluskal, T., Castillo, S., Villar-Briones, A., Orešič, M. MZmine 2: Modular framework for processing, visualizing, and analyzing mass spectrometry-based molecular profile data. *BMC Bioinformatics*. **11** (1), 395 (2010).
- 22 Watson, J. T., Sparkman, D. O. Electron Ionization. in *Introduction to mass spectrometry: Instrumentation, applications and strategies for data interpretation*. John Wiley & Sons, Ltd, 315–448 (2007).
- 23 Fernie, A. R. et al. Recommendations for reporting metabolite data. *The Plant Cell*. **23** (7), 2477 (2011).
- 24 Treutler, H. et al. Discovering regulated metabolite families in untargeted metabolomics studies. *Analytical Chemistry*. **88** (16), 8082–8090 (2016).
- 25 Wang, M. et al. Sharing and community curation of mass spectrometry data with Global Natural Products Social Molecular Networking. *Nature Biotechnology*. **34** (8), 828–837 (2016).
- 26 Naake, T., Fernie, A. R. MetNet: Metabolite network prediction from high-resolution mass spectrometry data in R aiding metabolite annotation. *Analytical Chemistry*. **91** (3), 1768–1772 (2019).
- 27 Chambers, J. M. *Statistical models in S*. CRC Press, Inc. (1991).
- 28 Misra, B. B. Data normalization strategies in metabolomics: Current challenges, approaches, and tools. *European Journal of Mass Spectrometry*. **26** (3), 165–174 (2020).
- 29 Livera, A. M. D. et al. Statistical methods for handling unwanted variation in metabolomics data. *Analytical Chemistry*. **87** (7), 3606–3615 (2015).
- 30 Sakia, R. M. The Box-Cox transformation technique: a review. **41** (2), 169–178 (1992).
- 31 van den Berg, R. A., Hoefsloot, H. C. J., Westerhuis, J. A., Smilde, A. K., van der Werf, M. J. Centering, scaling, and transformations: improving the biological information content of metabolomics data. *BMC Genomics*. **7**, 142 (2006).
- 31 Marees, A. T. et al. A tutorial on conducting genome-wide association studies: Quality control and statistical analysis. *International Journal of Methods in Psychiatric Research*. **27** (2), e1608 (2018).

- 32 Torkamaneh, D., Laroche, J., Bastien, M., Abed, A., Belzile, F. Fast-GBS: a new pipeline for the efficient and highly accurate calling of SNPs from genotyping-by-sequencing data. *BMC Bioinformatics*. **18** (1), 5 (2017).
- 33 Zhao, S., Agafonov, O., Azab, A., Stokowy, T., Hovig, E. Accuracy and efficiency of germline variant calling pipelines for human genome data. *Scientific Reports*. **10** (1), 20222 (2020).
- 34 Bradbury, P. J. et al. TASSEL: software for association mapping of complex traits in diverse samples. *Bioinformatics*. **23** (19), 2633–2635 (2007).
- 35 Bates, D., Mächler, M., Bolker, B., Walker, S. Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*. **67** (1), doi: 10.18637/jss.v067.i01 (2015).
- 36 Yin, L. et al. rMVP: A memory-efficient, visualization-enhanced, and parallel-accelerated tool for genome-wide association study. *Genomics, Proteomics & Bioinformatics*. doi: [10.1016/j.gpb.2020.10.007](https://doi.org/10.1016/j.gpb.2020.10.007) (2021).
- 37 Kanai, M. et al. Genetic analysis of quantitative traits in the Japanese population links cell types to complex human diseases. *Nature Genetics*. **50** (3), 390–400 (2018).
- 38 Salem, M. A. et al. An improved extraction method enables the comprehensive analysis of lipids, proteins, metabolites and phytohormones from a single sample of leaf tissue under water-deficit stress. *Plant Journal: for Cell and Molecular Biology*. **103** (4), 1614–1632 (2020).
- 39 Balcke, G. U. et al. Multi-omics of tomato glandular trichomes reveals distinct features of central carbon metabolism supporting high productivity of specialized metabolites. *The Plant Cell*. **29** (5), 960–983 (2017).
- 40 Leonova, T. et al. Does protein glycation impact on the drought-related changes in metabolism and nutritional properties of mature pea (*Pisum sativum* L.) seeds? *International Journal of Molecular Sciences*. **21** (2), 567 (2020).
- 41 Alfonsi, K. et al. Green chemistry tools to influence a medicinal chemistry and research chemistry based organisation. *Green Chemistry*. **10** (1), 31–36 (2008).
- 42 Bozek, K. et al. Organization and evolution of brain lipidome revealed by large-scale analysis of human, chimpanzee, macaque, and mouse tissues. *Neuron*. **85** (4), 695–702 (2015).
- 43 Delgado, R., Muñoz, Y., Peña-Cortés, H., Giavalisco, P., Bacigalupo, J. Diacylglycerol activates the light-dependent channel TRP in the photosensitive microvilli of *Drosophila melanogaster* photoreceptors. *The Journal of Neuroscience*. **34** (19), 6679 (2014).
- 44 Sharma, D. K. et al. UPLC-MS analysis of *Chlamydomonas reinhardtii* and *Scenedesmus obliquus* lipid extracts and their possible metabolic roles. *Journal of Applied Phycology*. **27** (3), 1149–1159 (2015).
- 45 Dunn, W. B., Wilson, I. D., Nicholls, A. W., Broadhurst, D. The importance of experimental design and QC samples in large-scale and MS-driven untargeted metabolomic studies of humans. *Bioanalysis*. **4** (18), 2249–2264 (2012).
- 46 Fan, S. et al. Systematic error removal using random forest for normalizing large-scale untargeted lipidomics data. *Analytical Chemistry*. **91** (5), 3590–3596 (2019).
- 47 Larsson, S. J., Lipka, A. E., Buckler, E. S. Lessons from Dwarf8 on the strengths and weaknesses of structured association mapping. *PLOS Genetics*. **9** (2), e1003246 (2013).
- 48 Platt, A., Vilhjálmsdóttir, B. J., Nordborg, M. Conditions under which genome-wide association studies will be positively misleading. *Genetics*. **186** (3), 1045–1052 (2010).
- 49 Nyholt, D. R. A simple correction for multiple testing for single-nucleotide polymorphisms in linkage disequilibrium with each other. *American Journal of Human Genetics*. **74** (4), 765–769 (2004).


- 50 Teo, Y. Y. Common statistical issues in genome-wide association studies: a review on power, data quality control, genotype calling and population structure. *Current Opinion in Lipidology*. **19** (2), 133–143 (2008).
- 51 Privé, F., Aschard, H., Ziyatdinov, A., Blum, M. G. B. Efficient analysis of large-scale genome-wide data with two R packages: bigstatsr and bigsnpr. *Bioinformatics*. **34** (16), 2781–2787 (2018).
- 52 Alseekh, S. et al. Domestication of crop metabolomes: desired and unintended consequences. *Trends in Plant Science*. **26** (6), 650–661 (2021).
- 53 Yano, K. et al. GWAS with principal component analysis identifies a gene comprehensively controlling rice architecture. *Proceedings of the National Academy of Sciences of the United States of America*. **116** (42), 21262 (2019).
- 55 Wu, S. et al. Mapping the Arabidopsis metabolic landscape by untargeted metabolomics at different environmental conditions. *Molecular Plant*. **11** (1), 118–134 (2018).
- 56 Ye, J. et al. An InDel in the promoter of *Al-ACTIVATED MALATE TRANSPORTER9* selected during tomato domestication determines fruit malate contents and aluminum tolerance. *The Plant Cell*. **29** (9), 2249–2268 (2017).
- 57 Zhang, W. et al. Genome assembly of wild tea tree *DASZ* reveals pedigree and selection history of tea varieties. *Nature Communications*. **11** (1), 3719 (2020).
- 58 Tohge, T., Fernie, A. R. Annotation of plant gene function via combined genomics, metabolomics and informatics. *Journal of Visualized Experiments: JoVE*. (64), e3487 (2012).







[Click here to access/download;Figure;figure_4.png](#)



UHPLC-MS settings for lipids	
Time [min]	Eluent A to B [%]*
0 - 1.00	45% A
1.00 - 4.00	lg 45% - 25% A
4.00 - 12.00	lg 25% - 11% A
12.00 - 15.00	lg 11% - 0% A
15.00 - 19.50	cw 0% A
19.50-19.51	0% - 45% A
19.51-24.00	eq 45%
UHPLC-MS/MS settings for polar and s	
Time [min]	Eluent A and B [%]*
0 - 1.00	99% A
1.00 - 11.00	lg 99% -60% A
11.00 - 13.00	lg 60% - 30% A
13.00 - 15.00	lg 30% - 1% A
15.00 - 16.00	cw 1% A
16.00 - 17.00	lg 1% - 99% A
17.00 - 20.00	eq 99% A
GC-MS settings for derivatized metabo	
Time [min]	Temperature [°C]
0 - 2.00	85
2.00 - 18.66	lg 80 - 330
18.66 - 24.66	cw 330
24.66	rapid cooling

Information
Eluent A: 1% 1M NH ₄ -Acetate, 0.1% acetic acid in water (UHPLC grade) Eluent B: 1% 1M NH ₄ -Acetate, 0.1% acetic acid in acetonitrile/2-propanol 7:3 (UHPLC grade) Flow rate: 400 µL/min Injection volume: 2 µL
Semi-polar metabolites
Information
Eluent A: 0.1% formic acid in water (UHPLC grade) Eluent B: 0.1% formic acid in acetonitrile (UHPLC grade) Flow rate: 400 µL/min Injection volume: 3 µL
Non-polar metabolites
Information
Carrier gas: Helium Flow rate: 2 mL/min Temperature gradient: 15 °C/min Injection volume: 1 µL



Click here to access/download
Table of Materials
Table of material (1).xlsx

Dear editor,

Please find attached our revised manuscript "Large-scale Multi-omics Genome-Wide Association Studies (Mo-GWAS): Guidelines for sample preparation and normalization"

By: Mustafa Bulut, Alisdair R. Fernie and Saleh Alseekh

We have responded to the editorial comments and revised our manuscript accordingly. We hope this revised version is deemed suitable for publication as a protocol article in JoVE.

Yours sincerely,

Saleh Alseekh and all other authors

Your MS Excel document "Supplementary Table 1 .xlsx" cannot be opened and processed. Please see the common list of problems, and suggested resolutions below.

External Data

If you are submitting an Excel file, please make sure that your document does not have links to external data. If it does, break the links, save the document and resend. To break the links please do the following.

-On the Edit menu, click Links.

-In the Source list, click the link you want to break.

-To select multiple linked objects, hold down CTRL and click each linked object. To select all links, press CTRL+A.

-Click Break Link.

Embedded Macros

Your submission should not contain macros. If they do, an alert box may appear when you open your document (this alert box prevents EM from automatically converting your Excel document into the PDF that Editors and Reviewers will use). You must adjust your Excel document to remove these macros.

Excel 2002/Excel XP files

At the present time, EM supports Excel files in Excel 2000 and earlier formats. If you are using a more recent version of MS Excel, try saving your Excel document in a format compatible with Excel 2000, and resubmit to EM.

Other Problems

If you are able to get your Excel document to open with no alert boxes appearing, and you have submitted it in Excel 2000 (or earlier) format, and you still see an error indication in your PDF file (where your Excel document should be appearing). please contact the journal via the 'Contact Us' button on the Navigation Bar.'

You will need to reformat your Excel document, and then re-submit it.