

# Journal of Visualized Experiments

## Quantification of interbacterial competition using single-cell fluorescence imaging --Manuscript Draft--

Article Type:	Methods Article - JoVE Produced Video
Manuscript Number:	JoVE62851R1
Full Title:	Quantification of interbacterial competition using single-cell fluorescence imaging
Corresponding Author:	Stephanie N Smith The University of North Carolina at Chapel Hill Chapel Hill, NC UNITED STATES
Corresponding Author's Institution:	The University of North Carolina at Chapel Hill
Corresponding Author E-Mail:	Stesmith@live.unc.edu
Order of Authors:	Stephanie N Smith, M.S. Alecia Septer
Additional Information:	
Question	Response
Please specify the section of the submitted manuscript.	Biology
Please indicate whether this article will be Standard Access or Open Access.	Open Access (\$3900)
Please indicate the <b>city, state/province, and country</b> where this article will be <b>filmed</b> . Please do not use abbreviations.	Chapel Hill, North Carolina, USA
Please confirm that you have read and agree to the terms and conditions of the author license agreement that applies below:	I agree to the <a href="#">Author License Agreement</a>
Please provide any comments to the journal here.	
Please confirm that you have read and agree to the terms and conditions of the video release that applies below:	I agree to the <a href="#">Video Release</a>

**TITLE:**

Quantification of Interbacterial Competition using Single-Cell Fluorescence Imaging

**AUTHORS AND AFFILIATIONS:**

Stephanie Smith<sup>1</sup>, Alecia N. Septer<sup>1\*</sup>

<sup>1</sup>Department of Earth, Marine, and Environmental Sciences, University of North Carolina, Chapel Hill, NC 27599

\*For correspondence: 3202 Venable Hall, Chapel Hill, NC 27599

Email:

Stephanie Smith ([stesmith@live.unc.edu](mailto:stesmith@live.unc.edu))

Alecia N. Septer ([asepter@email.unc.edu](mailto:asepter@email.unc.edu))

Email Address of the corresponding author:

Alecia N. Septer ([asepter@email.unc.edu](mailto:asepter@email.unc.edu))

**KEYWORDS:**

fluorescence microscopy, single-cell, bacterial competition, *Allivibrio fischeri*, interbacterial killing, contact-dependent, live-cell imaging, quantitative microscopy, image analysis.

**SUMMARY:**

This manuscript describes a method for using single-cell fluorescence microscopy to visualize and quantify bacterial competition in coculture.

**ABSTRACT:**

Interbacterial competition can directly impact the structure and function of microbiomes. This work describes a fluorescence microscopy approach that can be used to visualize and quantify competitive interactions between different bacterial strains at the single-cell level. The protocol described here provides methods for advanced approaches in slide preparation on both upright and inverted epifluorescence microscopes, live-cell and time-lapse imaging techniques, and quantitative image analysis using the open-source software FIJI. The approach in this manuscript outlines the quantification of competitive interactions between symbiotic *Vibrio fischeri* populations by measuring the change in area over time for two coincubated strains that are expressing different fluorescent proteins from stable plasmids. Alternative methods are described for optimizing this protocol in bacterial model systems that require different growth conditions. Although the assay described here uses conditions optimized for *V. fischeri*, this approach is highly reproducible and can easily be adapted to study competition among culturable isolates from diverse microbiomes.

**INTRODUCTION:**

This article outlines a method for quantifying bacterial competition at the single-cell level using fluorescence microscopy. The structure and function of microbial communities is often shaped by competitive interactions among microbes, and in many cases characterizing these interactions

requires observing different bacterial strains in coinubation<sup>1-8</sup>. Traditionally, bacterial competition is quantified at the population level by counting colony forming units (CFUs) of inhibitor and target strains before and after a coinubation period<sup>2,9</sup>. Mechanisms for microbial competition are broadly distributed among bacteria and may rely on either diffusion or cell-cell contact to inhibit target cells<sup>10-19</sup>.

Although bacterial strains are often observed in coinubation at the population level, this manuscript outlines an assay for single-cell quantification of bacterial competition. Further, this work includes suggestions for adapting the protocol for the use with other bacterial species, including those that are unable to be genetically manipulated to express different fluorescent proteins. While the specific techniques in this article are used to study contact-dependent intraspecific competition between strains of the symbiotic bacterium *Vibrio fischeri*<sup>20-22</sup>, they can be adapted for competition between many organisms. This article provides instructions for slide setup on both upright and inverted microscopes, and all analysis is described using the open-source software FIJI<sup>23</sup> so that the method can be used by researchers with access to different imaging setups and analysis programs. Given the importance of studying microbial competition at both the population and single-cell level, this method will be a valuable resource for researchers to quantify competitive interactions, particularly those that do not have access to proprietary analysis software.

## PROTOCOL:

### 1 Optimization of bacterial strains

1.1 Choose two bacterial strains for single-cell bacterial competition assays. Here, two strains of *V. fischeri* are used: a target strain (ES114<sup>24</sup>) and an inhibitor strain (MJ11<sup>25</sup>) that is known to kill the target strain using the type VI secretion system on chromosome II (T6SS2)<sup>1</sup>, which is a contact-dependent killing mechanism.

1.2 Determine the appropriate controls for the experiment. In this example, the appropriate control is to incubate both the wild-type and the T6SS mutant inhibitor strains with the target strain to quantify the effect of T6SS-mediated killing.

NOTE: Additional controls can include a target strain that expresses the necessary immunity gene(s) to prevent T6SS-dependent killing or an inhibitor mutant strain expressing wild-type copies of the mutated genes *in trans* to restore T6SS activity<sup>1</sup>.

1.3 When possible, transform strains with stable plasmids encoding genes for different fluorescent proteins (e.g., GFP or RFP) to visually distinguish strain types on the microscope. Here, the inhibitor strain is tagged with a GFP-encoding plasmid (pVSV102), and the target strain is tagged with a dsRed-encoding plasmid (pVSV208)<sup>26</sup>.

NOTE: If it is not possible to use stable plasmids, fluorescent tags can be introduced onto the bacterial chromosome for visualization<sup>27,28</sup>.

1.4 During the initial optimization period, image clonal cultures of the tagged strains under each of the fluorescent filters that will be used during the experiment to ensure that cells are only visible in the intended channel. For example, ensure that a GFP-tagged strain is only visible in the FITC channel.

## 2 Agarose pad preparation

2.1 Prepare agarose pad solution by dissolving 2% low-melt agarose (w/v) into mPBS. Heat the solution briefly in the microwave and vortex until the agarose is completely dissolved. Keep this solution warm by placing it in a 55 °C water bath until ready to use. See the **Discussion** section for more information about preparing agarose pads.

NOTE: Here, mPBS was prepared by adding 20 g/L NaCl to standard 1x PBS.

2.2 Wrap a piece of lab tape around a glass slide five times. Repeat this process a second time on the same slide so that the distance between the two pieces of tape is slightly smaller than the width of a coverslip (**Figure 1A**). For example, if using 25 mm<sup>2</sup> coverslips, the pieces of tape should be spaced approximately 20 mm apart.

NOTE: While the number of times the tape is wrapped around the slide can be modified to adjust the thickness of the agarose pad, it is important that the layers of tape are the same height on both sides of the slide so that the agarose pad remains flat.

2.3 Pipette warm agarose solution between the two pieces of tape and immediately top with a coverslip so that it rests on the pieces of tape. This will ensure that the surface of the agarose pad remains flat. The volume of agarose solution pipetted in this step should be enough that the coverslip makes contact with the liquid and pushes out any bubbles in the agarose solution. For this particular setup, 200 µL of warm agarose is sufficient.

2.4 Let the agarose pad solidify at room temperature for at least 1 h prior to the coinubation assay. Step 2.2 will produce an agarose pad of approximately 20 mm<sup>2</sup>.

2.5 Cut this agarose pad with a razor blade into four, 5 mm<sup>2</sup> pads to be used for imaging.

NOTE: Agarose pads can be made up to one week prior to the experiment and stored at 4°C in an empty, sterile Petri plate sealed with parafilm to prevent drying.

## 3 Prepare strains for co-incubation

3.1 Streak out each strain to be used in the coinubation assay from -80 °C stocks onto LBS agar plates supplemented with the appropriate antibiotics and incubate overnight at 24 °C. For this example, three strains are used: the wild-type inhibitor strain, the type VI secretion system mutant, and the target strain.

3.2 The next day, start overnight cultures in biological duplicate by picking two colonies from each strain and resuspending them in LBS medium supplemented with the appropriate antibiotics and incubate overnight at 24°C with shaking at 200 rpm.

3.3 On the following morning, subculture each biological replicate 1:1000 into fresh LBS medium without antibiotics and incubate at 24 °C with shaking for 4-5 h or until cells reach an OD<sub>600</sub> of ~1.5.

NOTE: The timing of steps 3.1, 3.2, and 3.3 may need to be optimized for different bacterial species as their growth rate may vary substantially. For this assay, cells were aimed to be in mid-log phase at the start of the coincubation assay.

#### 4 Coincubate bacterial strains

4.1 Starting with mid-log cultures from step 3.3, measure and record the optical density at 600 nm (OD<sub>600</sub>) for all samples.

4.2 Normalize each sample to an OD<sub>600</sub> = 1.0, which corresponds to approximately 10<sup>9</sup> CFU/mL for *V. fischeri*, by diluting the culture with LBS medium.

4.3 Mix the two competing strains together at a 1:1 ratio based on volume by adding 30 µL of each normalized strain to a labeled 1.5 mL tube. Vortex the mixed-strain culture for 1-2 s.

NOTE: In some cases, it may be appropriate to mix cocultures in different ratios. For example, when one strain grows much faster than the other, it may be necessary to start the slower growing strain at a numerical advantage in order to observe the competition. Optimization may also be required if OD<sub>600</sub> does not correspond to similar CFU/mL for both strains.

4.4 Repeat step 4.3 for each biological replicate and treatment. In the example shown here, this will result in a total of four mixed-strain tubes: two biological replicates with the wild-type inhibitor strain mixed with the target strain and two biological replicates with the type VI secretion system mutant strain mixed with the target strain.

4.5 To ensure competing cells are sufficiently dense for contact-dependent killing in the coincubation on the agar pad, concentrate each mixed culture 3-fold by centrifuging the mixed culture in a standard 1.5 mL centrifuge tube for 1 min at 21,130 x *g*, discarding the supernatant, and resuspending each pellet in 20 µL LBS medium. Repeat for each sample.

NOTE: Some bacterial cells are sensitive to damage by centrifugation at high rcf; in such cases the mixed culture can be centrifuged for 3 min at 4600 x *g*<sup>29</sup>. Additionally, when quantifying contact-dependent competition, it is important to ensure sufficient cell density on the slide to observe killing. In this article, “crowded” treatments, where killing is observed, had approximately 10 cells/20 µm<sup>2</sup>; see the **Discussion** section for more information.

## 5 Slide setup

5.1 When using an upright microscope, place a  $\sim 5 \text{ mm}^2$  agarose pad onto a standard 1 mm glass slide. Spot 2  $\mu\text{L}$  of a mixed culture onto the agarose pad and place a #1.5 coverslip ( $25 \text{ mm}^2$ ) over the spot. See **Figure 1B** for an example.

5.2 When using an inverted microscope, spot 2  $\mu\text{L}$  of a mixed culture onto the #1.5 coverslip bottom of a 35 mm Petri dish and place a  $\sim 5 \text{ mm}^2$  agarose pad over the coincubation spot. Place a 12 mm circular glass coverslip over the agarose pad. See **Figure 1C** for an example.

5.3 Repeat step 5.1 or 5.2, depending on the microscope setup used, for the remaining three mixed cultures, resulting in four slides or dishes to be imaged.

5.4 Allow slides to sit on the benchtop for approximately 5 min before proceeding to step 6. This allows cells to settle on the agar pad and eliminates movement during the imaging process.

## 6 Fluorescence microscopy

6.1 Begin by focusing on cells using white light (phase contrast or DIC) to minimize the effects of photo-bleaching. Based on the average size of a single bacterial cell, use a 60x or 100x oil objective.

6.2 Adjust the exposure time and acquisition settings for each channel so that cells are visible in the appropriate channel with minimal background detection.

NOTE: It is appropriate to use different exposure times for different channels, but the same exposure time should be used across all biological replicates and treatments for a given channel.

6.3 For each sample, select at least five fields of view (FOV) and acquire images in each appropriate channel using the acquisition settings from step 6.2 (See examples in **Figure 2**). Save the XY points from each FOV so that the same FOV can be imaged during the final time point. Imaging the same FOV at each time point is necessary to determine the proportion of area occupied by target or inhibitor cells during the analysis steps.

NOTE: In this example, the fluorescence of GFP is detected using a filter with an excitation wavelength of 467 – 498 nm and an emission filter of 513 – 556 nm and is false-colored green. Fluorescence of dsRed is detected using a filter with an excitation wavelength of 542 – 582 nm and an emission filter of 603 – 678 nm and is false-colored magenta.

6.4 After 2 h, repeat step 6.3 for each sample using the previously saved XY points (**Fig 2**).

NOTE: The timing of subsequent images may need to be optimized for organisms with different growth rates or competitive mechanisms.

## 7 Image analysis in FIJI

7.1 Download and install the FIJI image processing software using the instructions found here: <https://imagej.net/Fiji/Downloads>

### 7.2 Open FIJI and import image files for analysis.

NOTE: In most cases .TIFF files will be used for image analysis, although some image acquisition software will export using proprietary file types. FIJI can recognize most proprietary file types and images can be imported and analyzed as follows.

7.3 For each image acquired in steps 6.3 and 6.4, convert the image to grayscale, separate the channels, and begin by thresholding (Ctrl + Shift + T) and creating a binary mask of the preprocessed image (Figure 3A,B).

NOTE: Here, the default thresholding settings in FIJI are used. In some cases, it may be necessary to change those settings, in which case the same settings should be used for all images in that experiment.

7.4 Set scale on the image (Analyze | Set Scale) using the appropriate values for the microscopy setup<sup>23</sup>.

7.5 Set measurements (Analyze | Set Measurements) and select Area.

NOTE: Other measurements can be added if they are appropriate for the experiment. Only the object Area measurement is required for the example analysis shown here.

7.6 Analyze particles (Analyze | Analyze Particles) using the default settings (Figure 3C). If there are debris in the sample, it may be necessary to adjust the size or circularity to filter out non-cell particles. Select Show | Outlines so that the output of this analysis will include a numbered outline of all particles analyzed (Figure 3D).

NOTE: Comparing the outline in Figure 3D to the initial image is especially important in the optimization step to ensure that (1) all cells are being analyzed, and (2) that any debris is excluded from the analysis.

7.7 Export the measurements from step 7.4 (Figure 3E) into a spreadsheet software for further analysis and graphing.

7.8 Repeat steps 7.1 – 7.5 for all channels and images acquired during the experiment.

## 8 Calculating the percent change in target area over time

8.1 For each field of view analyzed in section 7, ensure that the exported file contain an individual area measurement for each particle that was analyzed. Beginning with the target strain's fluorescence channel, calculate the sum particle area for each individual field of view. For two biological replicates with five FOV each, this should result in ten sum areas per treatment at each time point.

8.2 Calculate the percent change in the target area over time for each FOV using the following equation:  $\left(\frac{(\text{final sum area})}{(\text{initial sum area})} * 100\right)$

8.3 Repeat this calculation for all treatments and graph the percent change in the target area (result of the equation from step 8.2) for each treatment (**Figure 4A**).

8.4 Determine whether there is a net increase in the target population (indicating growth), a net decrease in the target population (indicating death), or no change (indicating no growth or death) for each treatment.

NOTE: Percent change in the target area with values greater than 100 indicate net target growth, and values lower than 100 indicate net target death. Percent change values that remain at 100 indicate that the target population did not grow but also did not lyse. See discussion for suggested follow-up experiments.

## 9 Calculating the percent change in inhibitor area over time

9.1 Repeat steps 8.1 to 8.3, this time using the measurements collected from the inhibitor strain's fluorescence channel in section 7 (**Figure 4B**).

9.2 Determine whether there was a net increase in inhibitor population (growth); a net decrease in inhibitor population (death), or no change for each treatment. Values greater than 100 indicate net inhibitor growth, and values lower than 100 indicate net inhibitor death.

## REPRESENTATIVE RESULTS:

To visualize and quantify competitive interactions between bacteria at the single-cell level, a protocol was developed and optimized for *V. fischeri* by modifying our well-established CFU-based assay<sup>1,2</sup>. This method utilizes GFP- and dsRed-encoding stable plasmids to visually distinguish different strains of *V. fischeri*. The competitive outcome of these interactions can be quantified by analyzing the images acquired from this assay using the open-source software FIJI. As an example, the following experiment was performed using *V. fischeri* isolates. An inhibitor strain harbored a plasmid that encodes GFP, and a target strain harbored a plasmid that encodes dsRed. Given that the T6SS2 encoded by the inhibitor is a contact-dependent killing mechanism, treatments were included where cells were either crowded (high cell-cell contact) or disperse (low cell-cell contact) on a slide to highlight the impact of experimental setup on the final results of this assay. In the sample data, competing strains were mixed at a 1:1 ratio and incubated on an agarose pad for 2 h, and both initial and final (2 h) images were taken. As a control, a T6SS2



mutant strain was also coincubated with the target strain in both crowded and disperse conditions. Cultures of each strain were prepared and coincubated as described above and slides were prepared as shown in **Figure 1**.

**Figure 2** shows representative fluorescence microscopy images of each experimental treatment with the same field of view imaged at an initial and final time point. For each treatment, either a wild-type inhibitor or T6SS mutant strain harboring a GFP-encoding plasmid was mixed at a 1:1 ratio with the target strain harboring a dsRed-encoding plasmid. During a 2 h coincubation period with this experimental setup, growing *V. fischeri* cells may go through 1-2 divisions (**Figure 2; gray arrows**). In **Figure 2A**, cell-cell contact was forced between the target and inhibitor by concentrating the mixed culture before spotting onto the slide. Multiple target cells are observed to become rounded and/or disappear over the course of 2 h, consistent with target cells being eliminated by the inhibitor (**Figure 2; white arrows**). See the **Discussion** section for more information on interpreting rounding or lysing target cells. In **Figure 2B**, the same coincubation was spotted onto a slide, this time without concentrating the mixed culture so that the cells remained disperse and there was minimal contact between strains on the slide. Here, no target cells are observed to disappear or round, suggesting that the target strain was not inhibited in this treatment. **Figure 2C** and **Figure 2D** show the same crowded and disperse treatments described above, this time using a T6SS mutant as the inhibitor strain. Target cells were not observed to disappear or round when coincubated with a T6SS mutant in either crowded or disperse conditions, again suggesting that the target was not inhibited in either treatment.

**Figure 3** shows the FIJI analysis workflow used to quantify competition in this protocol. A representative image from the target channel was selected (**Figure 3A**) and a binary mask was created using the default threshold settings in FIJI (**Figure 3B**). The image scale was set appropriately for this microscopy setup. Particles were analyzed using the size parameter = 0 – infinity, circularity parameter = 0.00 – 1.00, and **Show Outlines** was selected (**Figure 3C**). The results of this particle analysis are shown as both a numbered outline of each particle (**Figure 3D**), and as a table with columns for the particle number, file name (label), and particle area in  $\mu\text{m}^2$  (area) (**Figure 3E**).

In **Figure 4**, data obtained from **Figure 3E** is graphed and analyzed. In **Figure 4A**, the percent change in target area between the initial and final timepoint is presented for each treatment according to step 8.2. If the percent change in the target area is greater than 100, this represents net increase in target (i.e., growth) and is observed in conditions where the target population is not significantly inhibited. However, if the percent change in the target area is lower than 100, this result indicates a net decrease in the target (i.e., death) and is observed in conditions where the target population is significantly inhibited. When the target was coincubated with a wild-type inhibitor in crowded conditions, the data show a net decrease in the target area. By contrast, when the target was coincubated with either a wild-type inhibitor in disperse conditions or a T6SS mutant in crowded or disperse conditions, the data show a net increase in the target area. The percent change in the target area when the target was coincubated with a wild-type inhibitor in crowded conditions was below 100 and significantly lower than all other treatments according to a one-way ANOVA followed by a Tukey's multiple comparisons test across all treatments ( $p <$

0.0001). These data indicate that target cell death is dependent on a functional T6SS in the inhibitor and underscores the importance of an experimental setup that allows sufficient cell-cell contact, in order to detect cell death from a contact-dependent killing mechanism.

**Figure 4B** presents the change in inhibitor area between the initial and final timepoint for each treatment. In this example, net growth of the inhibitor strain was observed across all treatments. However, the percent change in inhibitor area was significantly higher when a wild-type inhibitor was coincubated with the target in crowded conditions compared to all other treatments according to a one-way ANOVA followed by a Tukey's multiple comparisons test across all treatments ( $p < 0.0001$ ). Initially, we considered that the net increase in inhibitor area may be driven by the increase in available space to grow into as target cells are eliminated. However, this same increase in inhibitor growth was not observed in disperse treatments, where inhibitor cells had room to grow from the beginning of the coincubation. Alternatively, this result could suggest that nutrients released from lysing target cells allow for a greater increase in the inhibitor population. Taken together, these results suggest that the inhibitor strain eliminates the target in a T6SS-dependent manner only when high cell-cell contact is forced by crowding cells on the slide.

#### FIGURE AND TABLE LEGENDS:

**Figure 1: Agarose pad preparation and slide setup for coincubation assays.** (A) Setup for making 2% agarose pads. Five layers of lab tape (green) are wrapped around a cover slip at two points approximately 20 mm apart. Next, warm 2% agarose in mPBS (yellow) is pipetted between the pieces of tape and immediately covered with a 25 mm<sup>2</sup> cover slip and allowed to solidify for at least 1 h at room temperature. Use a razor blade to cut the agarose pad into ~5 mm<sup>2</sup> pieces and use tweezers to transfer the pad to a new slide for imaging. (B) When imaging on an upright microscope, place the 5 mm<sup>2</sup> agarose pad directly onto the slide, followed the mixed culture (blue) and a 12 mm circular #1.5 cover slip. (C) When imaging on an inverted microscope, spot the mixed culture directly onto the #1.5 glass cover slip bottom of a 35 mm Petri dish, and place an agarose pad on top of the culture followed by a second 12 mm circular cover slip to flatten the agarose pad.

**Figure 2: Time lapse images of coincubation spots in either crowded or disperse conditions.** (A) Representative images at initial and final time points where a mixed culture of target and wild-type inhibitor was concentrated 3x prior to spotting on the slide to force cell-cell contact between strains. White arrows in TRITC channel indicate examples of target cells that round or lyse throughout the course of the experiment. (B) Representative images where a mixed culture of target and wild-type inhibitor was spotted without concentrating so that cells are disperse and there is minimal cell-cell contact between strains. Gray arrows in FITC and TRITC channels indicate examples of cell division throughout the course of the experiment. (C) Representative images where mixed culture of target and T6SS<sup>-</sup> mutant was concentrated 3x prior to spotting on the slide to force cell-cell contact between strains. (D) Representative images where mixed culture of target and T6SS<sup>-</sup> mutant was spotted without concentrating so that cells are disperse and there is minimal cell-cell contact between strains. Scale bars = 5 μm and are consistent across all images; TRITC channel is false-colored magenta, FITC channel is false-colored green.

Deconvolution was performed on all images; background was subtracted, and brightness/contrast adjusted uniformly across all images.

**Figure 3: FIJI analysis workflow.** (A) Representative image for analysis. This workflow is repeated for both channels across all fields of view and samples. Scale bars = 5  $\mu\text{m}$  and are consistent across all images; TRITC channel is false-colored magenta, FITC channel is false-colored green. (B) Binary mask created by thresholding the image using the default settings in FIJI. (C) Example of settings for particle analysis used in this manuscript. Size range = 0-infinity  $\mu\text{m}^2$ ; circularity = 0.00 – 1.00; show = outlines. (D) Particle outline created as an output of particle analysis in (C). The particle outline in (D) should be compared to the original image (A) to ensure that all cells were captured in the particle analysis. (E) Results table created as an output from particle analysis in (C). Object number (column 1) corresponds to individual particles (one or more cells) outlined and labeled in red in panel (D). Label = file name of analyzed image; Area = total particle area in  $\mu\text{m}^2$ .

**Figure 4: Sample data for assessing whether target strain is inhibited.** The percent change in area between initial and final time points for the target strain (A) and inhibitor strain (B), at different initial cell densities. Slide density indicates either a starting cell density that is crowded (high cell contact between strains), or more disperse (low cell contact between strains) as described in Figure 2. Inhibitor genotype indicates that either a wild-type or the T6SS mutant (T6SS<sup>-</sup>) strain was coinoculated with the target strain. Asterisks indicate a significant difference in % change comparing all treatments (one-way ANOVA followed by a Tukey's multiple comparisons test comparing all treatments; ( $p < 0.0001$ )). Dashed line indicates no net change in strain area between the initial and final timepoint; a % change > 100 indicates net increase (i.e., growth) and % change < 100 indicates net decrease (i.e., cell death).

## DISCUSSION:

The protocol described above provides a powerful tool for quantifying and characterizing interbacterial competition at the single-cell level. This assay, which was developed by modifying our CFU-based competition assay on agar plates<sup>1,2</sup>, allowed for the visualization of single-cell competition among *V. fischeri* isolates and suggestions are provided for optimizing the method for a wide range of systems and microscopy setups. Although the method described here was optimized for the light-organ symbiont *V. fischeri*, it can be easily modified to accommodate many diverse, culturable microbes. It is important to note that competitive mechanisms can be regulated by any number of environmental variables, including temperature, salinity, and viscosity<sup>30-34</sup>. Previous work has confirmed that *V. fischeri* competes using a contact-dependent Type VI Secretion System that is active on surfaces<sup>30</sup>, making the conditions described in this assay suitable for studying competition between the example strains. It is also important to consider the initial density of cells on the slide when quantifying bacterial competition. Given that contact between target and inhibitor cells is often required for killing to occur, the mixed culture should be concentrated such that cell-cell contact is maximized and cells remain in a single plane on the slide. Cell cultures should be grown to a similar optical density (mid-log phase) and then concentrated to force contact rather than simply growing cultures to a higher optical density due to the physiological changes of cells in different growth phases. In other systems,

culture conditions and the experimental setup may need to be modified to ensure that the competitive mechanism is active and can be detected in the coincubation condition.

The agarose pads used in this assay provide several benefits: they provide stabilization so that cells do not move around freely, and they prevent the culture from drying out over the course of the experiment. Additionally, if chemical inducers, such as isopropyl- $\beta$ -D-thiogalactoside (IPTG), are required for the experiment, they can easily be added to the agarose solution. However, it is important to note that the agarose preparation will likely need to be adjusted for different systems. In the example described above, the agarose pad was prepared by dissolving 2% agarose (w/v) into 20 psu mPBS, which is the standard salinity used in *V. fischeri* growth medium. Furthermore, in some cases a carbon source may need to be added to the agarose pad in order for cells to grow and compete over longer experiments. In such a case, the mPBS in agarose pads can be replaced with any growth medium, although the nutrients in growth medium may come with the tradeoff of additional background fluorescence.

Without proprietary image analysis software, it can be very difficult to get individual cell counts when cell-cell contact is high, which as we show here is required to observe contact-dependent killing. This assay was designed to provide an alternative method for quantification that does not rely on individual cell counts. Instead, the total cell area for each fluorescence channel is used to quantify the extent of killing between coincubated strains. Because this method relies on area rather than individual cell counts, default thresholding settings are typically sufficient for outlining the total cell area. The accuracy of thresholding can be verified by dividing the total object area for a representative field of view by the average cell size for the model organism and comparing this estimated cell number to a manual cell count for the same image.

In coincubations between one inhibitor and one target (non-killer) strain, net growth of the inhibitor is predicted. As seen in **Figure 4**, inhibitor growth may be significantly higher in treatments where killing is observed, compared to treatments where killing is not observed, perhaps because nutrients released by lysing target cells allow the inhibitor strain to grow more quickly. In the example shown here, net target death is observed because T6SS-mediated competition results in target cell lysis where the target is physically eliminated. However, it is important to note that not all competitive mechanisms result in the physical elimination of target cells. If a target is incapacitated by a toxin that causes growth inhibition, the protocol outlined here may result in the visible target population remaining stable over time as target cells no longer grow but also do not lyse. In such a case, it would be appropriate to compare the results of this assay with follow up tests for target cell viability, such as plating for colony forming units (CFUs) or by performing live-dead assays by staining with propidium iodide or SYTOX green<sup>35,36</sup>.

Compared to coincubation assays that rely on CFU counts, this assay makes it possible to observe and quantify the spatial structure of competition between strains and track changes in target cell morphology over time. For example, inhibitor cells that kill using a T6SS are known to encode LysM-domain protein that degrade target cell wall, resulting in initial cell rounding and then lysis<sup>13</sup>, which we observed in the example shown in **Figure 2A**. Further, this protocol can be used to track competition at high resolution over very short time scales. In the example shown here,

a significant decrease in the target area is observed after only two hours when cells are crowded and cell-cell contact is forced between strains (**Figure 4**). The image analysis described here could also be performed using confocal microscopy, which would make it possible to study bacterial competition *in vivo* or in complex biofilms, without disrupting the spatial distribution of coincubated strains.

In summary, the assay described here aims to provide an accessible and easily modified approach for visualizing and quantifying bacterial competition at the single-cell level using fluorescence microscopy. This method can be applied to diverse bacterial isolates and can be used to visualize bacterial competition even in complex environments such as within a host or biofilm matrix.

#### **ACKNOWLEDGMENTS:**

A.N.S was supported by NIGMS grant R35 GM137886 and S.N.S was supported by the National Defense Science and Engineering Graduate Fellowship Program.

#### **DISCLOSURES:**

The authors have nothing to disclose.

#### **REFERENCES:**

- 1 Speare, L. et al. Bacterial symbionts use a type VI secretion system to eliminate competitors in their natural host. *Proceedings of the National Academy of Sciences*. **115** (36), E8528-E8537 (2018).
- 2 Speare, L., Septer, A. N. Coincubation assay for quantifying competitive interactions between *Vibrio fischeri* isolates. *Journal of Visualized Experiments*. (149) e59759 (2019).
- 3 Frost, I. et al. Cooperation, competition and antibiotic resistance in bacterial colonies. *The ISME journal*. **12** (6), 1582-1593 (2018).
- 4 Stubbendieck, R. M., Vargas-Bautista, C., Straight, P. D. Bacterial communities: interactions to scale. *Frontiers in Microbiology*. **7**, 1234 (2016).
- 5 Souza, D. P. et al. Bacterial killing via a type IV secretion system. *Nature Communications*. **6** (1), 1-9 (2015).
- 6 Anderson, M. C., Vonaesch, P., Saffarian, A., Marteyn, B. S., Sansonetti, P. J. *Shigella sonnei* encodes a functional T6SS used for interbacterial competition and niche occupancy. *Cell Host and Microbe*. **21** (6), 769-776. e763 (2017).
- 7 Basler, M., Ho, B., Mekalanos, J. Tit-for-tat: Type VI secretion system counterattack during bacterial cell-cell interactions. *Cell*. **152** (4), 884-894 (2013).
- 8 Guillemette, R., Ushijima, B., Jalan, M., Häse, C. C., Azam, F. Insight into the resilience and susceptibility of marine bacteria to T6SS attack by *Vibrio cholerae* and *Vibrio coralliilyticus*. *PLoS One*. **15** (1), e0227864 (2020).
- 9 Hachani, A., Lossi, N. S., Filloux, A. A visual assay to monitor T6SS-mediated bacterial competition. *Journal of Visualized Experiments*. (73), e50103, (2013).
- 10 Hibbing, M. E., Fuqua, C., Parsek, M. R., Peterson, S. B. Bacterial competition: surviving and thriving in the microbial jungle. *Nature Reviews Microbiology*. **8** (1), 15-25 (2010).
- 11 Ruhe, Z. C., Low, D. A., Hayes, C. S. Bacterial contact-dependent growth inhibition. *Trends in Microbiology*. **21** (5), 230-237 (2013).

528 12 Wood, D. W., Pierson III, L. S. The *phzI* gene of *Pseudomonas aureofaciens* 30–84 is  
529 responsible for the production of a diffusible signal required for phenazine antibiotic production.  
530 *Gene*. **168** (1), 49-53 (1996).

531 13 Smith, W. P. et al. The evolution of the type VI secretion system as a disintegration  
532 weapon. *PLoS Biology*. **18** (5), e3000720 (2020).

533 14 Chen, L., Zou, Y., She, P., Wu, Y. Composition, function, and regulation of T6SS in  
534 *Pseudomonas aeruginosa*. *Microbiological Research*. **172**, 19-25 (2015).

535 15 Sana, T. G., Lugo, K. A., Monack, D. M. T6SS: The bacterial" fight club" in the host gut. *PLoS*  
536 *Pathogens*. **13** (6), e1006325 (2017).

537 16 Basler, M. Type VI secretion system: secretion by a contractile nanomachine.  
538 *Philosophical Transactions of the Royal Society B: Biological Sciences*. **370** (1679), 20150021  
539 (2015).

540 17 Joshi, A. et al. Rules of engagement: the type VI secretion system in *Vibrio cholerae*.  
541 *Trends in Microbiology*. **25** (4), 267-279 (2017).

542 18 Nadell, C. D., Drescher, K., Foster, K. R. Spatial structure, cooperation and competition in  
543 biofilms. *Nature Reviews Microbiology*. **14** (9), 589-600 (2016).

544 19 Stubbendieck, R. M., Straight, P. D. Multifaceted interfaces of bacterial competition.  
545 *Journal of Bacteriology*. **198** (16), 2145-2155 (2016).

546 20 Septer, A. N. The *Vibrio*-squid symbiosis as a model for studying interbacterial  
547 competition. *Msystems*. **4** (3), (2019).

548 21 Tischler, A. H., Hodge-Hanson, K. M., Visick, K. L. *Vibrio fischeri*–squid symbiosis. *eLS*. 1-9,  
549 (2019).

550 22 Mandel, M. J., Dunn, A. K. Impact and influence of the natural *Vibrio*-squid symbiosis in  
551 understanding bacterial–animal interactions. *Frontiers in Microbiology*. **7**, 1982 (2016).

552 23 Schindelin, J. et al. Fiji: an open-source platform for biological-image analysis. *Nature*  
553 *Methods*. **9** (7), 676-682 (2012).

554 24 Boettcher, K., Ruby, E. Depressed light emission by symbiotic *Vibrio fischeri* of the sepiolid  
555 squid *Euprymna scolopes*. *Journal of Bacteriology*. **172** (7), 3701-3706 (1990).

556 25 Doino, J. A., McFall-Ngai, M. J. A transient exposure to symbiosis-competent bacteria  
557 induces light organ morphogenesis in the host squid. *The Biological Bulletin*. **189** (3), 347-355  
558 (1995).

559 26 Dunn, A. K., Millikan, D. S., Adin, D. M., Bose, J. L., Stabb, E. V. New *rfp*- and *pES213*-derived  
560 tools for analyzing symbiotic *Vibrio fischeri* reveal patterns of infection and *lux* expression in situ.  
561 *Applied and Environmental Microbiology*. **72** (1), 802-810 (2006).

562 27 Lambertsen, L., Sternberg, C., Molin, S. Mini-Tn7 transposons for site-specific tagging of  
563 bacteria with fluorescent proteins. *Environmental Microbiology*. **6** (7), 726-732 (2004).

564 28 Koch, B., Jensen, L. E., Nybroe, O. A panel of Tn7-based vectors for insertion of the *gfp*  
565 marker gene or for delivery of cloned DNA into Gram-negative bacteria at a neutral chromosomal  
566 site. *Journal of Microbiological Methods*. **45** (3), 187-195 (2001).

567 29 Peterson, B. W., Sharma, P. K., Van Der Mei, H. C., Busscher, H. J. Bacterial cell surface  
568 damage due to centrifugal compaction. *Applied and Environmental Microbiology*. **78** (1), 120-125  
569 (2012).

570 30 Speare, L., Smith, S., Salvato, F., Kleiner, M., Septer, A. N. Environmental viscosity  
571 modulates interbacterial killing during habitat transition. *MBio*. **11** (1), (2020).

572 31 Salomon, D., Gonzalez, H., Updegraff, B. L., Orth, K. *Vibrio parahaemolyticus* type VI  
573 secretion system 1 is activated in marine conditions to target bacteria, and is differentially  
574 regulated from system 2. *PloS One*. **8** (4), e61086 (2013).

575 32 Sana, T. G. et al. *Salmonella Typhimurium* utilizes a T6SS-mediated antibacterial weapon  
576 to establish in the host gut. *Proceedings of the National Academy of Sciences*. **113** (34), E5044-  
577 E5051 (2016).

578 33 Bachmann, V. et al. Bile salts modulate the mucin-activated type VI secretion system of  
579 pandemic *Vibrio cholerae*. *PLoS Neglected Tropical Diseases*. **9** (8), e0004031 (2015).



580 34 Ishikawa, T. et al. Pathoadaptive conditional regulation of the type VI secretion system in  
581 *Vibrio cholerae* O1 strains. *Infection and Immunity*. **80** (2), 575-584 (2012).

582 35 Johnson, M. B., Criss, A. K. Fluorescence microscopy methods for determining the viability  
583 of bacteria in association with mammalian cells. *Journal of Visualized Experiments*. (79), e50729  
584 (2013).

585 36 Stiefel, P., Schmidt-Emrich, S., Maniura-Weber, K., Ren, Q. Critical aspects of using  
586 bacterial cell viability assays with the fluorophores SYTO9 and propidium iodide. *BMC*  
587 *Microbiology*. **15** (1), 36 (2015).

588

Figure 1



**C.** [Click here to access/download;Figure;Fig-1.pdf](#) 











Figure 2







**E.**

	Label	Area
1	T6-crowded	3.046
2	T6-crowded	1.411
3	T6-crowded	0.558
4	T6-crowded	0.739
5	T6-crowded	39.791

Figure 4



[Click here to access/download;Figure;Fig-4.pdf](#)





[Click here to access/download](#)

**Table of Materials**  
**JoVE\_Materials\_SNSrevisions.xls**



**Editorial comments:**

Changes to be made by the Author(s):

1. Please take this opportunity to thoroughly proofread the manuscript to ensure that there are no spelling or grammar issues. Please define all abbreviations at first use.

[We have proofread the manuscript and have defined all abbreviations at first use.](#)

2. Please provide an email address for each author.

[Line 9-10, email addresses are provided for each author.](#)

3. Please increase your abstract's word count to 150-300 words.

[Lines 21 – 33, the abstract's word count was increased to 153 words.](#)

4. Please revise the following lines to avoid overlap with previously published work: 102-106.

[Lines 106 – 114 have been revised to avoid overlap with previously published work.](#)

5. JoVE cannot publish manuscripts containing commercial language. This includes trademark symbols (™), registered symbols (®), and company names before an instrument or reagent. Please remove all commercial language from your manuscript and use generic terms instead. All commercial products should be sufficiently referenced in the Table of Materials and Reagents.

For example: ORCA-Fusion Digital CMOS camera; NIS-Elements software etc

[References to commercial products were moved to the Table of Materials and Reagents and removed from the main text.](#)

6. Step 4.5: What do you mean by "Concentrate each mixed culture 3x by centrifuging 1 m at max speed (21,130 x g)"?

[Lines 135 – 138, this step was reworded to clarify how the protocol should be performed.](#)

7. Please ensure that all text in the protocol section is written in the imperative tense as if telling someone how to do the technique (e.g., "Do this," "Ensure that," etc.). The actions should be described in the imperative tense in complete sentences wherever possible. Avoid usage of phrases such as "could be," "should be," and "would be" throughout the Protocol. Any text that cannot be written in the imperative tense may be added as a "Note." However,

notes should be concise and used sparingly. Please include all safety procedures and use of hoods, etc. (e.g., step 7.1)

Line 179, step 7.1 was reworded so it is written in the imperative tense.

8. Please note that your protocol will be used to generate the script for the video and must contain everything that you would like shown in the video. Please ensure you answer the "how" question, i.e., how is the step performed? Alternatively, add references to published material specifying how to perform the protocol action. There should be enough detail in each step to supplement the actions seen in the video so that viewers can easily replicate the protocol.

The protocol has been clarified so that it can be used to generate the script for the video in a way that viewers can easily replicate the protocol.

9. After including a one line space between each protocol step, highlight up to 3 pages of protocol text for inclusion in the protocol section of the video. This will clarify what needs to be filmed. (Tip: Do not highlight notes or headings of sections in which all steps won't be filmed.)

One line space was added between each protocol step, and a section of the protocol text has been highlighted to include in the protocol section of the video.

10. Please include a scale bar for all images taken with a microscope to provide context to the magnification used. Define the scale in the appropriate Figure Legend.

Figures 2 and 3 were updated so that scale bars are on every microscope image, and the scale bar is defined in each figure legend.

11. Please add limitations of the technique to the discussion.

The discussion section has been updated to reflect the limitations to this technique; also addressed in reviewer 1 responses.

---

### **Reviewers' comments:**

#### **Reviewer #1:**

The manuscript by Stephanie Smith and Alecia N. Septer titled „Quantification of interbacterial competition using single-cell fluorescence microscopy" describes a method that can be used to quantify bacterial competition using fluorescence microscopy. The

manuscript is well written, the authors clearly present the method, explain all steps necessary to perform analysis, and give easily understandable examples. However, the manuscript suffers from some oversimplification which might lead to scientific inaccuracies if performed by others. The manuscript also lacks description of limitations that the method clearly introduces. Finally, I believe that some experimental data should be included in order to support statements given by the authors. Overall, it is an interesting method article that I believe can be published after some major revisions.

Questions/suggestions.

1. Lines 43-44. The authors in the manuscript suggest that their method "highlights the importance of considering starting cell density when drawing conclusions about competition via contact-dependent mechanisms". In my opinion this statement is unnecessary as contact-dependent mechanism itself means that the contact must be present and at low densities it is kept to a minimum.

From lines 45-46 we removed the statement that this method highlights the importance of considering starting cell density when drawing conclusions about competition via contact-dependent mechanisms.

2. In the introduction I would suggest to include a short description of currently used methods for quantification of interbacterial competition.

Lines 38-40 were reworded to clarify that traditional methods of quantification of interbacterial competition primarily rely on CFU counts of inhibitor and target populations before and after a coincubation period.

3. The introduction also contains sentence (Line 49) that the method can be used to study diffusible competitive mechanisms. However, this is the only place in the article where it is mentioned. As this was not evaluated I would suggest to remove this statement or include it in the discussion.

We removed the mention of using this method to study diffusible competitive mechanisms from line 49.

4. Lines 63-65 (Step 1.2). I would suggest to include a sentence to add an additional control (whether a target strain complemented with immunity gene (a strain) or inhibitor mutant strain with reinstated T6SS activity).

Lines 68-70 we included a note (1.2.1) that suggests using these additional controls.

5. Lines 66-69 (Step 1.2.2). In the specific cases where it is not possible to use stable plasmids expressing fluorescent proteins, the authors suggest to use stains to distinguish bacteria species. Such statement should be referenced with the appropriate literature or experiments indicating no impact on the killing ability or susceptibility in the interbacterial competition. In my opinion this point is extremely important. If there is no interference of fluorescent stains to bacterial viability and/or their susceptibility to killing there is a potential for the method to be applied for the evaluation of interbacterial competition among important clinical pathogens that are resistant to multiple antibiotics as they cannot be transformed with plasmids containing selective markers. This point should be emphasized more in the article. Probably in the discussion section.

The reviewer brings up an excellent point. We performed further experiments with *V. fischeri* strains to validate our suggested modification of using dyes to distinguish different bacterial species in coinocubation. However, we were unable to distinguish between differentially-stained bacterial strains because over the course of coinocubation, both populations appear to take up the stain even when the cells were washed multiple times before coinocubation. We have therefore removed this suggestion from this current work, and hope to optimize this method more in the future.

6. Line 78 (Step 2.1). It should be noted whether a low-melting or conventional agarose was used.

In line 82, we clarify that low-melting agarose is used to prepare agarose pads.

7. Lines 129-131 (Step 4.5). Centrifuging time should be corrected into "min" instead of "m". Also, I would advise against using such a high speed centrifuging forces as it has been suggested that they might cause damage to bacterial cells (as described by Peterson BW, Sharma PK, van der Mei HC, Busscher HJ. Bacterial cell surface damage due to centrifugal compaction. Appl Environ Microbiol. 2012 Jan;78(1):120-5. doi: 10.1128/AEM.06780-11. Epub 2011 Oct 28. PMID: 22038609; PMCID: PMC3255633).

Have you checked cell viability after centrifugation?

Line 137, "m" was corrected to "min". We added a note (4.5.1) to indicate that some cells may be sensitive to damage by centrifugation at high rcf and suggest a lower rcf for 3 min in this case. In *V. fischeri*, we have obtained CFU counts before and after centrifugation at the indicated rcf and do not observe a significant difference in cell viability after centrifugation.



8. Lines 66-69 (Step 4.5.1). the targeted density of cells on the slides should be indicated.

Lines 142 - 143, we indicate that "crowded" treatments should aim for a cell density of approximately 10 cells/20  $\mu\text{m}^2$ .

9. Lines 175-177 (Step 7.3). Authors should indicate that image must be converted to grayscale.

Lines 185 - 187, we indicate that the image should be converted to 8-bit grayscale prior to separating the channels and thresholding.

10. The authors should perform and provide data as a figure from colony counting assays where agarose pads are suspended in media and plated to enumerate bacteria before and after competition experiment. This would indicate how and if the microscopy results are comparable to the conventional bacterial competition evaluation method.

Lines 230 - 232, we indicate that this assay was developed by modifying our CFU-based competition assay and provide references that go into details about how these were performed and used to characterize T6SS-dependent killing in *V. fischeri*.

11. My question about Figure 2. Why do we see similar bacterial density before and after (2h) competition experiments. Do bacteria grow and divide during this period? If not, are the conditions appropriate for the competition as killing via T6SS requires energy? I would suggest to include carbon source into agarose pad. In addition, Figure 4 B shows the net growth of the inhibitor strain by ~300%. However, from Figure 2 this increase is not obvious. How can it be?

Lines 249 - 251 and Figure 2. In text we indicate that during a 2 h coinoculation, *V. fischeri* cells may go through 1-2 divisions. In Figure 2, we added arrows to indicate cells that have divided between the initial and final timepoint. In the discussion (lines 385 – 386), we indicate that in some cases it may be required to add a carbon source to the agarose pad in order for cells to grow and compete over longer time periods. However, we have found that addition of a carbon source is not necessary for these short-term experiments as *V. fischeri* and other copiotrophic bacterial species typically must be starved in carbon free media for days before they stop dividing. Adding carbon also could mask the effects of nutrient release from lysed cells, as discussed in the below comment.

12. The argument in lines 280-281: "This result suggests that as the inhibitor eliminates target cells, more space is available for the inhibitor to grow without competition for space." This statement does not sound correct as one would expect the similar increase in cell

numbers under the disperse conditions where the space to grow is plenty. However, I do not see such effect in Figure 4B.

Lines 295 - 300, we would like to thank the reviewer for pointing this out to us. The results section has been updated to reflect this comment and we hypothesize that perhaps the increased growth of the lethal strain when killing is detected could be driven by the added nutrients that are released from lysing target cells.

13. The authors state that the method can be applied to various competition mechanisms. I do not think this is correct as this depends on the toxins used by the attacker strain. In the case of toxins that causes cell stasis and/or growth inhibition this method might give false negative results. The authors make an argument that their method can differentiate it by evaluating population size of target bacteria (lines 379-383). However, no real experimental evidence with such inhibition types are presented. This should be acknowledged in the manuscript.

Lines 402 - 410, we clarify that this method alone is not sufficient to determine cell death in the absence of target cell lysis, and provide suggestions for follow-up assays to further investigate target cell viability.

14. The limitations of the method should be included in the discussion section.

Lines 402 - 410, we address limitations of the method as described in the previous comment.

## **Reviewer #2:**

Manuscript Summary:

This manuscript described a detailed method for preparing bacterial co-cultures, imaging protocols, and image quantification.

Major Concerns:

None.

Minor Concerns: The authors did a great job going through the protocols. Here are just a few small comments.

- In the title "single-cell fluorescence microscopy" maybe a little bit confusing, can revised to "single-cell fluorescence imaging".

Line 2, the title has been updated to reflect this suggestion and the word "microscopy" was changing to "imaging".

- Line 70-72: It is quite common that many bacteria can not be transformed, and this protocol can be applied to broader audience in microbiology if the authors describe in more details about the methods for DAPI or SYBR Green stain.

As described in Reviewer #1 response: Upon further experimentation, we chose to remove our suggestion of using stains to distinguish different bacterial species.

### **Reviewer #3:**

Manuscript Summary:

The submitted manuscript describes a method for visualizing and quantifying bacterial competition at the single-cell level using fluorescence microscopy. It is based on using changes in the areas occupied by the inhibiting (killer) and target bacterial strains in the microscopic field of view over a specific time period of co-incubation (typically a few hours). The protocols are provided in sufficient detail, covering slide preparation, bacterial strain preparation, image collection and quantitative image analysis using the FIJI software. The discussion deals with the important consideration of adjusting the bacterial cell density in the co-culture so that inhibiting and target cells are in contact during the timelapse experiment, thus guaranteeing a significant number of killing events. I believe that the protocol, as presented, is a good point to begin for a laboratory setting out to explore this phenomenon. The manuscript is well written and easy to follow. Representative results are shown and the data-processing workflow is presented in an easy-to-follow manner.

Major Concerns:

I have no major concerns.

Minor Concerns:

I have no minor concerns.