

Video Article

Propagation of Human Embryonic Stem (ES) Cells

Laurence Daheron¹

¹Center for Regenerative Medicine, MGH - Massachusetts General Hospital

Correspondence to: Laurence Daheron at ldaheron@partners.org

URL: <https://www.jove.com/video/119>

DOI: [doi:10.3791/119](https://doi.org/10.3791/119)

Keywords: Cellular Biology, Issue 1, ES, embryonic stem cells, tissue culture

Date Published: 11/30/2006

Citation: Daheron, L. Propagation of Human Embryonic Stem (ES) Cells. *J. Vis. Exp.* (1), e119, doi:10.3791/119 (2006).

Abstract

Video Link

The video component of this article can be found at <https://www.jove.com/video/119/>

References

1. Kawasaki, H. et al. Induction of midbrain dopaminergic neurons from ES cells by stromal cell-derived inducing activity. *Neuron* 28, 31-40 (2000).
2. Perrier, A.L. et al. Derivation of midbrain dopamine neurons from human embryonic stem cells. *Proc Natl Acad Sci U S A* 101, 12543-12548 (2004).
3. Pruszak, J., Isacson, O. Directed differentiation of human embryonic stem cells into dopaminergic neurons. *Human Embryonic Stem Cells: The Practical Handbook* Sullivan, S., Cowan, C., Eggan, K. (eds.); John Wiley and Sons (2007).
4. Pruszak, J., Sonntag, K.C., Aung, M.H., Sanchez-Pernaute, R. and Isacson, O. Markers and methods for cell sorting of human embryonic stem cell-derived neural cell populations. *Stem Cells* 25, 2257-2268 (2007).
5. Sonntag, K.C. et al. Enhanced yield of neuroepithelial precursors and midbrain-like dopaminergic neurons from human embryonic stem cells using the bone morphogenic protein antagonist noggin. *Stem Cells* 25, 411-418 (2007).
6. Kawasaki, H. et al. Generation of dopaminergic neurons and pigmented epithelia from primate ES cells by stromal cell-derived inducing activity. *Proc Natl Acad Sci U S A* 99, 1580-1585 (2002).
7. Ko, J.Y. et al. Human embryonic stem cell-derived neural precursors as a continuous, stable, and on-demand source for human dopamine neurons. *J Neurochem* 103, 1417-1429 (2007).
8. Hong, S., Kang, U.J., Isacson, O. and Kim, K.S. Neural precursors derived from human embryonic stem cells maintain long-term proliferation without losing the potential to differentiate into all three neural lineages, including dopaminergic neurons. *J Neurochem* (2007).
9. Zhang, S.C. Neural subtype specification from embryonic stem cells. *Brain Pathol* 16, 132-142 (2006).
10. Pankratz, M.T. et al. Directed neural differentiation of human embryonic stem cells via an obligated primitive anterior stage. *Stem Cells* 25, 1511-1520 (2007).
11. Lazzari, G. et al. Direct derivation of neural rosettes from cloned bovine blastocysts: a model of early neurulation events and neural crest specification in vitro. *Stem Cells* 24, 2514-2521 (2006).
12. Pomp, O., Brokhman, I., Ben-Dor, I., Reubinoff, B. and Goldstein, R.S. Generation of peripheral sensory and sympathetic neurons and neural crest cells from human embryonic stem cells. *Stem Cells* 23, 923-930 (2005).