

Video Article

Separation of *Plasmodium falciparum* Late Stage-infected Erythrocytes by Magnetic Means

Lorena Michelle Coronado^{1,2}, Nicole Michelle Tayler^{1,2}, Ricardo Correa^{1,2}, Rita Marissa Giovani³, Carmenza Spadafora¹

¹Centro de Biología Celular y Molecular de Enfermedades, Instituto de Investigaciones Científicas y Servicios de Alta Tecnología (INDICASAT AIP)

²Acharya Nagarjuna University

³Departamento de Medios y Creativo, Instituto de Investigaciones Científicas y Servicios de Alta Tecnología (INDICASAT AIP)

Correspondence to: Carmenza Spadafora at cspadafora@indicasat.org.pa

URL: <https://www.jove.com/video/50342>

DOI: [doi:10.3791/50342](https://doi.org/10.3791/50342)

Keywords: Infection, Issue 73, Infectious Diseases, Molecular Biology, Cellular Biology, Immunology, Medicine, Parasitology, *Plasmodium falciparum*, Cell Culture Techniques, Hemozoin, Magnetic Beads, Schizont Purification, paramagnetism, erythrocytes, red blood cells, malaria, parasitemia, parasites, isolation, cell culture

Date Published: 3/2/2013

Citation: Coronado, L.M., Tayler, N.M., Correa, R., Giovani, R.M., Spadafora, C. Separation of *Plasmodium falciparum* Late Stage-infected Erythrocytes by Magnetic Means. *J. Vis. Exp.* (73), e50342, doi:10.3791/50342 (2013).

Abstract

Unlike other *Plasmodium* species, *P. falciparum* can be cultured in the lab, which facilitates its study¹. While the parasitemia achieved can reach the ~40% limit, the investigator usually keeps the percentage at around 10%. In many cases it is necessary to isolate the parasite-containing red blood cells (RBCs) from the uninfected ones, to enrich the culture and proceed with a given experiment.

When *P. falciparum* infects the erythrocyte, the parasite degrades and feeds from haemoglobin^{2,3}. However, the parasite must deal with a very toxic iron-containing haem moiety^{4,5}. The parasite eludes its toxicity by transforming the haem into an inert crystal polymer called haemozoin^{6,7}. This iron-containing molecule is stored in its food vacuole and the metal in it has an oxidative state which differs from the one in haem⁸. The ferric state of iron in the haemozoin confers on it a paramagnetic property absent in uninfected erythrocytes. As the invading parasite reaches maturity, the content of haemozoin also increases⁹, which bestows even more paramagnetism on the latest stages of *P. falciparum* inside the erythrocyte.

Based on this paramagnetic property, the latest stages of *P. falciparum* infected-red blood cells can be separated by passing the culture through a column containing magnetic beads. These beads become magnetic when the columns containing them are placed on a magnet holder. Infected RBCs, due to their paramagnetism, will then be trapped inside the column, while the flow-through will contain, for the most part, uninfected erythrocytes and those containing early stages of the parasite.

Here, we describe the methodology to enrich the population of late stage parasites with magnetic columns, which maintains good parasite viability¹⁰. After performing this procedure, the unattached culture can be returned to an incubator to allow the remaining parasites to continue growing.

Video Link

The video component of this article can be found at <https://www.jove.com/video/50342/>

Protocol

All steps of the protocol, except for centrifugations, should be carried out inside a hood to keep the sample sterile.

1. Late Stage Isolation of *P. falciparum*-infected Erythrocytes

All late stages of *Plasmodium*-infected erythrocytes can be separated with this methodology, since hemozoin, which confers paramagnetism on the parasite, is a common metabolite to the genus. A high parasitemia (3-10%) in culture is recommended to get better yields with this protocol, although a typical culture will contain 2-4% hematocrit (volume percentage of red blood cells) and 1-8% parasitemia (percentage of infected red blood cells). Parasites can be cultured as described by Trager and Jansen¹.

1. A simple, multi-part setup must be assembled for every schizont isolation procedure. For this, LS columns, a Magnetic MidiMACS Separator and a MACS MultiStand from Miltenyi BioTec (**Figure 1**) are used. First, attach the magnetic separator to the metal stand and then place the magnetic column on the separator.
2. In a separate tube (use 50 ml conicals throughout), mix 23.2 ml of RPMI 1640 and 750 µl of 7.5% sodium bicarbonate to create the work solution. Have additional tubes on hand for discarding and collecting the flow-through of the parasite culture.
3. Add 3 ml of the work solution to the column to equilibrate it, and discard the flow-through into a tube.

4. Add 8 ml of the *P. falciparum* culture to the column, making sure that as soon as the last part of the work solution has been pushed out of the column by the culture, the receiving tube is replaced by the collecting one to begin recovery of the unbound culture. This is necessary to avoid diluting the culture any further.
5. Once the flow of the culture through the column has ceased, add 1 ml of the work solution to push the remaining RBCs out of the column.
6. Washing step: Add another 3 ml of the work solution to the column and collect the flowing liquid into the "discard" tube in order to avoid diluting the culture being saved as flow-through in the other tube.
7. Elution: Once the 3 ml have passed through the column, detach the latter from the stand and place it on top of a 15 ml tube.
8. Again, add 3 ml of work solution to the column. This step elutes the erythrocytes infected with late stage parasites that were trapped in the beads of the column.

Note: At this point, and depending on the amount of parasites required, start another round/s of collection, obviously subject to the limits of the parasitemia in the original culture. The column can be reused as long as the flow maintains a steady state.

9. Concentration of the collected parasites: Centrifuge the 15 ml tube containing the eluate for 5 min at 150 x g at room temperature to form a dark pellet from the parasitized erythrocytes.
10. Remove the supernatant leaving enough liquid to take a very small sample and determine the concentration and total yield through microscopy (See step 2 below).
11. At this moment, if the isolation and capture of late stage parasites is accomplished satisfactorily, incubate the collected culture again with fresh media.
12. Dilute the collected parasites in the volume and solution needed for subsequent experiments.

2. Purity and Yield Analysis

1. Verify the concentration of parasites obtained by using a hemocytometer and counting under a light microscope. To count, add 10 μ l of the mix of the collected parasites and the media under the cover slip of the hemocytometer. Count the number of parasites in the four quadrants of the hemocytometer and divide the total number by 4. Multiply that number by 10^4 to obtain the concentration of infected erythrocytes per ml.

Notes: a) The concentration of the mix will be decided by the user at the moment of resuspending the pellet, by adding more or less medium. A suggested working concentration is 5.5×10^4 schizontes per μ l (this is typically achieved by adding 100-300 μ l to the collected parasites). If 20 μ l of this concentration is added to a final volume of 100 μ l mix of media and erythrocytes, a parasitemia of about 1% in a typical 4% hematocrit culture will be obtained. b) Counting under the hemocytometer has to be quick, since the erythrocytes start lysing as the sample dries inside the chamber. Infected erythrocytes, which should be the majority, will show a dark spot inside, which differentiates them from non-infected ones.

2. Perform a thin-layer Giemsa stain to assess the purity of the collection by fixing the slides very rapidly in 10% methanol, drying them, and immersing them in a 20% solution of Giemsa stain diluted with distilled water. Stain the slides for 10 min, after which they can be dried by air and examined under the microscope. A parasitemia well above 70% should be achievable.

Representative Results

In **Figure 2**, the culture passing through the magnetic column is shown, before (**A**) and after the procedure (**B**). One to two infected erythrocytes are usually seen on a 100X magnification field as shown in **Figure 2**, with the arrows pointing to infected erythrocytes in **Figure 2A**. In a typical procedure, starting with a culture at 5% parasitemia (**Figure 2A**), the performance of this procedure usually produces erythrocytes with a parasitemia of 97%-100% (**Figure 2B**). Enrichment of parasitemia on the order of 20X is easily achievable, and several passes of the culture through the column may achieve a high yield of *P. falciparum* late-stage-infected erythrocytes, as needed or desired.




Figure 1. Typical layout of the magnetic components. The MIDI magnetic column, the magnetic separator and the stand are put together as depicted here.




Figure 2. A) Typical 5% unsynchronized parasite content before enrichment. A 1 μ l sample of the parasites collected through this magnetic procedure was Giemsa stained and viewed at 100X magnification. One to three infected erythrocytes are usually seen in any given field of scope (marked with arrows). **B)** Parasite content after enrichment. A 1 μ l sample of the collected parasites through this magnetic procedure is Giemsa stained and viewed at 100X magnification. Parasitemia is now 99%.

Discussion

In vitro cultures of the malaria parasite *P. falciparum* exhibit a limited parasitemia, with more than half of the red blood cells uninfected at the highest proliferation point of culture. For most research experiments, it is desirable to work only with the infected erythrocytes. To this end, a separation technique is necessary to divide the culture according to infection. Useful methods include the use of streptolysin O to permeabilize and lyse the uninfected RBCs¹¹ and a series of variations of differential centrifugations which take advantage of the different density of the two groups. Some of these methods include gelatin flotation¹² and the use of Plasmion¹³ but the most common substance used to create layers of various densities to trap the different red blood cells is the density gradient medium Percoll, which may present some toxicity to the parasites¹⁴.

We have used a technique first described by Paul and further analyzed by Trang^{15, 16}, based on the paramagnetic property of haemozoin, an inert crystal produced by the digestion of haemoglobin by the parasite. As reported elsewhere by our group, this method is non-invasive and does not seem to affect the parasite, as indicated by the normal after-treatment capability of invasion of fresh erythrocytes, as compared to the invasion rates of parasites subjected to enrichment by Percoll gradient centrifugation^{10, 14, 17}. In addition to this benefit, if the culture has a high number of infected erythrocytes, there can be more than one collection step, since the binding capacity of the column may be not enough to hold all the parasite content at once. Therefore, the collected culture can be passed through the column several times until enough parasites are collected for a given subsequent experiment.

The collection of *P. falciparum* parasites with the use of magnetic beads is simple, fast, and requires only a benchtop centrifuge, unlike the differential centrifugation techniques that need a large and costly floor centrifuge and which also requires meticulous handling of the samples.

Some applications of culture enrichment are the use of the parasites in invasion assays where the starting parasitemia is rigorously controlled in all samples; the use of the collected parasites to extract hemozoin, the inert crystal common to the Plasmodium family; the use of a high density of parasites to prepare for microscopy analysis; and the extraction of DNA of only parasites when they are mixed with other blood cells. This last application could be used to analyze clinical samples to genetically identify the strain of malaria they are infected with and make an informed therapeutic plan.

Disclosures

We have nothing to disclose.

Acknowledgements

This work was funded by grant PRB-009 to CS and a doctoral scholarship to LC, from the Secretaría Nacional de Ciencia y Tecnología (SENACYT), Panama.

References

1. Jensen, J.B. & Trager, W. *Plasmodium falciparum* in culture: use of outdated erythrocytes and description of the candle jar method. *J. Parasitology*. **63** (5), 883-886 (1977).
2. Guzman, I.Y., Francis S.E., Oksman, A., Smith, C.E., Duffin, K.L., & Goldberg, D.E. Order and specificity of the *Plasmodium falciparum* hemoglobin degradation pathway. *J. Clin. Invest.* **93**, 1602-1608 (1994).
3. Rosenthal, P.J. & Meshnick, S.R. Hemoglobin catabolism and iron utilization by malaria parasites. *Mol. Biochem. Parasitol.* **83** (2), 131-139 (1996).
4. Fitch, C.D., Chevli, R., Kanjanangulpan, P., Dutta, P., Chevli, K. & Chou, A.C. Intracellular ferriprotoporphyrin IX is a lytic agent. *Blood*. **62** (6), 1165-1168 (1983).
5. Hebbel, R.P. & Eaton, J.W. Pathobiology of heme interaction with the erythrocyte membrane. *Semin. Hematol.* **26** (2), 136-149 (1989).
6. Egan, T.J. Haemozoin formation. *Mol. Biochem. Parasitol.* **157** (2), 127-136 (2008).
7. Hempelmann, E. & Marques, H.M. Analysis of malaria pigment from *Plasmodium falciparum*. *J. Pharmacol. Toxicol. Methods*. **32** (1), 25-30 (1994).
8. Fitch, C.D. & Kanjanangulpan, P. The state of ferriprotoporphyrin IX in malaria pigment. *J. Biol. Chem.* **262** (32), 15552-15555 (1987).
9. Moore, L.R., Fujioka, H., Williams, P.S., Chalmers, J.J., Grimberg, B., Zimmerman, P.A., & Zborowski, M. Hemoglobin degradation in malaria-infected erythrocytes determined from live cell magnetophoresis. *FASEB J.* **20** (6), 747-9 (2006).
10. Spadafora, C., Gerena, L., & Kopydlowski, K.M. Comparison of the *in vitro* invasive capabilities of *Plasmodium falciparum* schizonts isolated by Percoll gradient or using magnetic based separation. *Malaria J.* **10**, 96 (2011).
11. Jackson, K.E., Spielmann, T., Hanssen, E., Adisa, A., Separovic, F., Dixon, M.W., Trenholme, K.R., Hawthorne, P.L., Gardiner, D.L., Gilberger, T., & Tilley, L. Selective permeabilization of the host cell membrane of *Plasmodium falciparum*-infected red blood cells with streptolysin O and equinatoxin II. *Biochem. J.* **403**, 167-175 (2007).
12. Goodyer, I.D., Johnson, J., Eisenthal, R., & Hayes, D.J. Purification of mature-stage *Plasmodium falciparum* by gelatine flotation. *Ann. Trop. Med. Parasitol.* **88**(2), 209-211 (1994).
13. Pasvol, G., Wilson, R.J., Smalley, M.E., & Brown, J. Separation of viable schizont-infected red cells of *Plasmodium falciparum* from human blood. *Ann. Trop. Med. Parasitol.* **72**, 87-88 (1978).
14. Pertoft, H. Fractionation of cells and subcellular particles with Percoll. *J. Biochem. Biophys. Methods*. **10**, 44(1-2), 1-30 (2000).
15. Paul, F., Roath, S., Melville, D., Warhurst, D.C., & Osisanya, J.O.S. Separation of malaria-infected erythrocytes from whole blood: use of a selective high-gradient magnetic separation technique. *The Lancet*. **318**, 8237, 70-71 (1981).
16. Trang, D.T., Huy, N.T., Kariu, T., Tajima, K., & Kamei, K. One-step concentration of malarial parasite-infected red blood cells and removal of contaminating white blood cells. *Malar. J.* **3**, 7 (2004).
17. Nillni, E.A., Londner, M.V., & Spira, D.T. A simple method for separation of uninfected erythrocytes from those infected with *Plasmodium berghei* and for isolation of artificially released parasites. *Z. Parasitenkd.* **64**, 279-284 (1981).