

Materials List for:

## Dissection of Organizer and Animal Pole Explants from Xenopus laevis Embryos and Assembly of a Cell Adhesion Assay

Souichi Ogata<sup>1</sup>, Ken W.Y. Cho<sup>1</sup>

<sup>1</sup>Department of Developmental and Cell Biology, University of California, Irvine (UCI)

Correspondence to: Ken W.Y. Cho at kwcho@uci.edu

URL: http://www.jove.com/video/575

DOI: doi:10.3791/575

## **Materials**

Name	Туре	Company	Catalog Number	Comments
1x MBS-H	Buffer			Modified Barth's medium: 1L recipe5.14gNaCl75mgKCl0.2gN free-acidAdjust the pH to 7.4 with NaOH. Autoclave.
CMF-MBS	Buffer			This is Ca2+/Mg2+-free version of MBS-H. 500ml recipe :2.57gNaCl37.5mgKCl0.free-acidAdjust the pH to 7.4 with NaOH. Autoclave.
1% agarose/1x Barth's plates				as many as the number of samples
1% agarose/CMF-MBS plates				as many as the number of samples
E-Cadherin/Fc chimera	Reagent	Sigma-Aldrich	E-2278	Dissolve 0.1mg of solid E-cadherin in 100ul of 1x MBS-H containing 40% glycerol to make 100x master stock (1.0mg/ml). Store it in –20C. Make 1/100 dilution with 1x MBS- H to make 10ug/ml working solution just before you use. You need 200ul per samples of this working solution.
Fibronectin, 0.1% solution	Reagent	Sigma-Aldrich	F-1141	This pre-made solution can be used as 50x stock. Make 1/50 dilution with 1x MBS-H make 20ug/ ml working solution just before you use. You need 200ul per samples of this working solution.I do NOT recommend using solid Fibronectin, because it is difficult to dissolve at 0.1% concentration to make the stock solution. Although it is more expensive, this pre-made 0.1% solution is



				better to obtain consistent results.
4% BSA /MBS-H	Buffer			BSA: molecular biology grade fraction V
Sigmacoted yellow tips	Tool			
No.1 cover slips				40x24mm or 60x24mm
Press-to-seal silicon insulator	Tool	Glace Bio-Labs	JTR20-2.5	2.5mm depth x 20mm diameter
NUNC Lab-Tek Chambered Coverglass	Tool	Fisher Scientific	12-565-471	2wells/slides
Grid slide glass	Tool			or Finder graticule for cell counting